

Anemia: Common Complications and Conditions

CCC: CCC-001 (ISC)

[Link to Codes](#)**MCG Health**Inpatient & Surgical Care
30th Edition

- Prevention
- Clinical Indications for Ongoing Inpatient Care
- Alternatives to Inpatient Care
- Discharge
- References
- Footnotes
- Definitions
- Codes

Prevention

- Prevention issues and possible preventive measures include:
 - Avoid excessive blood drawing, and use pediatric or small-volume blood collection tubes.(1)(2)(3)
 - Use closed blood sampling devices in ICU patient.(1)(2)
 - Use peptic ulcer disease prophylaxis in critically ill, mechanically ventilated, and coagulopathic patient.(4)(5)
- ☐ Monitor patient receiving medications that contribute to anemia; examples include(6)(7):
 - Immunosuppressive agents
 - Chemotherapy(3)(8)(9)
 - Antibiotics, including penicillins, cephalosporins, and sulfonamides(10)
 - Antiretroviral (HIV) medications (eg, zidovudine)
 - Antiviral medications (eg, ribavirin)
 - Medications that reduce platelet function (eg, aspirin, NSAIDs)
- ☐ Monitor and treat (eg, with erythropoietin-stimulating agents or intravenous iron, or by treating underlying disease) patient who has illness or condition associated with anemia; examples include(11)(12)(13):
 - Renal failure(14)(15)(16)
 - Heart failure(17)(18)(19)(20)(21)
 - Cancer(3)(8)
 - HIV disease(22)
 - Malnutrition(23)
 - Chronic inflammatory disorders (eg, autoimmune diseases)(23)(24)(25)(26)
 - Immunodeficiency disorders (eg, common variable immunodeficiency, autoimmune lymphoproliferative syndrome)(22)
 - Critical illness (eg, sepsis, myocardial infarction, acute ischemic stroke)(2)(27)(28)(29)
 - Pregnancy(30)

- Menorrhagia(31)(32)
- ☐ Perioperative blood conservation techniques (eg, intraoperative salvage of erythrocytes, tranexamic acid to reduce bleeding risk) and treatment of anemia (eg, with erythropoietin-stimulating agents or intravenous iron) in patient at risk for postoperative anemia; examples include(11)(33)(34)(35)(36)(37)(38):
 - Orthopedic surgery (eg, total joint replacement, open reduction and internal fixation of fractures)(35)(37)(39)(40)(41)(42)(43)(44)
 - Cardiac surgery(29)(36)(45)(46)
 - Vascular surgery(47)(48)(49)(50)
 - Thoracic surgery(12)
 - Colorectal surgery(51)
- Intraoperative treatment of anemia with transfusion(46)(52)
- Preoperative treatment of anemia prior to major elective surgery(34)(45)(52)(53)

Clinical Indications for Ongoing Inpatient Care

- Ongoing inpatient care is indicated for **1 or more** of the following(14)(52)(54)(55)(56):
 - Hemodynamic instability (1)(54)(55)(57)
 - Active hemolysis (eg, progressive anemia)[A](1)(55)(57)(58)(59)(60)
 - Active bleeding that cannot be controlled in observation care(1)(29)(55)(57)
 - Altered mental status (1)(54)(57)
 - Cardiac arrhythmias of immediate concern (1)(57)
 - Acute peripheral ischemia (eg, pulseless, cool, mottled, or cyanotic extremity)(57)
 - High-risk thrombocytopenia (low platelet count) (54)(57)(60)
 - Syncope or near syncope(1)(57)
 - Clinically significant signs or symptoms (eg, heart failure, chest pain, dyspnea, myocardial ischemia, Acute renal failure (stage 3 acute kidney injury), intestinal ischemia)(57)(61)

Alternatives to Inpatient Care

- Alternatives to inpatient stay extension include:
 - Outpatient care
 - Initial evaluation and laboratory testing(14)(54)(57)
 - Further outpatient evaluation as indicated; examples include:
 - Colonoscopy(62)
 - Double contrast barium enema
 - Upper GI endoscopy(62)
 - MRI or CT enterography(63)
 - Enteroscopy or wireless capsule endoscopy
 - Radionuclide red blood cell scan
 - Angiography
 - Bone marrow aspiration and biopsy(64)
 - Outpatient treatment as indicated; examples include(3)(11)(16)(26)(55):
 - Iron, folate, or vitamin B12 supplementation as appropriate[B](3)(14)(16)
 - Intravenous iron in outpatient infusion center for iron deficiency(14)(16)(65)
 - Erythropoietin-stimulating agents (ie, recombinant human erythropoietin, darbepoetin)(3)(14)

- Transfusion in outpatient infusion center for chronic anemia (eg, for hematocrit less than 25% (0.25) accompanied by fatigue)(14)(45)(57)(66)
- Palliative care(67)

Discharge

- Extended stay beyond goal length of stay for primary condition may be needed until **ALL** of the following are present(52)(54):
 - Hemodynamic stability
 - Active blood loss absent
 - Signs and symptoms of anemia absent or improved
 - Mental status normal or at baseline
 - Hgb/Hct level stable and acceptable for next level of care
 - Etiology of anemia requiring inpatient care absent

References

1. Hof L, Choorapoikayil S, Meybohm P, Zacharowski K. Patient Blood Management in intensive care patients. *Current Opinion in Critical Care* 2021;27(6):709-716. DOI: 10.1097/MCC.0000000000000880. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9] View abstract...
2. Whitehead NS, et al. Interventions to prevent iatrogenic anemia: a Laboratory Medicine Best Practices systematic review. *Critical Care* 2019;23(1):278. DOI: 10.1186/s13054-019-2511-9. [Context Link 1, 2, 3] View abstract...
3. Griffiths EA, et al. Hematopoietic Growth Factors. NCCN Clinical Practice Guidelines in Oncology [Internet] National Comprehensive Cancer Network (NCCN). v. 1.2025; 2024 Oct 11 Accessed at: <https://www.nccn.org/>. [accessed 2024 Oct 15] [Context Link 1, 2, 3, 4, 5, 6]
4. Ye Z, et al. Gastrointestinal bleeding prophylaxis for critically ill patients: a clinical practice guideline. *British Medical Journal* 2020;368:l6722. DOI: 10.1136/bmj.l6722. [Context Link 1] View abstract...
5. PEPTIC Investigators for the Australian and New Zealand Intensive Care Society Clinical Trials Group, et al. Effect of stress ulcer prophylaxis with proton pump inhibitors vs histamine-2 receptor blockers on in-hospital mortality among ICU patients receiving invasive mechanical ventilation: the PEPTIC randomized clinical trial. *Journal of the American Medical Association* 2020;323(7):616-626. DOI: 10.1001/jama.2019.22190. [Context Link 1] View abstract...
6. Abu-Zeinah G, DeSancho MT. Understanding sideroblastic anemia: an overview of genetics, epidemiology, pathophysiology and current therapeutic options. *Journal of Blood Medicine* 2020;11:305-318. DOI: 10.2147/JBM.S232644. [Context Link 1] View abstract...
7. Thomas MR, Scully M. How I treat microangiopathic hemolytic anemia in patients with cancer. *Blood* 2021;137(10):1310-1317. DOI: 10.1182/blood.2019003810. [Context Link 1] View abstract...
8. Abdel-Razeq H, Hashem H. Recent update in the pathogenesis and treatment of chemotherapy and cancer induced anemia. *Critical Reviews in Oncology/Hematology* 2020;145:102837. DOI: 10.1016/j.critrevonc.2019.102837. [Context Link 1, 2] View abstract...
9. Gilreath JA, Rodgers GM. How I treat cancer-associated anemia. *Blood* 2020;136(7):801-813. DOI: 10.1182/blood.2019004017. [Context Link 1] View abstract...
10. Berentsen S, Barcellini W. Autoimmune hemolytic anemias. *New England Journal of Medicine* 2021;385(15):1407-1419. DOI: 10.1056/NEJMra2033982. [Context Link 1] View abstract...
11. Pan K, Pang S, Robinson M, Goede D, Meenrajan S. A review of perioperative anemia: A modifiable and not so benign risk factor. *Journal of Family Medicine and Primary Care* 2022;11(9):5004-5009. DOI: 10.4103/jfmpc.jfmpc_2209_21. [Context Link 1, 2, 3] View abstract...
12. Lanigan M, Wilkey A. Current concepts in evaluation and management of preoperative anaemia in patients undergoing thoracic surgery. *Current Opinion in Anaesthesiology* 2023;36(1):89-95. DOI: 10.1097/ACO.0000000000001214. [Context Link 1, 2] View abstract...
13. DeLoughery TG, Jackson CS, Ko CW, Rockey DC. AGA Clinical practice update on management of Iron deficiency anemia: expert review. *Clinical Gastroenterology and Hepatology* 2024;22(8):1575-1583. DOI: 10.1016/j.cgh.2024.03.046. [Context Link 1] View abstract...

14. Kidney Disease: Improving Global Outcomes (KDIGO) Anemia Work Group. KDIGO clinical practice guideline for anemia in chronic kidney disease. *Kidney International. Supplement* 2012;2(4):297-335. (Reaffirmed 2025 Jul) [Context Link 1, 2, 3, 4, 5, 6, 7]
15. Fishbane S, Coyne DW. How I treat renal anemia. *Blood* 2020;136(7):783-789. DOI: 10.1182/blood.2019004330. [Context Link 1] View abstract...
16. Bazeley JW, Wish JB. Recent and Emerging Therapies for Iron Deficiency in Anemia of CKD: A Review. *American Journal of Kidney Diseases* 2022;79(6):868-876. DOI: 10.1053/j.ajkd.2021.09.017. [Context Link 1, 2, 3, 4] View abstract...
17. Anand I, Gupta P. How I treat anemia in heart failure. *Blood* 2020;136(7):790-800. DOI: 10.1182/blood.2019004004. [Context Link 1] View abstract...
18. Ponikowski P, et al. Ferric carboxymaltose for iron deficiency at discharge after acute heart failure: a multicentre, double-blind, randomised, controlled trial. *Lancet* 2020;396(10266):1895-1904. DOI: 10.1016/S0140-6736(20)32339-4. [Context Link 1] View abstract...
19. Kalra PR, et al. Intravenous ferric derisomaltose in patients with heart failure and iron deficiency in the UK (IRONMAN): an investigator-initiated, prospective, randomised, open-label, blinded-endpoint trial. *Lancet* 2022;400(10369):2199-2209. DOI: 10.1016/S0140-6736(22)02083-9. [Context Link 1] View abstract...
20. Alharbi A, et al. Outcomes and complications of heart failure with iron deficiency anemia: a nationwide analysis. *Hospital Practice* 2024;52(3):84-90. DOI: 10.1080/21548331.2024.2337615. [Context Link 1] View abstract...
21. Wang H, et al. Impact of intravenous Iron in patients with heart failure and Iron deficiency: an updated meta-analysis of randomized controlled trials. *BMC Cardiovascular Disorders* 2024;24(1):695. DOI: 10.1186/s12872-024-04368-y. [Context Link 1] View abstract...
22. Brodsky RA. Warm autoimmune hemolytic anemia. *New England Journal of Medicine* 2019;381(7):647-654. DOI: 10.1056/NEJMcp1900554. [Context Link 1, 2] View abstract...
23. Abomhya A, et al. Iron deficiency anemia: an overlooked complication of Crohn's disease. *Journal of Hematology* 2022;11(2):55-61. DOI: 10.14740/jh989. [Context Link 1, 2] View abstract...
24. Jacobson-Kelly AE, Stanek JR, Powers JM, Dotson JL, O'Brien SH. Trends in anemia, iron, therapy, and transfusion in hospitalized pediatric patients with inflammatory bowel disease. *Journal of Pediatrics* 2020;222:141-145.e1. DOI: 10.1016/j.jpeds.2020.02.071. [Context Link 1] View abstract...
25. Gordon M, et al. Interventions for treating iron deficiency anaemia in inflammatory bowel disease. *Cochrane Database of Systematic Reviews* 2021, Issue 1. Art. No.: CD013529. DOI: 10.1002/14651858.CD013529.pub2. [Context Link 1] View abstract...
26. Goyal A, et al. Anemia in children with inflammatory bowel disease: a position paper by the IBD Committee of the North American Society of Pediatric Gastroenterology, Hepatology and Nutrition. *Journal of Pediatric Gastroenterology and Nutrition* 2020;71(4):563-582. DOI: 10.1097/MPG.0000000000002885. [Context Link 1, 2, 3] View abstract...
27. Warner MA, et al. Early post-hospitalization hemoglobin recovery and clinical outcomes in survivors of critical illness: a population-based cohort study. *Journal of Intensive Care Medicine* 2022;37(8):1067-1074. DOI: 10.1177/08850666211069098. [Context Link 1] View abstract...
28. Desai A, et al. Impact of anemia on acute ischemic stroke outcomes: A systematic review of the literature. *PLoS ONE* 2023;18(1):Online. DOI: 10.1371/journal.pone.0280025. [Context Link 1] View abstract...
29. Coz Yataco AO, et al. Red blood cell transfusion in critically ill adults: an American College of Chest physicians clinical practice guideline. *Chest* 2025;167(2):477-489. DOI: 10.1016/j.chest.2024.09.016. [Context Link 1, 2, 3] View abstract...
30. Pavord S, et al. UK guidelines on the management of iron deficiency in pregnancy. *British Journal of Haematology* 2020;188(6):819-830. DOI: 10.1111/bjh.16221. [Context Link 1] View abstract...
31. Murji A, et al. Risks of preoperative anemia in women undergoing elective hysterectomy and myomectomy. *American Journal of Obstetrics and Gynecology* 2019;221(6):629.e1-629.e18. DOI: 10.1016/j.ajog.2019.07.018. [Context Link 1] View abstract...
32. Munro MG, et al. The relationship between heavy menstrual bleeding, iron deficiency, and iron deficiency anemia. *American Journal of Obstetrics and Gynecology* 2023;229(1):1-9. DOI: 10.1016/j.ajog.2023.01.017. [Context Link 1] View abstract...
33. Tibi P, et al. STS/SCA/AmSECT/SABM update to the clinical practice guidelines on patient blood management. *Annals of Thoracic Surgery* 2021;112(3):981-1004. DOI: 10.1016/j.athoracsur.2021.03.033. [Context Link 1] View abstract...
34. Warner MA, Shore-Lesserson L, Shander A, Patel SY, Perelman SI, Guinn NR. Perioperative anemia: prevention, diagnosis, and management throughout the spectrum of perioperative care. *Anesthesia and Analgesia* 2020;130(5):1364-1380. DOI: 10.1213/ANE.0000000000004727. [Context Link 1, 2] View abstract...

35. Chaudhry YP, et al. Predictors and outcomes of postoperative hemoglobin of <8 g/dL in total joint arthroplasty. *Journal of Bone and Joint Surgery. American Volume* 2022;104(2):166-171. DOI: 10.2106/JBJS.20.01766. [Context Link 1, 2] View abstract...
36. Corwin HL, et al. Management of perioperative iron deficiency in cardiac surgery: a modified RAND Delphi study. *Annals of Thoracic Surgery* 2022;113(1):316-323. DOI: 10.1016/j.athoracsur.2020.11.031. [Context Link 1, 2] View abstract...
37. Polisetty T, Cannon D, Grewal G, Vakharia R, Vegas A, Levy JC. Iron deficiency anemia is associated with increased medical and implant-related complications, and length of stay for patients undergoing total shoulder arthroplasty. *Journal of Shoulder and Elbow Surgery* 2023;32(5):e200-e205. DOI: 10.1016/j.jse.2022.11.005. [Context Link 1, 2] View abstract...
38. Fox TP, et al. Association between routine cell salvage use for lower segment caesarean section and post-operative iron infusion and anemia. *Archives of Gynecology and Obstetrics* 2023;1-7. DOI: 10.1007/s00404-023-07082-w. [Context Link 1] View abstract...
39. Harris A, Badin D, Hegde V, Oni J, Sterling R, Khanuja H. Preoperative anemia is an independent risk factor for increased complications and mortalities after total knee arthroplasty regardless of postoperative transfusions. *Journal of Arthroplasty* 2023;38(7 Suppl 2):S177-S181. DOI: 10.1016/j.arth.2023.01.042. [Context Link 1] View abstract...
40. Ortiz-Babilonia CD, et al. Anemia and its severity is associated with worse postoperative outcomes following open reduction internal fixation of ankle fractures. *Foot and Ankle International* 2022;43(12):1532-1539. DOI: 10.1177/10711007221131811. [Context Link 1] View abstract...
41. Pollock JR, et al. 30-day outcomes for total ankle arthroplasty in patients with anemia. *Foot and Ankle International* 2021;42(11):1463-1468. DOI: 10.1177/10711007211017512. [Context Link 1] View abstract...
42. Mo K, et al. Increased severity of anemia is associated with postoperative complications following a adult spinal deformity surgery. *World Neurosurgery* 2022;167:e541-e548. DOI: 10.1016/j.wneu.2022.08.045. [Context Link 1] View abstract...
43. Levitt EB, et al. Risk factors for prolonged hospital stay after femoral neck fracture. *Orthopedics* 2023;46(4):211-217. DOI: 10.3928/01477447-20230207-02. [Context Link 1] View abstract...
44. Ganta A, Linker JA, Pettit CJ, Esper GW, Egol KA, Konda SR. Chronic preinjury anemia is associated with increased risk of 1-year mortality in geriatric hip fracture patients. *Journal of the American Academy of Orthopedic Surgeons* 2024;DOI: 10.5435/JAAOS-D-24-00658. [Context Link 1] View abstract...
45. Cammalleri V, Muscoli S, Versaci F, Romeo F. Periprocedural anemia management in severe aortic stenosis patients undergoing transcatheter aortic valve implantation. *Journal of Cardiology* 2020;75(2):117-123. DOI: 10.1016/j.jcc.2019.08.016. [Context Link 1, 2, 3] View abstract...
46. Hosseini M, et al. Analysis of anemia, transfusions, and CABG outcomes in the Society of Thoracic Surgeons national database. *Annals of Thoracic Surgery* 2024;117(5):1053-1060. DOI: 10.1016/j.athoracsur.2024.01.009. [Context Link 1, 2] View abstract...
47. Johnson CE, et al. Impact of perioperative blood transfusion in anemic patients undergoing infra inguinal bypass. *Annals of Vascular Surgery* 2022;79:72-80. DOI: 10.1016/j.avsg.2021.07.014. [Context Link 1] View abstract...
48. Natour AK, et al. Impact of preoperative anemia on hospitalization, death, and overall survival in patients with peripheral artery disease undergoing endovascular therapy: a retrospective cohort study in the United States and Canada. *Journal of Endovascular Therapy* 2023;Online. DOI: 10.1177/15266028221149926. [Context Link 1] View abstract...
49. Li R, Sidawy A, Nguyen BN. Preoperative anemia is a risk factor for worse 30-day outcomes in both open and endovascular repair of non-ruptured infrarenal abdominal aortic aneurysm. *Vascular* 2024;17085381241273141. DOI: 10.1177/17085381241273141. [Context Link 1] View abstract...
50. Nisi F, et al. Prognostic Impact of Anemia and Blood Transfusions on Cardiovascular Outcomes in Patients Undergoing Vascular Surgery: A Scoping Review. *Journal of Cardiothoracic and Vascular Anesthesia* 2025;39(2):511-525. DOI: 10.1053/j.jvca.2024.10.033. [Context Link 1] View abstract...
51. Sydhom P, et al. Efficacy and safety of preoperative intravenous iron versus standard care in colorectal cancer patients with iron deficiency anemia: a systematic review and meta-analysis. *Annals of Medicine and Surgery* 2024;86(12):7105-7119. DOI: 10.1097/MS9.0000000000002727. [Context Link 1] View abstract...
52. American Society of Anesthesiologists Task Force on Perioperative Blood Management. Practice guidelines for perioperative blood management: an updated report by the American Society of Anesthesiologists Task Force on Perioperative Blood Management*. *Anesthesiology* 2015;122(2):241-275. DOI: 10.1097/ALN.0000000000000463. (Reaffirmed 2025 Jun) [Context Link 1, 2, 3, 4] View abstract...
53. Cahill CM, et al. Preoperative anemia management program reduces blood transfusion in elective cardiac surgical patients, improving outcomes and decreasing hospital length of stay. *Transfusion* 2021;61(9):2629-2636. DOI: 10.1111/trf.16564. [Context Link 1] View abstract...

54. Adamson JW, Longo DL. Anemia and polycythemia. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. *Harrison's Principles of Internal Medicine*. 21st ed. McGraw Hill Education; 2022:431-439. [Context Link 1, 2, 3, 4, 5, 6]
55. Nelson A. Anemia. In: Gershel JC, Rauch DA, editors. *Caring for the Hospitalized Child: A Handbook of Inpatient Pediatrics*. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2023:325-331. [Context Link 1, 2, 3, 4, 5, 6]
56. Vieth JT, Lane DR. Anemia. *Hematology/Oncology Clinics of North America* 2017;31(6):1045-1060. DOI: 10.1016/j.hoc.2017.08.008. [Context Link 1] View abstract...
57. Verbillion MB, Dupre AA. Anemia and polycythemia. In: Walls RM, editor. *Rosen's Emergency Medicine: Concepts and Clinical Practice*. 10th ed. Elsevier; 2023:1462-1479.e2. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12]
58. Branchford BR, Samelson-Jones BJ, Flood VH. Platelet and blood vessel disorders. In: Kliegman RM, et al., editors. *Nelson Textbook of Pediatrics*. 22nd ed. Elsevier; 2025:3049-3060. [Context Link 1, 2]
59. Luzzatto L, De Franceschi L. Hemolytic anemias. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. *Harrison's Principles of Internal Medicine*. 21st ed. McGraw Hill Education; 2022:776-791. [Context Link 1, 2]
60. Spring J, Munshi L. Hematology emergencies in critically ill adults: benign hematology. *Chest* 2022;161(5):1285-1296. DOI: 10.1016/j.chest.2021.12.650. [Context Link 1, 2] View abstract...
61. Padda J, et al. Acute anemia and myocardial infarction. *Cureus* 2021;13(8):Online. DOI: 10.7759/cureus.17096. [Context Link 1] View abstract...
62. Ko CW, et al. AGA clinical practice guidelines on the gastrointestinal evaluation of iron deficiency anemia. *Gastroenterology* 2020;159(3):1085-1094. DOI: 10.1053/j.gastro.2020.06.046. (Reaffirmed 2025 Jul) [Context Link 1, 2] View abstract...
63. ACR-SAR-SPR Practice Parameter for the Performance of Magnetic Resonance (MR) Enterography. Resolution 26 [Internet] American College of Radiology (ACR). 2020 Accessed at: <https://www.acr.org/>. [created 2015; accessed 2024 Sep 18] [Context Link 1]
64. Kulasekararaj A, et al. Guidelines for the diagnosis and management of adult aplastic anaemia: a British Society for Haematology guideline. *British Journal of Haematology* 2024;204(3):784-804. DOI: 10.1111/bjh.19236. (Reaffirmed 2024 Aug) [Context Link 1] View abstract...
65. Auerbach M, Henry D, DeLoughery TG. Intravenous ferric derisomaltose for the treatment of iron deficiency anemia. *American Journal of Hematology* 2021;96(6):727-734. DOI: 10.1002/ajh.26124. [Context Link 1] View abstract...
66. Carson JL, et al. Transfusion thresholds for guiding red blood cell transfusion. *Cochrane Database of Systematic Reviews* 2021, Issue 12. Art. No.: CD002042. DOI: 10.1002/14651858.CD002042.pub5. [Context Link 1] View abstract...
67. Neoh K, Page A, Chin-Yee N, Doree C, Bennett MI. Practice review: Evidence-based and effective management of anaemia in palliative care patients. *Palliative Medicine* 2022;36(5):783-794. DOI: 10.1177/02692163221081967. [Context Link 1] View abstract...

Footnotes

[A] Consistent with a diagnosis of active hemolysis as the etiology of anemia are laboratory findings such as elevated serum LDH and unconjugated bilirubin, reduced serum haptoglobin, and abnormal-appearing red blood cells on microscopic examination.(1)(55)(57)(58)(59) [A in Context Link 1]

[B] The addition of vitamin C to oral iron therapy may improve absorption.(26) [B in Context Link 1]

Definitions

Acute renal failure (stage 3 acute kidney injury)

- Acute[A] kidney injury (stage 3),[B] as indicated by **1 or more** of the following[C](1)(2)(3):
 - Acute[A] rise in serum creatinine to 3 times its baseline value[D] or higher
 - Acute[A] rise in serum creatinine to at least 1.5 times its baseline value[D] (increase of 50%) and serum creatinine is greater than 4 mg/dL (354 micromoles/L)

- In child younger than 18 years, estimated GFR less than 35 mL/min/1.73m² (0.58 mL/sec/1.73m²)  eGFR - Pediatric Calculator and **1 or more** of the following:
 - Acute[A] rise in serum creatinine to at least 1.5 times its baseline value[D] (increase of 50%)
 - Within past 48 hours, rise in serum creatinine greater than 0.3 mg/dL (26.5 micromoles/L)
- Decreased urine output,[E] as indicated by **ALL** of the following:
 - Assessment of urine output appropriate (eg, adequate volume status, no urinary obstruction)
 - Oligoanuria, as indicated by **1 or more** of the following:
 - Urine output less than 0.3 mL/kg/hr for 24 hours
 - Anuria (urine output less than 0.1 mL/kg/hr) for 12 hours
- Initiation of renal replacement therapy required(4)

References

1. Kidney Disease: Improving Global Outcomes (KDIGO) Acute Kidney Injury Work Group. KDIGO clinical practice guideline for acute kidney injury. Kidney International. Supplement 2012;2(1):1-138. (Reaffirmed 2025 Apr)
2. Weisbord SD, Palevsky PM. Prevention and management of acute kidney injury. In: Yu AS, Chertow GM, Luyckx VA, Marsden PA, Skorecki K, Taal MW, editors. Brenner and Rector's The Kidney. 11th ed. Philadelphia, PA: Elsevier; 2020:940-977.e15.
3. Palevsky PM, et al. KDOQI US commentary on the 2012 KDIGO clinical practice guideline for acute kidney injury. American Journal of Kidney Diseases 2013;61(5):649-72. DOI: 10.1053/j.ajkd.2013.02.349.
4. Gaudry S, Palevsky PM, Dreyfuss D. Extracorporeal kidney-replacement therapy for acute kidney injury. New England Journal of Medicine 2022;386(10):964-975. DOI: 10.1056/NEJMra2104090.

Footnotes

- A. A specialty society practice guideline defines acute kidney injury as a change in renal function that is known or presumed to have occurred over the previous 7 days.(1)
- B. For purposes of these criteria, the term acute renal failure can be viewed as stage 3 acute kidney injury in the KDIGO classification system.(1) Of note, in the ICD-10 diagnostic coding system (main coding scheme used in labeling diagnoses), the relevant codes covering the clinical entity of significant acute kidney injury use the term acute kidney failure (eg, code N17.9).
- C. In determining the stage of acute kidney injury (eg, stage 2 vs stage 3), patients should be staged according to the criteria that give them the highest stage.(1)
- D. If the baseline serum creatinine is not known, absent known or suspected chronic kidney disease, a baseline serum creatinine can be approximated by assuming an estimated GFR of 75 mL/min/1.73m² (1.25 mL/sec/1.73m²). (1)  eGFR - Adult Calculator  eGFR - Pediatric Calculator
- E. The use of urine output as an indicator of acute kidney injury is important, as change in urine output can occur before a noticeable rise in creatinine.(1)(3) It is also suggested that urine output can identify patients with acute kidney injury that may otherwise be missed.(1)(3) However, urine output as a metric is only valid under the appropriate conditions. For example, the patient must not be hypovolemic, there must not be urinary obstruction, and the collection and measurement of urinary output must be complete and accurate.(1)(3) Use of urinary output to define acute kidney injury can be inaccurate in obese patients due to the nonlinear relationship between body weight and urine output. For example, in a 100-kg (220-lb) patient, urine output of 45 mL/hr over 12 hours (adequate output) would qualify as stage 2 acute kidney injury.(1)(3)

Altered mental status

- Altered mental status (ie, different from baseline), as indicated by **1 or more** of the following(1)(2)(3)(4):

- Confusional state (eg, disorientation, difficulty following commands, deficit in attention)
- Lethargy (eg, awake or arousable, but with drowsiness; reduced awareness of self and environment)
- Obtundation (ie, arousable only with strong stimuli, lessened interest in environment, slowed responses to stimulation)
- Stupor (ie, may be arousable but patient does not return to normal baseline level of awareness)
- Coma (ie, not arousable)

References

1. Maher PJ. Confusion. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:126-131.e1.
2. Lei C, Smith C. Depressed consciousness and coma. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:117-125.e1.
3. Abraham G, Maher PJ. Delirium and dementia. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:1269-1280.e2.
4. Holler-Managan YF. Neurologic evaluation. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:3550-3560.

Cardiac arrhythmias of immediate concern

- Cardiac arrhythmias or findings of immediate concern, as indicated by **1 or more** of the following(1)(2)(3):
 - Heart rhythms that are inherently dangerous or unstable, as indicated by **1 or more** of the following(4)(5)(6):
 - Resuscitated ventricular fibrillation or cardiac arrest
 - Ventricular escape rhythm
 - Sustained ventricular tachycardia (30 seconds or more of ventricular rhythm at greater than 100 beats per minute)
 - Nonsustained ventricular tachycardia and **1 or more** of the following:
 - Suspected cardiac ischemia as cause or consequence of ventricular tachycardia
 - Acute myocarditis
 - Unstable cardiac conduction defects, as indicated by **1 or more** of the following(7)(8):
 - Type II second-degree atrioventricular block
 - Third-degree atrioventricular block
 - New-onset left bundle branch block with suspected myocardial ischemia(9)
 - Any heart rhythm and **1 or more** of the following(5)(8)(10)(11)(12)(13):
 - Continuous long-term ECG monitoring needed (eg, initiation of drug requiring monitoring beyond observation care)
 - Patient has automatic implantable cardioverter-defibrillator that is repeatedly firing, malfunctioning, or in need of immediate adjustment of settings beyond scope of observation care.
 - Heart rhythms of concern due to **1 or more** of the following(8):
 - Hypotension
 - Respiratory distress
 - Association with other significant symptoms (eg, syncope)(10)(11)

References

1. Curtis AB, Tomaselli GF. Approach to the patient with cardiac arrhythmias. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1145-1162.e7.
2. Dalal AS, Van Hare GF. Disturbances of rate and rhythm of the heart. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:2844-2859.e1.

3. Joglar JA, et al. 2023 HRS Expert Consensus Statement on the management of arrhythmias during pregnancy. Heart Rhythm 2023;20(10):Online. DOI: 10.1016/j.hrthm.2023.05.017.
4. Garan H. Ventricular arrhythmias. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:332-340.
5. Al-Khatib SM, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. Circulation 2018;138(13):e272-e391. DOI: 10.1161/CIR.0000000000000549. (Reaffirmed 2025 Jun)
6. Myerburg RJ, Lampert R. Cardiac arrest and life-threatening arrhythmias. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:312-317.
7. Miller JM, Ellenbogen KA. Therapy for cardiac arrhythmias. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1208-1244.
8. Patton KK, Olgin JE. Bradyarrhythmias and atrioventricular block. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1312-1320.
9. Tan NY, Witt CM, Oh JK, Cha YM. Left bundle branch block: current and future perspectives. Circulation. Arrhythmia and Electrophysiology 2020;13(4):e008239. DOI: 10.1161/CIRCEP.119.008239.
10. Joglar JA, et al. 2023 ACC/AHA/ACCP/HRS Guideline for the diagnosis and management of atrial fibrillation: a report of the American College of Cardiology/American Heart Association joint committee on clinical practice guidelines. Journal of the American College of Cardiology 2024;149(1):Online. DOI: 10.1016/j.jacc.2023.08.017. (Reaffirmed 2025 Mar)
11. Zimetbaum P, Goldman L. Supraventricular ectopy and tachyarrhythmias. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:322-332.
12. Santucci PA, Wilber DJ. Electrophysiologic procedures and surgery. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:340-347.
13. Pendharkar SS, et al. AHA telemetry guidelines improve telemetry utilization in the inpatient setting. American Journal of Managed Care 2020;26(11):476-481. DOI: 10.37765/ajmc.2020.88525.

Hemodynamic instability

- Hemodynamic instability, as indicated by **1 or more** of the following:
 - Vital sign abnormality not corrected by appropriate treatment, as indicated by **1 or more** of the following[A]:
 - Tachycardia that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)[A]
 - Hypotension that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)[A]
 - Orthostatic hypotension that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)[A]
 - Severe hypotension in adult patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 80 mm Hg(1)[A]
 - Shock index greater than 1.0 (shock index is the heart rate divided by systolic blood pressure)[A](2)(3)(4)
 - Mean arterial pressure[B] less than 65 mm Hg  MAP Calculator[A](1)(6)
 - IV inotropic or vasopressor medication required to maintain adequate blood pressure or perfusion(3)(6)
 - Hypoperfusion due to hypotension, as indicated by **ALL** of the following:
 - Hypotension
 - Evidence of hypoperfusion, as indicated by **1 or more** of the following:
 - Lactate of 2.0 mmol/L (18 mg/dL) or more secondary to hypotension (ie, hypoperfusion)[C](3)(6)(8)
 - Metabolic acidosis (arterial or venous[D] pH less than 7.35) not otherwise explained(3)(6)(8)

- Significant end organ dysfunction due to hypotension (eg, Altered mental status, myocardial ischemia, 2-fold or higher acute increase in serum creatinine)(3)(6)
- Severe hypotension in pediatric patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 80 mm Hg in child 6 to 17 years of age^[A](14)
 - Systolic blood pressure less than 75 mm Hg in child 6 months to 5 years of age^[A](14)
 - Shock index greater than 0.9 in child 13 to 17 years of age (shock index is the heart rate divided by systolic blood pressure)^[A](2)(15)
 - Shock index greater than 1.0 in child 7 to 12 years of age (shock index is the heart rate divided by systolic blood pressure)^[A](2)(15)
 - Shock index greater than 1.2 in child 4 to 6 years of age (shock index is the heart rate divided by systolic blood pressure)^[A](2)(15)
 - IV inotropic or vasopressor medication required to maintain adequate blood pressure or perfusion(5)(6)
 - Hypoperfusion due to hypotension, as indicated by **ALL** of the following:
 - Hypotension
 - Evidence of hypoperfusion, as indicated by **1 or more** of the following:
 - Lactate of 2.0 mmol/L (18 mg/dL) or more secondary to hypotension (ie, hypoperfusion)^[C](6)(8)
 - Metabolic acidosis (arterial or venous pH less than 7.35) not otherwise explained^[D](5)(6)(14)
 - Significant end organ dysfunction due to hypotension (eg, Altered mental status, myocardial ischemia, 2-fold or higher acute increase in serum creatinine)(5)(6)(14)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Koch E, Lovett S, Nghiem T, Riggs RA, Rech MA. Shock index in the emergency department: utility and limitations. Open Access Emergency Medicine 2019;11:179-199. DOI: 10.2147/OAEM.S178358.
3. Angus DC. Approach to the patient with shock. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:658-665.
4. Balhara KS, Hsieh YH, Hamade B, Circh R, Kelen GD, Bayram JD. Clinical metrics in emergency medicine: the shock index and the probability of hospital admission and inpatient mortality. Emergency Medicine Journal 2017;34(2):89-94. DOI: 10.1136/emmermed-2015-205532.
5. Weiss SL, Tissieres P, Peters MJ. Shock. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:600-611.
6. Singer M, et al. The Third International Consensus definitions for sepsis and septic shock (Sepsis-3). Journal of the American Medical Association 2016;315(8):801-810. DOI: 10.1001/jama.2016.0287.
7. Wardi G, Brice J, Correia M, Liu D, Self M, Tainter C. Demystifying lactate in the emergency department. Annals of Emergency Medicine 2020;75(2):287-298. DOI: 10.1016/j.annemergmed.2019.06.027.
8. Kraut JA, Madias NE. Lactic acidosis. New England Journal of Medicine 2014;371(24):2309-2319. DOI: 10.1056/NEJMr1309483.
9. Olivieri L, Chasm R. Diabetic ketoacidosis in the pediatric emergency department. Emergency Medicine Clinics of North America 2013;31(3):755-773. DOI: 10.1016/j.emc.2013.05.004.
10. Maloney GE, Glauser JM. Diabetes mellitus and disorders of glucose homeostasis. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:1543-1558.e2.
11. Arnold TD, Miller M, van Wessel KP, Evans JA, Balogh ZJ. Base deficit from the first peripheral venous sample: a surrogate for arterial base deficit in the trauma bay. Journal of Trauma 2011;71(4):793-7; discussion 797. DOI: 10.1097/TA.0b013e31822ad694.
12. Malinoski DJ, Todd SR, Slone S, Mullins RJ, Schreiber MA. Correlation of central venous and arterial blood gas measurements in mechanically ventilated trauma patients. Archives of Surgery 2005;140(11):1122-5. DOI: 10.1001/archsurg.140.11.1122.
13. Zakrisson T, McFarlan A, Wu YY, Keshet I, Nathens A. Venous and arterial base deficits: do these agree in occult shock and in the elderly? A Bland-Altman analysis. Journal of Trauma and Acute Care Surgery 2013;74(3):936-9. DOI: 10.1097/TA.0b013e318282747a.
14. Alexander PMA, et al. Cardiovascular dysfunction criteria in critically ill children: the PODIUM consensus conference. Pediatrics 2022;149(1 Suppl 1):S39-S47. DOI: 10.1542/peds.2021-052888F.

15. Ramgopal S. The shock index among children presenting to the emergency department: analysis of nationally representative sample. *Journal of Emergency Medicine* 2024;67(2):e146-e156. DOI: 10.1016/j.jemermed.2024.03.022.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. The mean arterial pressure (MAP) takes into account both SBP and DBP readings.(3)(5)
- C. There are numerous causes of an elevated lactate level. The most common are cardiogenic or hypovolemic shock, severe heart failure, severe trauma, or sepsis. However, there are also other etiologies to consider such as vigorous exercise, seizures, liver disease, or medication use (eg, metformin, beta2-agonists); therefore, interpretation of elevated lactate levels requires consideration of the clinical context (eg, well-appearing post exercise vs hypotensive). The severity and persistence of the elevation can sometimes be helpful in this differentiation. In most instances of lactic acidosis, blood pH is less than 7.35, with a serum bicarbonate level of 20 mEq/L (mmol/L) or lower. However, a coexisting respiratory or metabolic alkalosis can mask these findings.(7)(8)
- D. Analyses have concluded that venous and arterial pH measurements are highly correlated, with small differences between them that are usually not clinically significant.(9)(10)(11)(12)(13) If a mixed acid-base disorder is suspected, an arterial blood gas is indicated.(10)

Hemodynamic stability

- Hemodynamic stability, as indicated by **1 or more** of the following:
 - Hemodynamic abnormalities at baseline or acceptable for next level of care
 - Patient hemodynamically stable, as indicated by **ALL** of the following(1)(2):
 - Tachycardia absent
 - Hypotension absent
 - No evidence of inadequate perfusion (eg, no myocardial ischemia)
 - No other hemodynamic abnormalities (eg, no Orthostatic hypotension)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. *Goldman-Cecil Medicine*. 27th ed. Elsevier; 2024:32-35.
2. Ramgopal S, Sepanski RJ, Martin-Gill C. Empirically derived age-based vital signs for children in the out-of-hospital setting. *Annals of Emergency Medicine* 2023;81(4):402-412. DOI: 10.1016/j.annemergmed.2022.09.019.

High-risk thrombocytopenia (low platelet count)

- Thrombocytopenia (low platelet count) and **1 or more** of the following(1)(2)(3):
 - Severe or life-threatening bleeding (eg, intracranial, major gastrointestinal, or extensive mucosal bleeding), with any reduced platelet count
 - Platelet count less than 20,000/mm³ (20 x10⁹/L) with any active bleeding
 - Platelet count less than 10,000/mm³ (10 x10⁹/L) with minor purpura or petechiae
 - Platelet count less than 5000/mm³ (5 x10⁹/L)
 - Low platelet count with hemolytic anemia

References

1. Konkle BA. Bleeding and thrombosis. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:450-457.
2. Konkle BA. Disorders of platelets and vessel wall. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:903-910.
3. Neunert C, et al. American Society of Hematology 2019 guidelines for immune thrombocytopenia. Blood Advances 2019;3(23):3829-3866. DOI: 10.1182/bloodadvances.2019000966. (Reaffirmed 2025 Jun)

Hypotension

- Hypotension, as indicated by **1 or more** of the following:
 - Hypotension in adult patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 90 mm Hg[A][B](1)
 - Mean arterial pressure[C] less than 70 mm Hg[A][B]  MAP Calculator(1)(2)
 - Decrease in baseline systolic blood pressure of 30 mm Hg or more, with significant signs or symptoms due to lower blood pressure (eg, near syncope, syncope, chest pain)[A][B](1)
 - Hypotension in pediatric patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 110 mm Hg in child 13 to 17 years of age[A][D](3)
 - Systolic blood pressure less than 100 mm Hg in child 6 to 12 years of age[A][D](3)
 - Systolic blood pressure less than 95 mm Hg in child 3 to 5 years of age[A][D](3)
 - Systolic blood pressure less than 90 mm Hg in child 1 or 2 years of age[A][D](3)
 - Systolic blood pressure less than 80 mm Hg in infant 6 to 11 months of age[A][D](3)
 - Systolic blood pressure less than 70 mm Hg in infant 3 to 5 months of age[A][D](3)
 - Systolic blood pressure less than 65 mm Hg in infant 1 or 2 months of age[A][D](3)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Puskarich MA, Jones AE. Shock. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:34-41.e1.
3. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of blood pressure requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline blood pressure, medications, and clinical impact. For example, an elderly patient with heart failure may be prescribed medication with the intentional goal of a reduced pressure. If the patient's baseline SBP is 92 mm Hg to 96 mm Hg, absent signs or symptoms of hypotension, an emergency department SBP of 86 mm Hg should not automatically be categorized as hypotensive. This would hold for mean arterial pressure (MAP) as well. The numeric values included in this definition are

provided to allow for consistency in terms of a technical definition of the term hypotension. Whether a blood pressure below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

C. The mean arterial pressure (MAP) takes into account both SBP and DBP readings.

D. Interpretation of blood pressure requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline blood pressure, medications, and clinical impact. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term hypotension. Whether a blood pressure below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Hypotension absent

- Hypotension absent,[A] as indicated by **1 or more** of the following:
 - Hypotension absent in adult patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure greater than or equal to 90 mm Hg[A](1)
 - Mean arterial pressure[B] greater than or equal to 70 mm Hg  MAP Calculator[A](1)(2)
 - Blood pressure at patient's baseline (eg, healthy adult with low systolic blood pressure), at intentional therapeutic goal (eg, patient with heart failure), or acceptable for next level of care (eg, blood pressure stable and no significant signs or symptoms due to low blood pressure)
 - Hypotension absent in pediatric patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure greater than or equal to 110 mm Hg in child 13 to 17 years of age[A](3)
 - Systolic blood pressure greater than or equal to 100 mm Hg in child 6 to 12 years of age[A](3)
 - Systolic blood pressure greater than or equal to 95 mm Hg in child 3 to 5 years of age[A](3)
 - Systolic blood pressure greater than or equal to 90 mm Hg in child 1 or 2 years of age[A](3)
 - Systolic blood pressure greater than or equal to 80 mm Hg in infant 6 to 11 months of age[A](3)
 - Systolic blood pressure greater than or equal to 70 mm Hg in infant 3 to 5 months of age[A](3)
 - Systolic blood pressure greater than or equal to 65 mm Hg in infant 1 or 2 months of age[A](3)
 - Blood pressure at patient's baseline (eg, healthy child with low systolic blood pressure), at intentional therapeutic goal, or acceptable for next level of care (eg, blood pressure stable and no significant signs or symptoms due to low blood pressure)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Puskarich MA, Jones AE. Shock. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:34-41.e1.
3. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. The mean arterial pressure (MAP) takes into account both SBP and DBP readings.

Hypoxemia

- Hypoxemia, as indicated by **1 or more** of the following(1):
 - Patient without baseline need for supplemental oxygen with oxygen saturation (SpO₂) of less than 90% or arterial blood gas partial pressure of oxygen (PaO₂) of less than 60 mm Hg (8.0 kPa) on room air[A][B]
 - Patient without baseline need for supplemental oxygen who now requires supplemental oxygen to keep SpO₂ greater than 89% or PaO₂ greater than 59 mm Hg (7.9 kPa)[A]
 - Patient with baseline need for supplemental oxygen who now requires increased supplemental oxygen to maintain oxygenation at baseline or acceptable level

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Jamali H, et al. Racial disparity in oxygen saturation measurements by pulse oximetry: evidence and implications. Annals of the American Thoracic Society 2022;19(12):1951-1964. DOI: 10.1513/AnnalsATS.202203-270CME.
3. Rojas-Camayo J, et al. Reference values for oxygen saturation from sea level to the highest human habitation in the Andes in acclimatised persons. Thorax 2018;73(8):776-778. DOI: 10.1136/thoraxjnl-2017-210598.

Footnotes

- A. Pulse oximetry (SpO₂) may underestimate arterial saturation (SaO₂), especially in people with dark skin pigmentation, thick skin, cold body temperature, or who are wearing colored nail polish, and thus may not accurately detect hypoxemia.(2) Where appropriate, pulse oximetry should be measured after warming fingers or removing nail polish. Pulse oximetry results should be assessed within the context of the patient's clinical presentation, and performance of an arterial blood gas considered for a more accurate assessment of oxygenation if indicated.(2) Specific target values may require adjustment when care is delivered at high altitude.(3)
- B. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Orthostatic hypotension

- Orthostatic hypotension,[A][B] as indicated by **1 or more** of the following(1)(2)(3):
 - Fall in SBP of 20 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position
 - Fall in DBP of 10 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position

References

1. Shibao C, Lipsitz LA, Biaggioni I, American Society of Hypertension Writing Group. Evaluation and treatment of orthostatic hypotension. Journal of the American Society of Hypertension 2013 Jul-Aug;7(4):317-324. DOI: 10.1016/j.jash.2013.04.006.
2. Dalal AS, Van Hare GF. Syncope. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:592-599.
3. Fang JC, O'Gara PT. History and physical examination: an evidence-based approach. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:123-140.

Footnotes

- A. Concomitant measurements of the heart rate are important to measure to help diagnose subtypes of orthostatic hypotension (eg, the lack of a compensatory increase in heart rate is typical of autonomic failure and an exaggerated tachycardia may be reflective of volume depletion). However, the heart rate is not a component of the definition of orthostatic hypotension, which relies upon blood pressure alone.(1)(2)(3)
- B. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Respiratory distress

- Respiratory distress, as indicated by **ALL** of the following(1)(2)(3)(4):
 - Patient with **1 or more** of the following:
 - Dyspnea (difficulty breathing)
 - Tachypnea
 - Abnormal breathing pattern (eg, chest retractions)
 - Other evidence of difficulty breathing
 - Evidence of respiratory compromise, as indicated by **1 or more** of the following:
 - Hypoxemia
 - Altered mental status
 - Other evidence of respiratory compromise (eg, respiratory acidosis)

References

1. Braithwaite SA, Wessel AL. Dyspnea. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:194-201.e2.
2. Naureckas ET, Solway J. Disturbances of respiratory function. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2133-2139.
3. Matthay MA, Ware LB. Acute respiratory failure. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:642-652.
4. Rakoczy K. Acute respiratory failure. In: Gershel JC, Rauch DA, editors. Caring for the Hospitalized Child: A Handbook of Inpatient Pediatrics. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2023:749-754.

Tachycardia

- Tachycardia,[A][B] as indicated by **1 or more** of the following:
 - Heart rate greater than 100 beats per minute in adult[A][B](1)
 - Heart rate greater than 85 beats per minute in child 13 to 17 years of age[A][C](2)
 - Heart rate greater than 95 beats per minute in child 6 to 12 years of age[A][C](2)
 - Heart rate greater than 110 beats per minute in child 1 to 5 years of age[A][C](2)
 - Heart rate greater than 120 beats per minute in infant 3 to 11 months of age[A][C](2)
 - Heart rate greater than 150 beats per minute in infant 1 or 2 months of age[A][C](2)

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. For example, an elderly patient on a beta-blocker medication with a baseline resting heart rate of 60 beats per minute may be clinically tachycardic at a heart rate of 94 beats per minute. Likewise, a patient who is upset, in pain, or nervous in the emergency department with a heart rate of 106 beats per minute may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as chest pain or hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.
- C. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. A patient who is upset, in pain, or nervous in the emergency department with an elevated heart rate may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Tachycardia absent

- Tachycardia[A][B] absent, as indicated by **1 or more** of the following:
 - Heart rate less than or equal to 100 beats per minute in adult[A][B](1)
 - Heart rate less than or equal to 85 beats per minute in child 13 to 17 years of age[A][B](2)
 - Heart rate less than or equal to 95 beats per minute in child 6 to 12 years of age[A][B](2)
 - Heart rate less than or equal to 110 beats per minute in child 1 to 5 years of age[A][B](2)
 - Heart rate less than or equal to 120 beats per minute in infant 3 to 11 months of age[A][B](2)
 - Heart rate less than or equal to 150 beats per minute in infant 1 or 2 months of age[A][B](2)

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no

or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

- B. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. For example, an elderly patient on a beta-blocker medication with a baseline resting heart rate of 60 beats per minute may be clinically tachycardic at a heart rate of 94 beats per minute. Likewise, a patient who is upset, in pain, or nervous in the emergency department with a heart rate of 106 beats per minute may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as chest pain or hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Tachypnea

- Tachypnea,^{[A][B]} as indicated by **1 or more** of the following:
 - Respiratory rate greater than 20 breaths per minute in adult^{[A][B]}(1)
 - Respiratory rate greater than 18 breaths per minute in child 13 to 17 years of age^{[A][B]}(2)
 - Respiratory rate greater than 22 breaths per minute in child 6 to 12 years of age^{[A][B]}(2)
 - Respiratory rate greater than 25 breaths per minute in child 3 to 5 years of age^{[A][B]}(2)
 - Respiratory rate greater than 30 breaths per minute in child 1 or 2 years of age^{[A][B]}(2)
 - Respiratory rate greater than 40 breaths per minute in infant 6 to 11 months of age^{[A][B]}(2)
 - Respiratory rate greater than 45 breaths per minute in infant 3 to 5 months of age^{[A][B]}(2)
 - Respiratory rate greater than 55 breaths per minute in infant 1 or 2 months of age^{[A][B]}(2)

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of respiratory rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline respiratory rate and whether the respiratory rate is indicative of significant underlying respiratory or other pathology. A mildly increased respiratory rate (eg, 20 to 22 breaths per minute in an adult), absent other indications of serious illness (eg, hypoxemia, metabolic acidosis, decreased tidal volume, pain), may be transitory and not clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachypnea. Whether a respiratory rate above the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment. In addition, the measurement of the respiratory rate is subject to error. A medical textbook states that because there can be significant breath-to-breath variation in respiratory rate, the rate should be assessed for at least 30 seconds, preferably for a full minute.⁽¹⁾ This same textbook also states that new technologies designed to measure the respiratory rate have not proved to be clinically useful.⁽¹⁾

Codes

ICD-10 Diagnosis: D46.0, D46.1, D46.20, D46.21, D46.22, D46.4, D50.0, D50.8, D50.9, D51.0, D51.1, D51.3, D51.8, D51.9, D52.0, D52.1, D52.8, D52.9, D53.0, D53.1, D53.2, D53.8, D53.9, D55.0, D55.1, D55.21, D55.29, D55.3, D55.8, D55.9, D58.8, D58.9, D59.0, D59.10, D59.11, D59.12, D59.13, D59.19, D59.2, D59.4, D59.8, D59.9, D61.03, D61.09, D61.1, D61.2, D61.3, D61.89, D61.9, D62, D63.0, D63.1, D63.8, D64.0, D64.1, D64.2, D64.3, D64.4, D64.81, D64.89, D64.9 [Hide]

ICD-10 Procedure: 30233H1, 30233N1, 30233P1, 30243H1, 30243N1, 30243P1

MCG Health
Inpatient & Surgical Care 30th Edition
Copyright © 2026 MCG Health, LLC
All Rights Reserved

Last Update: 1/25/2026 6:33:19 AM
Build Number: 30.0.2026012500524.025256