

Pain: Common Complications and Conditions

CCC: CCC-050 (ISC)

[Link to Codes](#)**MCG Health**Inpatient & Surgical Care
30th Edition

- Prevention
- Clinical Indications for Ongoing Inpatient Care
- Alternatives to Inpatient Care
- Discharge
- References
- Footnotes
- Definitions
- Codes

Prevention

- Prevention issues and possible preventive measures include(1)(2)(3)(4)(5)(6)(7)(8)(9)(10):
 - ☐ Frequent pain assessment using validated diagnostic tool; examples include(4)(11)(12):
 - Edmonton Symptom Assessment System
 - Verbal or visual analog scale for pain
 - Memorial Pain Assessment Card
 - Face, Legs, Activity, Cry, Consolability Scale for infants and toddlers(13)
 - Geriatric Pain Measure(14)
 - Burn Specific Pain Anxiety Scale(9)
 - Screening and treatment for coexisting opioid use disorder(10)(15)(16)(17)
 - ☐ Early referral and treatment of pain; examples include(6):
 - Pain or palliative care referral
 - Enhanced Recovery After Surgery (ERAS) protocols(18)(19)(20)
 - Analgesia and relaxation techniques prior to procedures and therapies (preventive analgesia)
 - Nonopioid anesthesia planning for high-risk patient (eg, obstructive sleep apnea)(21)
 - ☐ Nonpharmacologic prevention methods; examples include(22)(23)(24)(25):
 - Music(26)(27)
 - Hypnosis(9)(28)
 - Biofeedback
 - Guided imagery
 - Virtual reality(29)(30)
 - Distraction techniques (eg, watching cartoons, solving puzzles)(26)(31)
 - Non-nutritive sucking in infants(23)

- Multimodal analgesia; examples include(13)(20)(24)(32)(33)(34)(35)(36):
 - Intravenous acetaminophen or NSAID(37)(38)
 - Regional neural blockade(39)(40)
 - Neuraxial analgesia (epidural or intrathecal)(41)(42)
 - Periarticular joint injection
 - Cryoanalgesia(43)
 - Intravenous or topical lidocaine(44)
 - Preoperative dexamethasone(45)
 - Bisphosphonates for bone pain(46)
 - Topical anesthetic or capsaicin(47)
 - Transdermal buprenorphine(48)
 - Ketamine(9)(49)
 - Gabapentin(36)
 - Transcutaneous electrical nerve stimulation(50)
 - Buprenorphine and methadone for coexisting opioid use disorder(15)(51)
 - Antidepressants(52)(53)
- ☐ Prophylaxis of common side effects of opioids
 - Prevent or ease constipation; examples include(54)(55)(56):
 - Maintain hydration.
 - Maintain activity as much as possible.
 - Laxatives: stimulants and osmotic agents (eg, senna, lactulose, bisacodyl, polyethylene glycerol)
 - Prevent or ease nausea and vomiting (eg, antiemetics, dexamethasone).(57)
 - Prevent or ease pruritus with GABA analogues (eg, gabapentin, pregabalin).(58)

Clinical Indications for Ongoing Inpatient Care

- Ongoing inpatient care is indicated for **1 or more** of the following(1)(2)(3)(4)(5)(6)(7)(8):
 - ☐ Inpatient intravenous opioid treatment required,[A] as indicated by **ALL** of the following:
 - Pain insufficiently responsive to nonopioid analgesia (eg, NSAIDs)[B](59)
 - Alternative routes (eg, oral, transdermal, or submucosal) for opioid treatment not appropriate or not sufficient(35)(60)
 - ☐ Pain control regimen for next level of care not established, as indicated by **1 or more** of the following:
 - Sufficient pain control not achieved (eg, active adjustments to medications, dose, and route are ongoing)
 - Regimen used to achieve sufficient control is not readily performable at next level of care (eg, adjustment to medications or route necessary to be feasible at next level of care).
 - Suspected new or deteriorating disease process as etiology (eg, new site of metastatic disease) requiring inpatient evaluation and treatment (eg, urgent radiation therapy)
 - Palliative procedure planned that necessitates inpatient care for assessment and monitoring needs (eg, celiac block, stent to relieve obstruction)(61)
 - ☐ Severe side effects secondary to pain treatments, as indicated by **1 or more** of the following:
 - Hypotension
 - Dehydration that is severe or persistent
 - Vomiting that is severe or persistent
 - Altered mental status that is severe or persistent  GCS Calculator

- Ileus or obstipation with inadequate gastrointestinal function to maintain nutrition(54)(56)
- Acute urinary retention(62)
- Respiratory depression requiring reversal and readjustment of opioid therapy(63)(64)
- Seizures(65)
- Cardiac arrhythmia (eg, torsades de pointes due to prolonged QT interval on methadone)(66)(67)
- Spinal hematoma with neuraxial procedures(68)

Alternatives to Inpatient Care

- Alternatives to inpatient stay extension include(3)(4)(6)(69):
 - Outpatient care(54)
 - Multimodal analgesia (48)(50)(70)
 - Cognitive behavioral therapies (eg, imagery, meditation, prayer, biofeedback)
 - Muscle relaxants(71)
 - Antidepressants(52)(53)
 - Antiseizure medications (eg, GABA agonists)
 - Corticosteroid injection
 - Botulism injection
 - Peripheral nerve block
 - Neuromodulation
 - Bisphosphonates(46)
 - Topical anesthetics (eg, lidocaine patch)
 - Transdermal buprenorphine(48)
 - Transcutaneous electrical nerve stimulation(50)
 - Radiation therapy
 - Physical therapy
 - Outpatient procedure, including cordotomy, rhizotomy, or regional anesthesia procedure(72)
 - Laxatives: stimulants and osmotic agents (eg, senna, lactulose, bisacodyl, polyethylene glycerol)(55)(56)
 - Peripherally acting mu-opioid receptor antagonists (eg, methylbuprenorphine, naloxegol, alvimopan) for severe constipation(56)
 - Secretory agent (lubiprostone) for severe constipation
 - Home care(73)
 - Physical therapy
 - Physical techniques (cold, heat, massage)
 - Multimodal analgesia
 - Cognitive behavioral therapies (eg, imagery, meditation, prayer, biofeedback)
 - Hospice care[C] as indicated(5)
 - Recovery facility
 - Frequent assessment and analgesic adjustment
 - Cognitive behavioral therapies (eg, imagery, meditation, prayer, biofeedback)
 - Multimodal analgesia
 - Physical and occupational therapy
 - Hospice care[C] as indicated

Discharge

- Extended stay beyond goal length of stay for primary condition may be needed until **ALL** of the following are present(3)(4)(8)(60)(74)(75):
 - Hemodynamic stability
 - Pain absent or adequately managed
 - Adverse effects of medication (eg, nausea, vomiting, sedation, constipation) absent or managed and appropriate for next level of care
 - Discharge pain medication[D] regimen performable at next level of care (eg, medications available, can be administered)(15)
 - Education concerning safety and appropriate use of pain medications completed and understood

References

1. Dowell D, Ragan KR, Jones CM, Baldwin GT, Chou R. CDC Clinical Practice Guideline for Prescribing Opioids for Pain - United States, 2022. MMWR - Recommendations and Reports 2022;71(3):1-95. DOI: 10.15585/mmwr.rr7103a1. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9] View abstract...
2. Miner JR, Burton JH. Pain management. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:62-79.e2. [Context Link 1, 2]
3. Peterson SJB, Weisman SJ. Pediatric pain management. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:677-700. [Context Link 1, 2, 3, 4]
4. Swarm RA, et al. Adult Cancer Pain. NCCN Clinical Practice Guidelines in Oncology [Internet] National Comprehensive Cancer Network (NCCN). v. 2.2025; 2025 May 21 Accessed at: <https://www.nccn.org/>. [accessed 2025 Jun 04] [Context Link 1, 2, 3, 4, 5, 6, 7]
5. Walling AM, et al. Palliative Care. NCCN Clinical Practice Guidelines in Oncology [Internet] National Comprehensive Cancer Network (NCCN). v.2.2025; 2025 Mar 31 Accessed at: <https://www.nccn.org/>. [accessed 2025 Apr 29] [Context Link 1, 2, 3, 4]
6. Pain Management Best Practices Inter-Agency Task Force. Pain Management Best Practices Inter-Agency Task Force Report: Updates, Gaps, Inconsistencies, and Recommendations. [Internet] U.S. Department of Health and Human Services. 2019 May Accessed at: <https://www.hhs.gov/opioids/prevention/pain-management-options/index.html>. [accessed 2025 Sep 17] [Context Link 1, 2, 3, 4, 5]
7. Finnerup NB. Nonnarcotic methods of pain management. New England Journal of Medicine 2019;380(25):2440-2448. DOI: 10.1056/NEJMr1807061. [Context Link 1, 2] View abstract...
8. Anderson WG, Liao S. Improving Pain Management for Hospitalized Patients. [Internet] Society of Hospital Medicine. Accessed at: <https://www.hospitalmedicine.org/>. [accessed 2025 Sep 17] [Context Link 1, 2, 3]
9. Romanowski KS, et al. American Burn Association Guidelines on the Management of Acute Pain in the Adult Burn Patient: A Review of the Literature, a Compilation of Expert Opinion and Next Steps. Journal of Burn Care and Research 2020;41(6):1152-1164. DOI: 10.1093/jbcr/iraa120. (Reaffirmed 2025 Jul) [Context Link 1, 2, 3, 4] View abstract...
10. Calcaterra SL, et al. Management of opioid use disorder, opioid withdrawal, and opioid overdose prevention in hospitalized adults: A systematic review of existing guidelines. Journal of Hospital Medicine 2022;17(9):679-692. DOI: 10.1002/jhm.12908. [Context Link 1, 2] View abstract...
11. Chiarotto A, Maxwell LJ, Ostelo RW, Boers M, Tugwell P, Terwee CB. Measurement properties of Visual Analogue Scale, Numeric Rating Scale and Pain Severity subscale of the Brief Pain Inventory in patients with low back pain: a systematic review. Journal of Pain 2019;20(3):245-263. DOI: 10.1016/j.jpain.2018.07.009. [Context Link 1, 2, 3] View abstract...
12. Giordano V, et al. Pain and sedation scales for neonatal and pediatric patients in a preverbal stage of development: a systematic review. JAMA Pediatrics 2019;173(12):1186-1197. DOI: 10.1001/jamapediatrics.2019.3351. [Context Link 1] View abstract...
13. Gai N, Naser B, Hanley J, Peliowski A, Hayes J, Aoyama K. A practical guide to acute pain management in children. Journal of Anesthesia 2020;34(3):421-433. DOI: 10.1007/s00540-020-02767-x. [Context Link 1, 2] View abstract...
14. Schofield P. The assessment of pain in older people: UK National Guidelines. Age and Ageing 2018;47(suppl_1):i1-i22. DOI: 10.1093/ageing/afx192. (Reaffirmed 2025 Jun) [Context Link 1] View abstract...

15. Calcaterra SL, et al. Management of opioid use disorder and associated conditions among hospitalized adults: A Consensus Statement from the Society of Hospital Medicine. *Journal of Hospital Medicine* 2022;17(9):744-756. DOI: 10.1002/jhm.12893. [Context Link 1, 2, 3] View abstract...
16. Locke T, Salisbury-Afshar E, Coyle DT. Treatment updates for pain management and opioid use disorder. *Medical Clinics of North America* 2023;107(6):1035-1046. DOI: 10.1016/j.mcna.2023.06.017. [Context Link 1] View abstract...
17. Hatten BW, et al. Critical Issues Related to Opioids in Adult Patients Presenting to the Emergency Department. [Internet] American College of Emergency Physicians. 2024 Dec Accessed at: <https://www.acep.org/>. [accessed 2025 Sep 05] [Context Link 1]
18. Kennedy GT, et al. Enhanced recovery after surgery (ERAS) protocol reduces perioperative narcotic requirement and length of stay in patients undergoing mastectomy with implant-based reconstruction. *American Journal of Surgery* 2020;220(1):147-152. DOI: 10.1016/j.amjsurg.2019.10.007. [Context Link 1] View abstract...
19. Wainwright TW, et al. Consensus statement for perioperative care in total hip replacement and total knee replacement surgery: Enhanced Recovery After Surgery (ERAS®) Society recommendations. *Acta Orthopaedica* 2020;91(1):3-19. DOI: 10.1080/17453674.2019.1683790. [Context Link 1] View abstract...
20. Anne S, et al. Clinical practice guideline: opioid prescribing for analgesia after common otolaryngology operations. *Otolaryngology Head and Neck Surgery* 2021;164(2_suppl):S1-S42. DOI: 10.1177/0194599821996297. (Reaffirmed 2025 Jun) [Context Link 1, 2] View abstract...
21. Fiore JF, et al. Opioid versus opioid-free analgesia after surgical discharge: a systematic review and meta-analysis of randomised trials. *Lancet* 2022;399(10343):2280-2293. DOI: 10.1016/S0140-6736(22)00582-7. [Context Link 1] View abstract...
22. Vekhter D, Robbins MS, Minen M, Buse DC. Efficacy and feasibility of behavioral treatments for migraine, headache, and pain in the acute care setting. *Current Pain and Headache Reports* 2020;24(10):66. DOI: 10.1007/s11916-020-00899-z. [Context Link 1] View abstract...
23. Pillai Riddell RR, et al. Non-pharmacological management of infant and young child procedural pain. *Cochrane Database of Systematic Reviews* 2023, Issue 6. Art. No.: CD006275. DOI: 10.1002/14651858.CD006275.pub4. [Context Link 1, 2] View abstract...
24. Keefer L, Hashash JG, Szigethy E, Mayer EA. AGA clinical practice update on pain management in inflammatory bowel disease: Commentary. *Gastroenterology* 2024;166(6):1182-1189. DOI: 10.1053/j.gastro.2024.03.034. [Context Link 1, 2] View abstract...
25. Salamon KS, Russell C, DeVinney D, Soprano CM. Quality improvement protocol: improving the use of nonpharmacological pain management strategies within the inpatient hospital setting. *Journal of Clinical Medicine* 2024;13(6):DOI: 10.3390/jcm13061680. [Context Link 1] View abstract...
26. Goren K, et al. The impact of music, play, and pet therapies in managing pain and anxiety in paediatric patients in hospital: a rapid systematic review. *Paediatrics Child Health* 2023;28(4):218-224. DOI: 10.1093/pch/pxad010. [Context Link 1, 2] View abstract...
27. Trigueros-Murillo A, Martinez-Calderon J, Casuso-Holgado MJ, Gonzalez-Garcia P, Heredia-Rizo AM. Effects of music-based interventions on cancer-related pain, fatigue, and distress: an overview of systematic reviews. *Supportive Care in Cancer* 2023;31(8):Online. DOI: 10.1007/s00520-023-07938-6. [Context Link 1] View abstract...
28. Geagea D, et al. Clinical hypnosis for procedural pain and distress in children: a scoping review. *Pain Medicine* 2023;24(6):661-702. DOI: 10.1093/pm/pnac186. [Context Link 1] View abstract...
29. Smith V, et al. The effectiveness of virtual reality in managing acute pain and anxiety for medical inpatients: systematic review. *Journal of Medical Internet Research* 2020;22(11):e17980. DOI: 10.2196/17980. [Context Link 1] View abstract...
30. Groninger H, Violanti D, Mete M. Virtual reality for pain management in hospitalized patients with cancer: A randomized controlled trial. *Cancer* 2024;130(14):2552-2560. DOI: 10.1002/cncr.35282. [Context Link 1] View abstract...
31. Chan E, et al. Virtual reality for pediatric needle procedural pain: two randomized clinical trials. *Journal of Pediatrics* 2019;209:160-167.e4. DOI: 10.1016/j.jpeds.2019.02.034. [Context Link 1] View abstract...
32. O'Rourke MJ, et al. Preoperative management of opioid and nonopioid analgesics: Society for Perioperative Assessment and Quality Improvement (SPAQI) consensus statement. *Mayo Clinic Proceedings* 2021;96(5):1325-1341. DOI: 10.1016/j.mayocp.2020.06.045. [Context Link 1] View abstract...
33. Hyland SJ, Wetshtein AM, Grable SJ, Jackson MP. Acute pain management pearls: a focused review for the hospital clinician. *Healthcare (Basel, Switzerland)* 2022;11(1):Online. DOI: 10.3390/healthcare11010034. [Context Link 1] View abstract...
34. Liu H, et al. Multimodal pain management and postoperative outcomes in inpatient and outpatient shoulder arthroplasties: a population-based study. *Regional Anesthesia and Pain Medicine* 2024;Online. DOI: 10.1136/rapm-2023-104984. [Context Link 1] View abstract...

35. Davis MP, et al. Opioid conversion in adults with cancer: MASCC-ASCO-AAHPM-HPNA-NICSO guideline. *Supportive Care in Cancer* 2025;33(3):243. DOI: 10.1007/s00520-025-09286-z. [Context Link 1, 2] View abstract...
36. Garelli A, et al. Implementation of a multimodal pain protocol in adult burn patients. *Burns* 2025;51(5):107515. DOI: 10.1016/j.burns.2025.107515. [Context Link 1, 2] View abstract...
37. Barrington JW, et al. Impact of intravenous acetaminophen on lengths of stay and discharge status after total knee arthroplasty. *Journal of Knee Surgery* 2019;32(1):111-116. DOI: 10.1055/s-0038-1636908. [Context Link 1] View abstract...
38. Ma CH, et al. Systemic nonsteroidal anti-inflammatories for analgesia in postoperative critical care patients: a systematic review and meta-analysis of randomized control trials. *Critical Care Explorations* 2023;5(7):Online. DOI: 10.1097/CCE.0000000000000938. [Context Link 1] View abstract...
39. Albrecht E, Chin KJ. Advances in regional anaesthesia and acute pain management: a narrative review. *Anaesthesia* 2020;75 Suppl 1:e101-e110. DOI: 10.1111/anae.14868. [Context Link 1] View abstract...
40. Garlich JM, et al. Time to block: early regional anesthesia improves pain control in geriatric hip fractures. *Journal of Bone and Joint Surgery. American Volume* 2020;102(10):866-872. DOI: 10.2106/JBJS.19.01148. [Context Link 1] View abstract...
41. Hammal F, Chiu C, Kung JY, Bradley N, Dillane D. Pain management for hospitalized patients with rib fractures: A systematic review of randomized clinical trials. *Journal of Clinical Anesthesia* 2024;92:Online. DOI: 10.1016/j.jclinane.2023.111276. [Context Link 1] View abstract...
42. Ontario Health. Intrathecal drug delivery systems for cancer pain: a health technology assessment. *Ontario Health Technology Assessment Series* 2024;24(2):1-162. [Context Link 1] View abstract...
43. Park R, Coomber M, Gilron I, Shanthanna H. Cryoanalgesia for postsurgical pain relief in adults: A systematic review and meta-analysis. *Annals of Medicine and Surgery* 2021;69:Online. DOI: 10.1016/j.amsu.2021.102689. [Context Link 1] View abstract...
44. Chong PH, Yeo ZZ. Parenteral lidocaine for complex cancer pain in the home or inpatient hospice setting: a review and synthesis of the evidence. *Journal of Palliative Medicine* 2021;24(8):1154-1160. DOI: 10.1089/jpm.2020.0622. [Context Link 1] View abstract...
45. Heesen M, et al. Effect of intravenous dexamethasone on postoperative pain after spinal anaesthesia - a systematic review with meta-analysis and trial sequential analysis. *Anaesthesia* 2019;74(8):1047-1056. DOI: 10.1111/anae.14666. [Context Link 1] View abstract...
46. Gradishar WJ, et al. Breast Cancer. NCCN Clinical Practice Guidelines in Oncology [Internet] National Comprehensive Cancer Network (NCCN). v. 4.2025; 2025 Apr 17 Accessed at: <https://www.nccn.org/>. [accessed 2025 Apr 28] [Context Link 1, 2]
47. Goodwin B, et al. Topical capsaicin for the management of painful diabetic neuropathy: a narrative systematic review. *Pain Management* 2023;13(5):309-316. DOI: 10.2217/pmt-2023-0006. [Context Link 1] View abstract...
48. Pergolizzi JV, et al. Transdermal buprenorphine for acute pain in the clinical setting: a narrative review. *Journal of Pain Research* 2021;14:871-879. DOI: 10.2147/JPR.S280572. [Context Link 1, 2, 3] View abstract...
49. Schwenk ES, et al. Consensus guidelines on the use of intravenous ketamine infusions for acute pain management from the American Society of Regional Anesthesia and Pain Medicine, the American Academy of Pain Medicine, and the American Society of Anesthesiologists. *Regional Anesthesia and Pain Medicine* 2018;43(5):456-466. DOI: 10.1097/AAP.0000000000000806. [Context Link 1] View abstract...
50. Smith TJ, Wang EJ, Loprinzi CL. Cutaneous electroanalgesia for relief of chronic and neuropathic pain. *New England Journal of Medicine* 2023;389(2):158-164. DOI: 10.1056/NEJMr2110098. [Context Link 1, 2, 3] View abstract...
51. Herscher M, Fine M, Navalurkar R, Hirt L, Wang L. Diagnosis and management of opioid use disorder in hospitalized patients. *Medical Clinics of North America* 2020;104(4):695-708. DOI: 10.1016/j.mcna.2020.03.003. [Context Link 1] View abstract...
52. Dickenson AH, Patel R, Argoff CE. Anti-depressants. In: Benzon H, editor. *Practical Management of Pain*. 6th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:743-751.e4. [Context Link 1, 2]
53. Birkinshaw H, et al. Antidepressants for pain management in adults with chronic pain: a network meta-analysis. *Cochrane Database of Systematic Reviews* 2023, Issue 5. Art. No.: CD014682. DOI: 10.1002/14651858.CD014682.pub2. [Context Link 1, 2] View abstract...

54. Crockett SD, et al. American Gastroenterological Association Institute Guideline on the medical management of opioid-induced constipation. *Gastroenterology* 2019;156(1):218-226. DOI: 10.1053/j.gastro.2018.07.016. (Reaffirmed 2025 Feb) [Context Link 1, 2, 3] View abstract...
55. Horrigan J, Bhumi S, Miller D, Jafri M, Tadros M. Management of opioid-induced constipation in older adults. *Journal of Clinical Gastroenterology* 2023;57(1):39-47. DOI: 10.1097/MCG.0000000000001801. [Context Link 1, 2] View abstract...
56. Liu JJ, Brenner DM. Opioid-related constipation. *Gastroenterology Clinics of North America* 2022;51(1):107-121. DOI: 10.1016/j.gtc.2021.10.007. [Context Link 1, 2, 3, 4] View abstract...
57. Berger MJ, et al. Antiemesis. NCCN Clinical Practice Guidelines in Oncology [Internet] National Comprehensive Cancer Network (NCCN). v. 2.2025; 2025 May 12 Accessed at: <https://www.nccn.org/>. [accessed 2025 May 19] [Context Link 1]
58. Boehike C, Joos L, Coune B, et al. Pharmacological interventions for pruritus in adult palliative care patients. *Cochrane Database of Systematic Reviews* 2024, Issue 1. Art. No.: CD008320. DOI: 10.1002/14651858. [Context Link 1] View abstract...
59. Paice JA, et al. Use of opioids for adults with pain from cancer or cancer treatment: ASCO guideline. *Journal of Clinical Oncology* 2023;41(4):914-930. DOI: 10.1200/JCO.22.02198. [Context Link 1] View abstract...
60. Herzig SJ, Mosher HJ, Calcaterra SL, Jena AB, Nuckols TK. Improving the safety of opioid use for acute noncancer pain in hospitalized adults: a consensus statement from the Society of Hospital Medicine. *Journal of Hospital Medicine* 2018;13(4):263-271. DOI: 10.12788/jhm.2980. [Context Link 1, 2] View abstract...
61. Sola RG, Pulido P. Neurosurgical treatment of pain. *Brain Sciences* 2022;12(11):Online. DOI: 10.3390/brainsci12111584. [Context Link 1] View abstract...
62. Dobrek L. Lower urinary tract disorders as adverse drug reactions-a literature review. *Pharmaceuticals (Basel, Switzerland)* 2023;16(7):Online. DOI: 10.3390/ph16071031. [Context Link 1] View abstract...
63. Grape S, El-Boghdady K, Albrecht E. Management of adverse effects of intrathecal opioids in acute pain. *Best Practice and Research. Clinical Anaesthesiology* 2023;37(2):199-207. DOI: 10.1016/j.bpa.2023.02.002. [Context Link 1] View abstract...
64. van Lemmen M, et al. Opioid overdose: limitations in naloxone reversal of respiratory depression and prevention of cardiac arrest. *Anesthesiology* 2023;139(3):342-353. DOI: 10.1097/ALN.0000000000004622. [Context Link 1] View abstract...
65. Omoto T, Asaka J, Nihei S, Kudo K. Identifying risk factors for opioid-induced neurotoxicity in cancer patients receiving oxycodone. *Supportive Care in Cancer* 2023;31(4):Online. DOI: 10.1007/s00520-023-07676-9. [Context Link 1] View abstract...
66. Tisdale JE, et al. Drug-induced arrhythmias: a scientific statement from the American Heart Association. *Circulation* 2020;142(15):e214-e233. DOI: 10.1161/CIR.0000000000000905. (Reaffirmed 2025 Jul) [Context Link 1] View abstract...
67. Krantz MJ, Palmer RB, Haigney MCP. Cardiovascular complications of opioid use: JACC state-of-the-art review. *Journal of the American College of Cardiology* 2021;77(2):205-223. DOI: 10.1016/j.jacc.2020.11.002. [Context Link 1] View abstract...
68. Pozza DH, Tavares I, Cruz CD, Fonseca S. Spinal cord injury and complications related to neuraxial anaesthesia procedures: a systematic review. *International Journal of Molecular Sciences* 2023;24(5):Online. DOI: 10.3390/ijms24054665. [Context Link 1] View abstract...
69. Davidoff K, Heneghan C, Wolfe J, Ullrich C. Pediatric palliative care. In: Kliegman RM, et al., editors. *Nelson Textbook of Pediatrics*. 22nd ed. Elsevier; 2025:58-71. [Context Link 1]
70. Supernaw RB. Simple analgesics. In: Waldman SD, editor. *Pain Management*. 2nd ed. Philadelphia, PA: Elsevier Saunders; 2011:878-883. [Context Link 1]
71. Bhullar R, Koutalians EP, Argoff CE, Dubin A. Skeletal muscle relaxants. In: Benzon H, editor. *Practical Management of Pain*. 6th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:763-768.e2. [Context Link 1]
72. Heptonstall N, Scott-Warren J, Berman R, Filippiadis D, Bell J. Role of interventional radiology in pain management in oncology patients. *Clinical Radiology* 2023;78(4):245-253. DOI: 10.1016/j.crad.2022.05.022. [Context Link 1] View abstract...
73. Shepperd S, Goncalves-Bradley DC, Straus SE, Wee B. Hospital at home: home-based end-of-life care. *Cochrane Database of Systematic Reviews* 2021, Issue 3. Art. No.: CD009231. DOI: 10.1002/14651858.CD009231.pub3. [Context Link 1] View abstract...
74. Chou R, et al. Management of postoperative pain: a clinical practice guideline from the American Pain Society, the American Society of Regional Anesthesia and Pain Medicine, and the American Society of Anesthesiologists' Committee on Regional Anesthesia, Executive Committee, and Administrative Council. *Journal of Pain* 2016;17(2):131-57. DOI:

10.1016/j.jpain.2015.12.008. [Context Link 1] View abstract...

75. Elhady D, Novy D. Pain management at the end of life and home care for the terminally ill patient. In: Benzon H, editor. Practical Management of Pain. 6th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:1076-1092.e4. [Context Link 1]

Footnotes

[A] Many validated instruments are available to quantify pain.(4)(6)(11) A visual analog or numeric rating scale graded from 0 to 10 is often used; moderate pain correlates with a score of 4 to 6 and severe pain with 7 to 10.(4) Different scales may be more appropriate for different disease processes or patient populations.(11) [A in Context Link 1, 2]

[B] An evidence-based guideline concerning the use of opioids for pain makes several recommendations.(1) These recommendations do not apply to patients with pain due to sickle cell disease or cancer, or opioid use as part of palliative or end-of-life care. The guideline states that, because nonopioid therapies are at least as effective as opioids for many common types of pain, clinicians should maximize the use of nonpharmacologic and nonopioid pharmacologic therapies and only consider opioid therapy for pain if benefits are anticipated to outweigh risks to the patient (eg, pain insufficiently controlled via nonopioid therapy).(1) The guideline also recommends that, when starting opioid therapy for pain, clinicians should prescribe immediate-release opioids instead of long-acting opioids and should prescribe the lowest effective dosage.(1) For patients already being treated with an opioid, increasing the dosage above 50 morphine milligram equivalents (MME) per day often poses greater risk than benefit in improving pain and functioning.(1) The guideline also concludes that treatment with opioids for a few days or less is often sufficient for many causes of nonsurgical acute pain, but certain circumstances (eg, severe traumatic injuries) might require the use of opioids for durations of greater than 7 days.(1) [B in Context Link 1]

[C] Hospice enrollment may be indicated for terminal illness causing severe pain.(5) [C in Context Link 1, 2]

[D] An evidence-based guideline recommends that clinicians regularly re-evaluate benefits and risks of continued opioid therapy with patients.(1) The guideline also recommends that, to minimize withdrawal signs and symptoms, patients taking opioids continuously for more than 3 days but less than 1 week should have their daily dose reduced by 50% for 2 days prior to discontinuation, and that for patients taking opioids continuously for more than 1 week, the daily dose should be reduced 20% every 2 days.(1) [D in Context Link 1]

Definitions

Acute kidney injury (stage 2)

- Acute kidney injury (stage 2) with significant clinical findings, as indicated by **ALL** of the following(1)(2)(3):
 - Acute[A] kidney injury (stage 2), as indicated by **1 or more** of the following[B]:
 - Acute[A] rise in serum creatinine between 2 and 2.9 times its baseline value[C] (increase of 3 times baseline would qualify as stage 3 acute kidney injury)
 - Decreased urine output,[D] as indicated by **ALL** of the following:
 - Assessment of urine output appropriate (eg, adequate volume status, no urinary obstruction)
 - Urine output less than 0.5 mL/kg/hr for 12 hours
 - Significant clinical findings, as indicated by **1 or more** of the following[B]:
 - Hemodynamic instability
 - Worsening clinical status despite observation care (eg, rising creatinine or urine output remains less than 0.5 mL/kg/hr)
 - Altered mental status
 - Cardiac arrhythmias of immediate concern

- Severe hypertension (SBP greater than 180 mm Hg or DBP greater than 120 mm Hg in adults or greater than the 95th percentile for age, gender, and height plus 30 mm Hg in pediatric patients)(4)(5)
- Significant respiratory findings (eg, pulmonary edema) that are severe (eg, Hypoxemia) or persist despite observation care
- Clinically significant electrolyte abnormality that is severe (eg, hyperkalemia with severe ECG findings)[E] or persists despite observation care
- Clinically significant metabolic abnormality (eg, metabolic acidosis) that is severe or persists despite observation care

References

1. Kidney Disease: Improving Global Outcomes (KDIGO) Acute Kidney Injury Work Group. KDIGO clinical practice guideline for acute kidney injury. Kidney International. Supplement 2012;2(1):1-138. (Reaffirmed 2025 Apr)
2. Weisbord SD, Palevsky PM. Prevention and management of acute kidney injury. In: Yu AS, Chertow GM, Luyckx VA, Marsden PA, Skorecki K, Taal MW, editors. Brenner and Rector's The Kidney. 11th ed. Philadelphia, PA: Elsevier; 2020:940-977.e15.
3. Palevsky PM, et al. KDOQI US commentary on the 2012 KDIGO clinical practice guideline for acute kidney injury. American Journal of Kidney Diseases 2013;61(5):649-72. DOI: 10.1053/j.ajkd.2013.02.349.
4. Jones DW, et al. 2025 AHA/ACC/AANP/AAPA/ACCP/ACPM/AGS/AMA/ASPC/NMA/PCNA/SGIM guideline for the prevention, detection, evaluation and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. Circulation 2025;DOI: 10.1161/CIR.0000000000001356.
5. Flynn JT, et al. Clinical practice guideline for screening and management of high blood pressure in children and adolescents. Pediatrics 2017;140(3):e20171904. DOI: 10.1542/peds.2017-1904. (Reaffirmed 2025 Jul)
6. Mirvis DM, Goldberger AL. Electrocardiography. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:141-174.e3.
7. Mount DB. Fluid and electrolyte disturbances. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:338-356.

Footnotes

- A. A specialty society practice guideline defines acute kidney injury as a change in renal function that is known or presumed to have occurred over the previous 7 days.(1)
- B. In determining the stage of acute kidney injury (eg, stage 2 vs stage 3), patients should be staged according to the criteria that give them the highest stage.(1)
- C. If the baseline serum creatinine is not known, absent known or suspected chronic kidney disease, a baseline serum creatinine can be approximated by assuming an estimated GFR of 75 mL/min/1.73m² (1.25 mL/sec/1.73m²). (1)  eGFR - Adult Calculator  eGFR - Pediatric Calculator
- D. The use of urine output as an indicator of acute kidney injury is important, as change in urine output can occur before a noticeable rise in creatinine.(1)(3) It is also suggested that urine output can identify patients with acute kidney injury that may otherwise be missed.(1)(3) However, urine output as a metric is only valid under the appropriate conditions. For example, the patient must not be hypovolemic, there must not be urinary obstruction, and the collection and measurement of urinary output must be complete and accurate.(1)(3) Use of urinary output to define acute kidney injury can be inaccurate in obese patients due to the nonlinear relationship between body weight and urine output. For example, in a 100-kg (220-lb) patient, urine output of 45 mL/hr over 12 hours (adequate output) would qualify as stage 2 acute kidney injury.(1)(3)
- E. ECG findings include peaked T-waves, PR interval greater than 0.20 seconds, QRS interval greater than 0.12 seconds, and arrhythmia.(6)(7)

Acute renal failure (stage 3 acute kidney injury)

- Acute^[A] kidney injury (stage 3),^[B] as indicated by **1 or more** of the following^[C](1)(2)(3):
 - Acute^[A] rise in serum creatinine to 3 times its baseline value^[D] or higher

- Acute^[A] rise in serum creatinine to at least 1.5 times its baseline value^[D] (increase of 50%) and serum creatinine is greater than 4 mg/dL (354 micromoles/L)
- In child younger than 18 years, estimated GFR less than 35 mL/min/1.73m² (0.58 mL/sec/1.73m²)  eGFR - Pediatric Calculator and **1 or more** of the following:
 - Acute^[A] rise in serum creatinine to at least 1.5 times its baseline value^[D] (increase of 50%)
 - Within past 48 hours, rise in serum creatinine greater than 0.3 mg/dL (26.5 micromoles/L)
- Decreased urine output,^[E] as indicated by **ALL** of the following:
 - Assessment of urine output appropriate (eg, adequate volume status, no urinary obstruction)
 - Oligoanuria, as indicated by **1 or more** of the following:
 - Urine output less than 0.3 mL/kg/hr for 24 hours
 - Anuria (urine output less than 0.1 mL/kg/hr) for 12 hours
- Initiation of renal replacement therapy required⁽⁴⁾

References

1. Kidney Disease: Improving Global Outcomes (KDIGO) Acute Kidney Injury Work Group. KDIGO clinical practice guideline for acute kidney injury. Kidney International. Supplement 2012;2(1):1-138. (Reaffirmed 2025 Apr)
2. Weisbord SD, Palevsky PM. Prevention and management of acute kidney injury. In: Yu AS, Chertow GM, Luyckx VA, Marsden PA, Skorecki K, Taal MW, editors. Brenner and Rector's The Kidney. 11th ed. Philadelphia, PA: Elsevier; 2020:940-977.e15.
3. Palevsky PM, et al. KDOQI US commentary on the 2012 KDIGO clinical practice guideline for acute kidney injury. American Journal of Kidney Diseases 2013;61(5):649-72. DOI: 10.1053/j.ajkd.2013.02.349.
4. Gaudry S, Palevsky PM, Dreyfuss D. Extracorporeal kidney-replacement therapy for acute kidney injury. New England Journal of Medicine 2022;386(10):964-975. DOI: 10.1056/NEJMra2104090.

Footnotes

- A. A specialty society practice guideline defines acute kidney injury as a change in renal function that is known or presumed to have occurred over the previous 7 days.⁽¹⁾
- B. For purposes of these criteria, the term acute renal failure can be viewed as stage 3 acute kidney injury in the KDIGO classification system.⁽¹⁾ Of note, in the ICD-10 diagnostic coding system (main coding scheme used in labeling diagnoses), the relevant codes covering the clinical entity of significant acute kidney injury use the term acute kidney failure (eg, code N17.9).
- C. In determining the stage of acute kidney injury (eg, stage 2 vs stage 3), patients should be staged according to the criteria that give them the highest stage.⁽¹⁾
- D. If the baseline serum creatinine is not known, absent known or suspected chronic kidney disease, a baseline serum creatinine can be approximated by assuming an estimated GFR of 75 mL/min/1.73m² (1.25 mL/sec/1.73m²).⁽¹⁾  eGFR - Adult Calculator  eGFR - Pediatric Calculator
- E. The use of urine output as an indicator of acute kidney injury is important, as change in urine output can occur before a noticeable rise in creatinine.⁽¹⁾⁽³⁾ It is also suggested that urine output can identify patients with acute kidney injury that may otherwise be missed.⁽¹⁾⁽³⁾ However, urine output as a metric is only valid under the appropriate conditions. For example, the patient must not be hypovolemic, there must not be urinary obstruction, and the collection and measurement of urinary output must be complete and accurate.⁽¹⁾⁽³⁾ Use of urinary output to define acute kidney injury can be inaccurate in obese patients due to the nonlinear relationship between body weight and urine output. For example, in a 100-kg (220-lb) patient, urine output of 45 mL/hr over 12 hours (adequate output) would qualify as stage 2 acute kidney injury.⁽¹⁾⁽³⁾

Altered mental status

- Altered mental status (ie, different from baseline), as indicated by **1 or more** of the following(1)(2)(3)(4):
 - Confusional state (eg, disorientation, difficulty following commands, deficit in attention)
 - Lethargy (eg, awake or arousable, but with drowsiness; reduced awareness of self and environment)
 - Obtundation (ie, arousable only with strong stimuli, lessened interest in environment, slowed responses to stimulation)
 - Stupor (ie, may be arousable but patient does not return to normal baseline level of awareness)
 - Coma (ie, not arousable)

References

1. Maher PJ. Confusion. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:126-131.e1.
2. Lei C, Smith C. Depressed consciousness and coma. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:117-125.e1.
3. Abraham G, Maher PJ. Delirium and dementia. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:1269-1280.e2.
4. Holler-Managan YF. Neurologic evaluation. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:3550-3560.

Altered mental status that is severe or persistent

- Altered mental status (ie, different from baseline) that is severe or persistent, as indicated by **1 or more** of the following(1)(2)(3)(4):
 - Confusional state (eg, disorientation, difficulty following commands, deficit in attention) that persists despite appropriate treatment (eg, of underlying cause, despite observation care)
 - Lethargy (eg, awake or arousable, but with drowsiness; reduced awareness of self and environment) that persists despite appropriate treatment (eg, of underlying cause, despite observation care)
 - Obtundation (ie, arousable only with strong stimuli, lessened interest in environment, slowed responses to stimulation)
 - Stupor (ie, may be arousable but patient does not return to normal baseline level of awareness)
 - Coma (ie, not arousable)

References

1. Maher PJ. Confusion. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:126-131.e1.
2. Lei C, Smith C. Depressed consciousness and coma. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:117-125.e1.
3. Abraham G, Maher PJ. Delirium and dementia. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:1269-1280.e2.
4. Holler-Managan YF. Neurologic evaluation. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:3550-3560.

Cardiac arrhythmias of immediate concern

- Cardiac arrhythmias or findings of immediate concern, as indicated by **1 or more** of the following(1)(2)(3):
 - Heart rhythms that are inherently dangerous or unstable, as indicated by **1 or more** of the following(4)(5)(6):
 - Resuscitated ventricular fibrillation or cardiac arrest
 - Ventricular escape rhythm
 - Sustained ventricular tachycardia (30 seconds or more of ventricular rhythm at greater than 100 beats per minute)
 - Nonsustained ventricular tachycardia and **1 or more** of the following:

- Suspected cardiac ischemia as cause or consequence of ventricular tachycardia
- Acute myocarditis
- Unstable cardiac conduction defects, as indicated by **1 or more** of the following(7)(8):
 - Type II second-degree atrioventricular block
 - Third-degree atrioventricular block
 - New-onset left bundle branch block with suspected myocardial ischemia(9)
- Any heart rhythm and **1 or more** of the following(5)(8)(10)(11)(12)(13):
 - Continuous long-term ECG monitoring needed (eg, initiation of drug requiring monitoring beyond observation care)
 - Patient has automatic implantable cardioverter-defibrillator that is repeatedly firing, malfunctioning, or in need of immediate adjustment of settings beyond scope of observation care.
- Heart rhythms of concern due to **1 or more** of the following(8):
 - Hypotension
 - Respiratory distress
 - Association with other significant symptoms (eg, syncope)(10)(11)

References

1. Curtis AB, Tomaselli GF. Approach to the patient with cardiac arrhythmias. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1145-1162.e7.
2. Dalal AS, Van Hare GF. Disturbances of rate and rhythm of the heart. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:2844-2859.e1.
3. Joglar JA, et al. 2023 HRS Expert Consensus Statement on the management of arrhythmias during pregnancy. Heart Rhythm 2023;20(10):Online. DOI: 10.1016/j.hrthm.2023.05.017.
4. Garan H. Ventricular arrhythmias. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:332-340.
5. Al-Khatib SM, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. Circulation 2018;138(13):e272-e391. DOI: 10.1161/CIR.0000000000000549. (Reaffirmed 2025 Jun)
6. Myerburg RJ, Lampert R. Cardiac arrest and life-threatening arrhythmias. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:312-317.
7. Miller JM, Ellenbogen KA. Therapy for cardiac arrhythmias. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1208-1244.
8. Patton KK, Olgin JE. Bradyarrhythmias and atrioventricular block. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1312-1320.
9. Tan NY, Witt CM, Oh JK, Cha YM. Left bundle branch block: current and future perspectives. Circulation. Arrhythmia and Electrophysiology 2020;13(4):e008239. DOI: 10.1161/CIRCEP.119.008239.
10. Joglar JA, et al. 2023 ACC/AHA/ACCP/HRS Guideline for the diagnosis and management of atrial fibrillation: a report of the American College of Cardiology/American Heart Association joint committee on clinical practice guidelines. Journal of the American College of Cardiology 2024;149(1):Online. DOI: 10.1016/j.jacc.2023.08.017. (Reaffirmed 2025 Mar)
11. Zimetbaum P, Goldman L. Supraventricular ectopy and tachyarrhythmias. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:322-332.
12. Santucci PA, Wilber DJ. Electrophysiologic procedures and surgery. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:340-347.
13. Pendharkar SS, et al. AHA telemetry guidelines improve telemetry utilization in the inpatient setting. American Journal of Managed Care 2020;26(11):476-481. DOI: 10.37765/ajmc.2020.88525.

Dehydration that is severe or persistent

- Dehydration and **1 or more** of the following(1)(2)(3)(4):
 - Clinical findings of severe dehydration, as indicated by **1 or more** of the following:
 - Hemodynamic instability
 - Acute renal failure (stage 3 acute kidney injury)
 - Acute kidney injury (stage 2)
 - Serum sodium greater than 150 mEq/L (mmol/L)
 - Dehydration that is persistent, as indicated by **ALL** of the following:
 - Oral rehydration therapy not tolerated or insufficient to adequately correct dehydration
 - Dehydration persists despite appropriate treatment (eg, IV fluids, observation care)

References

- Padilipsky P, White W. Pediatric infectious diarrheal disease and dehydration. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:2151-2168.e3.
- Al-Awqati Q, Radhakrishnan J. Disorders of sodium and water. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:733-744.e1.
- Greenbaum LA. Deficit therapy. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:529-532.
- Greenbaum LA. Maintenance and replacement therapy. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:525-529.

Hemodynamic instability

- Hemodynamic instability, as indicated by **1 or more** of the following:
 - Vital sign abnormality not corrected by appropriate treatment, as indicated by **1 or more** of the following[A]:
 - Tachycardia that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)[A]
 - Hypotension that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)[A]
 - Orthostatic hypotension that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)[A]
 - Severe hypotension in adult patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 80 mm Hg(1)[A]
 - Shock index greater than 1.0 (shock index is the heart rate divided by systolic blood pressure)[A](2)(3)(4)
 - Mean arterial pressure[B] less than 65 mm Hg  MAP Calculator[A](1)(6)
 - IV inotropic or vasopressor medication required to maintain adequate blood pressure or perfusion(3)(6)
 - Hypoperfusion due to hypotension, as indicated by **ALL** of the following:
 - Hypotension
 - Evidence of hypoperfusion, as indicated by **1 or more** of the following:
 - Lactate of 2.0 mmol/L (18 mg/dL) or more secondary to hypotension (ie, hypoperfusion)[C](3)(6)(8)
 - Metabolic acidosis (arterial or venous[D] pH less than 7.35) not otherwise explained(3)(6)(8)
 - Significant end organ dysfunction due to hypotension (eg, Altered mental status, myocardial ischemia, 2-fold or higher acute increase in serum creatinine)(3)(6)
 - Severe hypotension in pediatric patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 80 mm Hg in child 6 to 17 years of age[A](14)

- Systolic blood pressure less than 75 mm Hg in child 6 months to 5 years of age^{[A](14)}
- Shock index greater than 0.9 in child 13 to 17 years of age (shock index is the heart rate divided by systolic blood pressure)^{[A](2)(15)}
- Shock index greater than 1.0 in child 7 to 12 years of age (shock index is the heart rate divided by systolic blood pressure)^{[A](2)(15)}
- Shock index greater than 1.2 in child 4 to 6 years of age (shock index is the heart rate divided by systolic blood pressure)^{[A](2)(15)}
- IV inotropic or vasopressor medication required to maintain adequate blood pressure or perfusion⁽⁵⁾⁽⁶⁾
- Hypoperfusion due to hypotension, as indicated by **ALL** of the following:
 - Hypotension
 - Evidence of hypoperfusion, as indicated by **1 or more** of the following:
 - Lactate of 2.0 mmol/L (18 mg/dL) or more secondary to hypotension (ie, hypoperfusion)^{[C](6)(8)}
 - Metabolic acidosis (arterial or venous pH less than 7.35) not otherwise explained^{[D](5)(6)(14)}
 - Significant end organ dysfunction due to hypotension (eg, Altered mental status, myocardial ischemia, 2-fold or higher acute increase in serum creatinine)⁽⁵⁾⁽⁶⁾⁽¹⁴⁾

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Koch E, Lovett S, Nghiem T, Riggs RA, Rech MA. Shock index in the emergency department: utility and limitations. Open Access Emergency Medicine 2019;11:179-199. DOI: 10.2147/OAEM.S178358.
3. Angus DC. Approach to the patient with shock. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:658-665.
4. Balhara KS, Hsieh YH, Hamade B, Circh R, Kelen GD, Bayram JD. Clinical metrics in emergency medicine: the shock index and the probability of hospital admission and inpatient mortality. Emergency Medicine Journal 2017;34(2):89-94. DOI: 10.1136/emmermed-2015-205532.
5. Weiss SL, Tissieres P, Peters MJ. Shock. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:600-611.
6. Singer M, et al. The Third International Consensus definitions for sepsis and septic shock (Sepsis-3). Journal of the American Medical Association 2016;315(8):801-810. DOI: 10.1001/jama.2016.0287.
7. Wardi G, Brice J, Correia M, Liu D, Self M, Tainter C. Demystifying lactate in the emergency department. Annals of Emergency Medicine 2020;75(2):287-298. DOI: 10.1016/j.annemergmed.2019.06.027.
8. Kraut JA, Madias NE. Lactic acidosis. New England Journal of Medicine 2014;371(24):2309-2319. DOI: 10.1056/NEJMr1309483.
9. Olivieri L, Chasm R. Diabetic ketoacidosis in the pediatric emergency department. Emergency Medicine Clinics of North America 2013;31(3):755-773. DOI: 10.1016/j.emc.2013.05.004.
10. Maloney GE, Glauser JM. Diabetes mellitus and disorders of glucose homeostasis. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:1543-1558.e2.
11. Arnold TD, Miller M, van Wessem KP, Evans JA, Balogh ZJ. Base deficit from the first peripheral venous sample: a surrogate for arterial base deficit in the trauma bay. Journal of Trauma 2011;71(4):793-7; discussion 797. DOI: 10.1097/TA.0b013e31822ad694.
12. Malinoski DJ, Todd SR, Slone S, Mullins RJ, Schreiber MA. Correlation of central venous and arterial blood gas measurements in mechanically ventilated trauma patients. Archives of Surgery 2005;140(11):1122-5. DOI: 10.1001/archsurg.140.11.1122.
13. Zakrisson T, McFarlan A, Wu YY, Keshet I, Nathens A. Venous and arterial base deficits: do these agree in occult shock and in the elderly? A Bland-Altman analysis. Journal of Trauma and Acute Care Surgery 2013;74(3):936-9. DOI: 10.1097/TA.0b013e318282747a.
14. Alexander PMA, et al. Cardiovascular dysfunction criteria in critically ill children: the PODIUM consensus conference. Pediatrics 2022;149(1 Suppl 1):S39-S47. DOI: 10.1542/peds.2021-052888F.
15. Ramgopal S. The shock index among children presenting to the emergency department: analysis of nationally representative sample. Journal of Emergency Medicine 2024;67(2):e146-e156. DOI: 10.1016/j.jemermed.2024.03.022.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. The mean arterial pressure (MAP) takes into account both SBP and DBP readings.(3)(5)
- C. There are numerous causes of an elevated lactate level. The most common are cardiogenic or hypovolemic shock, severe heart failure, severe trauma, or sepsis. However, there are also other etiologies to consider such as vigorous exercise, seizures, liver disease, or medication use (eg, metformin, beta2-agonists); therefore, interpretation of elevated lactate levels requires consideration of the clinical context (eg, well-appearing post exercise vs hypotensive). The severity and persistence of the elevation can sometimes be helpful in this differentiation. In most instances of lactic acidosis, blood pH is less than 7.35, with a serum bicarbonate level of 20 mEq/L (mmol/L) or lower. However, a coexisting respiratory or metabolic alkalosis can mask these findings.(7)(8)
- D. Analyses have concluded that venous and arterial pH measurements are highly correlated, with small differences between them that are usually not clinically significant.(9)(10)(11)(12)(13) If a mixed acid-base disorder is suspected, an arterial blood gas is indicated.(10)

Hemodynamic stability

- Hemodynamic stability, as indicated by **1 or more** of the following:
 - Hemodynamic abnormalities at baseline or acceptable for next level of care
 - Patient hemodynamically stable, as indicated by **ALL** of the following(1)(2):
 - Tachycardia absent
 - Hypotension absent
 - No evidence of inadequate perfusion (eg, no myocardial ischemia)
 - No other hemodynamic abnormalities (eg, no Orthostatic hypotension)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Ramgopal S, Sepanski RJ, Martin-Gill C. Empirically derived age-based vital signs for children in the out-of-hospital setting. Annals of Emergency Medicine 2023;81(4):402-412. DOI: 10.1016/j.annemergmed.2022.09.019.

Hypotension

- Hypotension, as indicated by **1 or more** of the following:
 - Hypotension in adult patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 90 mm Hg[A][B](1)
 - Mean arterial pressure[C] less than 70 mm Hg[A][B]  MAP Calculator(1)(2)
 - Decrease in baseline systolic blood pressure of 30 mm Hg or more, with significant signs or symptoms due to lower blood pressure (eg, near syncope, syncope, chest pain)[A][B](1)
 - Hypotension in pediatric patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 110 mm Hg in child 13 to 17 years of age[A][D](3)
 - Systolic blood pressure less than 100 mm Hg in child 6 to 12 years of age[A][D](3)
 - Systolic blood pressure less than 95 mm Hg in child 3 to 5 years of age[A][D](3)

- Systolic blood pressure less than 90 mm Hg in child 1 or 2 years of age[A][D](3)
- Systolic blood pressure less than 80 mm Hg in infant 6 to 11 months of age[A][D](3)
- Systolic blood pressure less than 70 mm Hg in infant 3 to 5 months of age[A][D](3)
- Systolic blood pressure less than 65 mm Hg in infant 1 or 2 months of age[A][D](3)

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Puskarch MA, Jones AE. Shock. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:34-41.e1.
3. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of blood pressure requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline blood pressure, medications, and clinical impact. For example, an elderly patient with heart failure may be prescribed medication with the intentional goal of a reduced pressure. If the patient's baseline SBP is 92 mm Hg to 96 mm Hg, absent signs or symptoms of hypotension, an emergency department SBP of 86 mm Hg should not automatically be categorized as hypotensive. This would hold for mean arterial pressure (MAP) as well. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term hypotension. Whether a blood pressure below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.
- C. The mean arterial pressure (MAP) takes into account both SBP and DBP readings.
- D. Interpretation of blood pressure requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline blood pressure, medications, and clinical impact. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term hypotension. Whether a blood pressure below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Hypotension absent

- Hypotension absent,[A] as indicated by **1 or more** of the following:
 - Hypotension absent in adult patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure greater than or equal to 90 mm Hg[A](1)
 - Mean arterial pressure[B] greater than or equal to 70 mm Hg  MAP Calculator[A](1)(2)
 - Blood pressure at patient's baseline (eg, healthy adult with low systolic blood pressure), at intentional therapeutic goal (eg, patient with heart failure), or acceptable for next level of care (eg, blood pressure stable and no significant signs or symptoms due to low blood pressure)
 - Hypotension absent in pediatric patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure greater than or equal to 110 mm Hg in child 13 to 17 years of age[A](3)
 - Systolic blood pressure greater than or equal to 100 mm Hg in child 6 to 12 years of age[A](3)
 - Systolic blood pressure greater than or equal to 95 mm Hg in child 3 to 5 years of age[A](3)
 - Systolic blood pressure greater than or equal to 90 mm Hg in child 1 or 2 years of age[A](3)

- Systolic blood pressure greater than or equal to 80 mm Hg in infant 6 to 11 months of age[A](3)
- Systolic blood pressure greater than or equal to 70 mm Hg in infant 3 to 5 months of age[A](3)
- Systolic blood pressure greater than or equal to 65 mm Hg in infant 1 or 2 months of age[A](3)
- Blood pressure at patient's baseline (eg, healthy child with low systolic blood pressure), at intentional therapeutic goal, or acceptable for next level of care (eg, blood pressure stable and no significant signs or symptoms due to low blood pressure)

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Puskarch MA, Jones AE. Shock. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:34-41.e1.
3. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. The mean arterial pressure (MAP) takes into account both SBP and DBP readings.

Hypoxemia

- Hypoxemia, as indicated by **1 or more** of the following(1):
 - Patient without baseline need for supplemental oxygen with oxygen saturation (SpO₂) of less than 90% or arterial blood gas partial pressure of oxygen (PaO₂) of less than 60 mm Hg (8.0 kPa) on room air[A][B]
 - Patient without baseline need for supplemental oxygen who now requires supplemental oxygen to keep SpO₂ greater than 89% or PaO₂ greater than 59 mm Hg (7.9 kPa)[A]
 - Patient with baseline need for supplemental oxygen who now requires increased supplemental oxygen to maintain oxygenation at baseline or acceptable level

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Jamali H, et al. Racial disparity in oxygen saturation measurements by pulse oximetry: evidence and implications. Annals of the American Thoracic Society 2022;19(12):1951-1964. DOI: 10.1513/AnnalsATS.202203-270CME.
3. Rojas-Camayo J, et al. Reference values for oxygen saturation from sea level to the highest human habitation in the Andes in acclimatised persons. Thorax 2018;73(8):776-778. DOI: 10.1136/thoraxjnl-2017-210598.

Footnotes

- A. Pulse oximetry (SpO₂) may underestimate arterial saturation (SaO₂), especially in people with dark skin pigmentation, thick skin, cold body temperature, or who are wearing colored nail polish, and thus may not accurately detect hypoxemia.(2) Where appropriate, pulse oximetry should be measured after warming fingers or removing nail polish. Pulse oximetry results should be assessed within the context of the patient's clinical presentation, and performance of an arterial blood

gas considered for a more accurate assessment of oxygenation if indicated.(2) Specific target values may require adjustment when care is delivered at high altitude.(3)

- B. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Multimodal analgesia

- Multimodal analgesia involves the utilization of 2 or more analgesic agents with different mechanisms of action in order to provide additive or synergistic pain control, while minimizing side effects and reliance on opioids.(1)(2)(3)

References

1. D'Souza RS, Johnson RL. Regional and multimodal treatments of perioperative pain. In: Benzon H, editor. Practical Management of Pain. 6th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:355-373.e8.
2. George S, Johns M. Review of nonopioid multimodal analgesia for surgical and trauma patients. American Journal of Health-System Pharmacy 2020;77(24):2052-2063. DOI: 10.1093/ajhp/zxaa301.
3. Finnerup NB. Nonnarcotic methods of pain management. New England Journal of Medicine 2019;380(25):2440-2448. DOI: 10.1056/NEJMra1807061.

Orthostatic hypotension

- Orthostatic hypotension,[A][B] as indicated by **1 or more** of the following(1)(2)(3):
 - Fall in SBP of 20 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position
 - Fall in DBP of 10 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position

References

1. Shibao C, Lipsitz LA, Biaggioni I, American Society of Hypertension Writing Group. Evaluation and treatment of orthostatic hypotension. Journal of the American Society of Hypertension 2013 Jul-Aug;7(4):317-324. DOI: 10.1016/j.jash.2013.04.006.
2. Dalal AS, Van Hare GF. Syncope. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:592-599.
3. Fang JC, O'Gara PT. History and physical examination: an evidence-based approach. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:123-140.

Footnotes

- A. Concomitant measurements of the heart rate are important to measure to help diagnose subtypes of orthostatic hypotension (eg, the lack of a compensatory increase in heart rate is typical of autonomic failure and an exaggerated tachycardia may be reflective of volume depletion). However, the heart rate is not a component of the definition of orthostatic hypotension, which relies upon blood pressure alone.(1)(2)(3)
- B. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Respiratory distress

- Respiratory distress, as indicated by **ALL** of the following(1)(2)(3)(4):
 - Patient with **1 or more** of the following:
 - Dyspnea (difficulty breathing)
 - Tachypnea
 - Abnormal breathing pattern (eg, chest retractions)
 - Other evidence of difficulty breathing
 - Evidence of respiratory compromise, as indicated by **1 or more** of the following:
 - Hypoxemia
 - Altered mental status
 - Other evidence of respiratory compromise (eg, respiratory acidosis)

References

- Braithwaite SA, Wessel AL. Dyspnea. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:194-201.e2.
- Naureckas ET, Solway J. Disturbances of respiratory function. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2133-2139.
- Matthay MA, Ware LB. Acute respiratory failure. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:642-652.
- Rakoczy K. Acute respiratory failure. In: Gershel JC, Rauch DA, editors. Caring for the Hospitalized Child: A Handbook of Inpatient Pediatrics. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2023:749-754.

Tachycardia

- Tachycardia,[A][B] as indicated by **1 or more** of the following:
 - Heart rate greater than 100 beats per minute in adult[A][B](1)
 - Heart rate greater than 85 beats per minute in child 13 to 17 years of age[A][C](2)
 - Heart rate greater than 95 beats per minute in child 6 to 12 years of age[A][C](2)
 - Heart rate greater than 110 beats per minute in child 1 to 5 years of age[A][C](2)
 - Heart rate greater than 120 beats per minute in infant 3 to 11 months of age[A][C](2)
 - Heart rate greater than 150 beats per minute in infant 1 or 2 months of age[A][C](2)

References

- Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
- Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

- B. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. For example, an elderly patient on a beta-blocker medication with a baseline resting heart rate of 60 beats per minute may be clinically tachycardic at a heart rate of 94 beats per minute. Likewise, a patient who is upset, in pain, or nervous in the emergency department with a heart rate of 106 beats per minute may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as chest pain or hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.
- C. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. A patient who is upset, in pain, or nervous in the emergency department with an elevated heart rate may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Tachycardia absent

- Tachycardia[A][B] absent, as indicated by **1 or more** of the following:
 - Heart rate less than or equal to 100 beats per minute in adult[A][B](1)
 - Heart rate less than or equal to 85 beats per minute in child 13 to 17 years of age[A][B](2)
 - Heart rate less than or equal to 95 beats per minute in child 6 to 12 years of age[A][B](2)
 - Heart rate less than or equal to 110 beats per minute in child 1 to 5 years of age[A][B](2)
 - Heart rate less than or equal to 120 beats per minute in infant 3 to 11 months of age[A][B](2)
 - Heart rate less than or equal to 150 beats per minute in infant 1 or 2 months of age[A][B](2)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. For example, an elderly patient on a beta-blocker medication with a baseline resting heart rate of 60 beats per minute may be clinically tachycardic at a heart rate of 94 beats per minute. Likewise, a patient who is upset, in pain, or nervous in the emergency department with a heart rate of 106 beats per minute may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as chest pain or hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Tachypnea

- Tachypnea,[A][B] as indicated by **1 or more** of the following:
 - Respiratory rate greater than 20 breaths per minute in adult[A][B](1)
 - Respiratory rate greater than 18 breaths per minute in child 13 to 17 years of age[A][B](2)
 - Respiratory rate greater than 22 breaths per minute in child 6 to 12 years of age[A][B](2)
 - Respiratory rate greater than 25 breaths per minute in child 3 to 5 years of age[A][B](2)
 - Respiratory rate greater than 30 breaths per minute in child 1 or 2 years of age[A][B](2)
 - Respiratory rate greater than 40 breaths per minute in infant 6 to 11 months of age[A][B](2)
 - Respiratory rate greater than 45 breaths per minute in infant 3 to 5 months of age[A][B](2)
 - Respiratory rate greater than 55 breaths per minute in infant 1 or 2 months of age[A][B](2)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of respiratory rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline respiratory rate and whether the respiratory rate is indicative of significant underlying respiratory or other pathology. A mildly increased respiratory rate (eg, 20 to 22 breaths per minute in an adult), absent other indications of serious illness (eg, hypoxemia, metabolic acidosis, decreased tidal volume, pain), may be transitory and not clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachypnea. Whether a respiratory rate above the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment. In addition, the measurement of the respiratory rate is subject to error. A medical textbook states that because there can be significant breath-to-breath variation in respiratory rate, the rate should be assessed for at least 30 seconds, preferably for a full minute.(1) This same textbook also states that new technologies designed to measure the respiratory rate have not proved to be clinically useful.(1)

Vomiting that is severe or persistent

- Vomiting and **1 or more** of the following(1)(2)(3)(4):
 - Pattern or content of vomitus suggests severe underlying cause or complication (eg, projectile, feculent, bilious, coffee ground, bloody).
 - Appropriate antiemetic treatment (eg, in observation care) does not sufficiently reduce vomiting.
 - Treatment regimen necessary to adequately control vomiting requires inpatient level of care (eg, not available in outpatient setting).

References

1. Guttman J. Nausea and vomiting. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:227-239.e2.
2. Hasler WL. Nausea, vomiting, and indigestion. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:291-297.

3. Maqbool A, Liacouras CA. Major symptoms and signs of digestive tract disorders. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:2224-2238.
4. Hammershaimb EA, Kotloff KL. Acute gastroenteritis in children. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:2355-2376.

Codes

ICD-10 Diagnosis: B02.24, F45.42, G89.0, G89.11, G89.12, G89.18, G89.21, G89.22, G89.28, G89.29, G89.4, G90.50, G90.511, G90.512, G90.513, G90.519, G90.521, G90.522, G90.523, G90.529, G90.59, R52 [Hide]

MCG Health
Inpatient & Surgical Care 30th Edition
Copyright © 2026 MCG Health, LLC
All Rights Reserved

Last Update: 1/25/2026 6:33:53 AM
Build Number: 30.0.2026012500524.025256