

Renal Failure, Acute

ORG: M-326 (ISC)

[Link to Codes](#)

MCG Health

Inpatient & Surgical Care
30th Edition

- Care Planning - Inpatient Admission and Alternatives
 - Clinical Indications for Admission to Inpatient Care
 - Supplemental Medicare Criteria
 - Alternatives to Admission
- Hospitalization
 - Optimal Recovery Course
 - Goal Length of Stay - **2 days**
 - Extended Stay
- Discharge
 - Discharge Planning
 - Discharge Destination
- Evidence Summary
 - Criteria
 - Hospitalization
 - Length of Stay
 - Rationale
 - Related CMS Coverage Guidance
- References
- Footnotes
- Definitions
- Codes

Care Planning - Inpatient Admission and Alternatives

Clinical Indications for Admission to Inpatient Care

Note: *Some patients may be appropriate for Observation care. For consideration of observation care, see Renal Failure, Acute: Observation Care* [↗](#) ISC.

- Admission is indicated for **1 or more** of the following^{[A][B](1)(2)(3):}^N
 - ☐ Acute renal failure (stage 3 acute kidney injury),^{[A][B]} as indicated by **1 or more** of the following:
 - Acute^[A] rise in serum creatinine to 3 times its baseline value^[C] or higher⁽¹⁾⁽²⁾⁽⁶⁾

- Acute[A] rise in serum creatinine to at least 1.5 times its baseline value[C] (increase of 50%) and serum creatinine is greater than 4 mg/dL (354 micromoles/L)(1)(2)(6)(7)
 - In child younger than 18 years, estimated glomerular filtration rate less than 35 mL/min/1.73m² (0.58 mL/sec/1.73m²)  eGFR - Pediatric Calculator and **1 or more** of the following:
 - Acute[A] rise in serum creatinine to at least 1.5 times its baseline value[C] (increase of 50%)(8)
 - Within past 48 hours, rise in serum creatinine greater than 0.3 mg/dL (26.5 micromoles/L)(8)
 - Decreased urine output,[D] as indicated by **ALL** of the following:
 - Assessment of urine output appropriate (eg, adequate volume status, no urinary obstruction)(1)
 - Oligoanuria, as indicated by **1 or more** of the following(1):
 - Urine output less than 0.3 mL/kg/hr for 24 hours(1)(8)
 - Anuria (urine output less than 0.1 mL/kg/hr) for 12 hours(1)(8)
 - Initiation of renal replacement therapy required(7)(9)(10)(11)
- ☐ Acute[A] kidney injury (eg, rise in serum creatinine from baseline) requiring inpatient care, as indicated by **1 or more** of the following(12):
- Hemodynamic instability (1)(8)
 - Altered mental status (1)(8)
 - Cardiac arrhythmias of immediate concern (1)(8)(13)
 - Severe hypertension (SBP greater than 180 mm Hg or DBP greater than 120 mm Hg in adults or greater than the 95th percentile for age, gender, and height plus 30 mm Hg in pediatric patients)(8)(14)(15)
 - Significant respiratory findings (eg, pulmonary edema) that are severe (eg, Hypoxemia) or persist despite observation care(2)(8)
 - Clinically significant electrolyte abnormality that is severe (eg, hyperkalemia with severe ECG findings)[E] or persists despite observation care(2)(8)(18)
 - Clinically significant metabolic abnormality (eg, metabolic acidosis) that is severe or persists despite observation care(2)(8)
 - Suspected etiology needing inpatient care (eg, Goodpasture syndrome, ANCA-associated vasculitis, cryoglobulinemic vasculitis, hemolytic uremic syndrome, lupus nephritis, hepatorenal syndrome, Kawasaki disease, possible need for plasmapheresis)(2)(8)(19)
 - Other finding or manifestation of acute renal disease necessitating inpatient care (eg, hemoptysis or concern for pulmonary hemorrhage, pericarditis, Anasarca)(3)(8)(12)(20)
 - Acute[A] kidney injury that persists or worsens despite observation care, as indicated by **1 or more** of the following:
 - Acute[A] rise in serum creatinine between 2 and 2.9 times its baseline value[C] and creatinine continues to rise during observation care (ie, has not plateaued)(1)(7)(8)
 - Decreased urine output,[D] as indicated by **ALL** of the following:
 - Assessment of urine output appropriate (eg, adequate volume status, no urinary obstruction)(1)(8)
 - Urine output less than 0.5 mL/kg/hr for 12 hours(1)(8)
 - Urine output has not improved in observation care (ie, remains less than 0.5 mL/kg/hr)(1)(8)

Supplemental Medicare Criteria

Note: These criteria are to be considered for Medicare patients if none of the standard Clinical Indications for Admission to Inpatient Care apply.

- Patient with Medicare coverage requires inpatient admission, as indicated by **1 or more** of the following(21):
 - Admitting clinician expects patient to require hospital care for less than 2 midnights but, based on complex medical factors documented in medical record, judges that inpatient care is necessary (case-by-case exception). The medical record must contain sufficient documentation to make clear the rationale for the exception.

- o Patient has need for intubation and mechanical ventilation that is new (ie, did not present to hospital already on mechanical ventilation).
- o Treatment plan for hospital admission includes procedure designated by CMS as inpatient only (ie, on Inpatient Only List).
- o Patient has already received medically necessary hospital care that meets 2-midnight benchmark (excluding activities such as triage/intake, delays in provision of care, or time added due to patient or family convenience). The medical record must contain sufficient documentation to make clear the medical necessity for hospital care across 2 or more midnights.

Alternatives to Admission

- Alternatives include(2)(3)(8):
 - o Outpatient care in emergency department, rapid treatment site, urgent care center, ambulatory surgery center, or dialysis center
 - Parenteral fluids or medications(22)
 - Evaluation for obstructive nephropathy if suspected (eg, renal ultrasound)(22)(23)
 - Cessation of nephrotoxic drugs(23)(24)
 - Referral for nephrology consultation(23)(25)
 - Palliative care consultation(26)(27)
 - o Home care
 - Nursing visit for assessment of clinical status
 - Laboratory testing
 - Parenteral fluids or medications
 - o Recovery facility
 - Parenteral fluid and medications correction
 - Electrolyte correction
 - Frequent assessment and treatment
 - Management of comorbidities
 - o Observation. See Renal Failure, Acute: Observation Care [ISC](#) guideline as appropriate.

Hospitalization

Optimal Recovery Course

Day	Level of Care	Clinical Status	Activity	Routes	Interventions	Medications
1	• ICU or floor[F] • Social Determinants of Health Assessment • Pediatric Readmission Risk Assessment • Adult Readmission Risk Assessment • Extended Stay Risk Assessment	• Clinical Indications met[G] • Possible lethargy, impaired sensorium • Possible multiple metabolic abnormalities[H]	• Activity as tolerated	• Parenteral or oral hydration • Parenteral or oral medications[I] • Oral diet as tolerated	• Urinalysis and other urine tests, CBC, renal function tests, hepatitis B test, other lab tests • CXR, ECG, renal ultrasound • Possible insertion of temporary venous or peritoneal dialysis access • Manage metabolic abnormalities	• Discontinue potentially nephrotoxic medications • Possible medical therapies[J]

- Discharge planning

- Manage obstructive nephropathy if indicated
- Possible dialysis or renal replacement therapy

2	• Floor • Social Determinants of Health Assessment • Pediatric Readmission Risk Assessment • Adult Readmission Risk Assessment • Extended Stay Risk Assessment	• Hypotension absent • Electrolyte abnormalities absent or improved • Acid-base abnormalities absent or improved • Mental status at baseline or improved • Renal function stable or improving	• Activity as tolerated	• Parenteral or oral hydration • Parenteral or oral medications • Oral diet as tolerated	• Possible dialysis or renal replacement therapy • Possible renal biopsy • Monitor electrolytes, renal function tests, acid-base, and volume status	• Possible medical therapies
3	• Social Determinants of Health Assessment • Pediatric Readmission Risk Assessment • Adult Readmission Risk Assessment • Extended Stay Risk Assessment • Floor to discharge • Complete discharge planning	• Hemodynamic stability • Mental status at baseline • Tachypnea absent • Hypoxemia absent • Etiology requiring inpatient treatment absent • Electrolyte abnormalities absent or acceptable for next level of care • Acid-base abnormalities absent • Renal function stable and acceptable for next level of care • Dialysis not needed or access and plan established • Discharge plans and education understood	• Ambulatory or acceptable for next level of care[K]	• Oral hydration[L] • Oral medications or regimen acceptable for next level of care • Oral diet or acceptable for next level of care • Possible calorie count, nutritional assessment	• Establish long-term treatment plan[M] • Monitor electrolytes, renal function tests, acid-base, and volume status • Dietary and fluid management counseling	• Possible medical therapies

(1)(5)(8)NN

Recovery Milestones are indicated in **bold**.

Goal Length of Stay: 2 days

Note: Goal Length of Stay assumes optimal recovery, decision making, and care. Patients may be discharged to a lower level of care (either later than or sooner than the goal) when it is appropriate for their clinical status and care needs.

Extended Stay

Minimal (a few hours to 1 day), Brief (1 to 3 days), Moderate (4 to 7 days), and Prolonged (more than 7 days).

- Extended stay beyond goal length of stay may be needed for(5):
 - Failure to meet discharge criteria (recovery milestones within final day in Optimal Recovery Course)(1)
 - Expect brief stay extension.
 - Continued complications of uremia(9)
 - Patient with more severe or persistent renal failure may require longer ICU and acute hospital care.
 - Expect brief to moderate stay extension.
 - Care of comorbid complications(1)(2)(31)(32)
 - Patient with complex comorbidities, including heart failure, hemorrhage, burns, diabetes, infection, sepsis, peritonitis, malignancy, hepatorenal syndrome, cirrhosis, malnutrition, vasculitis, or rhabdomyolysis, may require prolonged hospitalization.
 - Expect brief to moderate stay extension.
 - Postpartum acute renal failure(2)
 - Expect brief stay extension.
 - New need for dialysis (eg, not previously arranged and prepared)(1)(28)
 - Patient requiring inpatient dialysis may require longer acute hospital care.
 - Expect brief stay extension.

See Common Complications and Conditions [ISC](#) for further information.

Discharge

Discharge Planning

- Discharge planning includes[N]:
 - Assessment of needs and planning for care, including(34)(35):
 - Develop and modify treatment plan (involving multiple providers) as needed.
 - Evaluate and address preadmission functioning as needed.
 - Evaluate and address psychosocial status issues as indicated. See Psychosocial Assessment [SR](#) for further information.
 - Evaluate and address social determinants of health (eg, housing, food). See Social Determinants of Health Screening Tool [SR](#) for further information.
 - Evaluate and address patient or caregiver preferences as indicated.
 - Identify skilled services needed at next level of care, with specific attention to:
 - Coordination of care with outpatient center (eg, infusion, radiation, dialysis, surgery)(36)
 - Dialysis-related care, including laboratory tests(36)

- Electrolyte management(37)
- Genitourinary status assessment(37)
- Nutrition and hydration management(38)
- Early identification of anticipated discharge destination; options include(35)(39):
 - Home; considerations include:
 - Home safety assessment. See Home Safety Assessment [SR](#) for further information.
 - ☐ Patient safe to go home; examples include(40)(41)(42):
 - Medical status stable for patient's condition
 - Functional care can safely be provided with available resources.
 - Mental status stable for patient's condition
 - Medication availability confirmed and reconciliation complete
 - Patient/caregiver education completed with written discharge instructions provided
 - Community resources identified and referrals made, as needed
 - Home care arranged, if indicated
 - Necessary medical equipment delivery arranged or available in home, if indicated
 - Necessary medical supplies ordered, or patient/caregiver can obtain, if indicated
 - Access to follow-up care
 - Self-management ability if appropriate. See Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (IADL) Assessment [SR](#) for further information.
 - Caregiver need, ability, and availability
 - Post-acute skilled care or custodial care as indicated. See Discharge Planning Tool [SR](#) for further information.
- Transitions of care plan complete, including(35)(39)(43):
 - Patient and caregiver education complete.
 - See Teach-Back Tool [SR](#) for further information.
 - See Renal Failure, Acute: Patient Education for Clinicians [SR](#) for further information.
 - ☐ Medication reconciliation completion includes(44)(45):
 - Compare patient's discharge list of medications (prescribed and over-the-counter) against provider's admission or transfer orders.
 - Assess each medication for correlation to disease state or medical condition.
 - Report medication discrepancies to prescribing provider, attending physician, and primary care provider, and ensure accurate medication order is identified.
 - Provide reconciled medication list to all treating providers.
 - Confirm that patient or caregiver can acquire medication.
 - Educate patient and caregiver.
 - Provide complete medication list to patient and caregiver.
 - Importance of presenting personal medication list to all providers at each care transition, including all provider appointments
 - Reason, dosage, and timing of medication (eg, use "teach-back" techniques)(46)
 - Encourage communication between patient, caregiver, and pharmacy for obtaining prescriptions, setting up home medication delivery, and reviewing for drug-drug interactions.
 - See Medication Reconciliation Tool [SR](#) for further information.
 - Plan communicated to patient, caregiver, and all members of care team, including(47):
 - Inpatient care and service providers

- Primary care provider
- All post-discharge care and service providers
- Appointments planned or scheduled, which may include:
 - Primary care provider(48)
 - Dialysis(12)(49)(50)
 - Dietitian(38)
 - General surgeon(51)
 - Nephrologist(52)
 - Specialists for management of comorbidities as needed(52)
 - Urologist(53)
 - Vascular surgeon(54)
 - Other
- Outpatient testing and procedure plans made, which may include:
 - Laboratory testing(38)
 - Ultrasound(55)
 - Other
- Referrals made for assistance or support, which may include:
 - Community services(56)
 - Financial, for follow-up care, medication, and transportation(57)
 - Self-help or support groups(56)
 - Tobacco use treatment(38)(58)
 - Other
- Medical equipment and supplies coordinated (ie, delivered or delivery confirmed), which may include:
 - Blood pressure cuff(59)
 - Dialysis equipment and supplies(60)
 - Weight scale
 - Other

Discharge Destination

- Post-hospital levels of admission may include:
 - Home.
 - Home healthcare. See Home Care Indications for Admission Section [HC](#) in Renal Failure, Acute guideline in Home Care.
 - Recovery facility care. See Recovery Facility Care Indications for Admission Section [RFC](#) in Renal Failure, Acute guideline in Recovery Facility Care.

Evidence Summary

Criteria

The evidence for the clinical indications found in this guideline includes 8 published peer reviewed articles, 2 specialty society or other evidence-based guidelines, and 7 book sections.

Analysis of national hospital discharge data for a commercially insured population shows 55.4% of patients seen in an emergency department with a principal diagnosis of acute renal failure were admitted to inpatient care.(4) **(EG 3)** Analysis of hospital discharge data for a Medicare-insured population shows 80.4% of patients seen in an emergency department with a principal diagnosis of acute renal failure were admitted to inpatient care.(4) **(EG 3)** Analysis of an all-payer database shows a mean length of stay of 5.1 days for patients admitted to inpatient care.(5) **(EG 3)** A consensus guideline states that patients with acute renal failure, defined as stage 3 acute kidney injury, are appropriate for admission to inpatient care or the ICU to facilitate diagnostic workup, nephrology evaluation, and management of volume overload or severe metabolic or electrolyte abnormalities.(1) **(EG 2)** Consensus guidelines state that patients with acute kidney failure who require urgent renal replacement therapy should be admitted to inpatient care, often the ICU.(1)(3) **(EG 2)**

Hospitalization

A multinational, pragmatic, randomized controlled trial of adult patients with kidney failure who received at least 3 months of high-flux hemodialysis compared high-dose hemodiafiltration with high-flux hemodialysis (1360 patients, with 683 in the high-dose hemodiafiltration group and 677 in the high-flux hemodialysis group), and found that high-dose hemodiafiltration was associated with decreased risk of death from any cause (hazard ratio 0.77, 95% confidence interval 0.65 to 0.93) when compared with high-flux hemodialysis.(30) **(EG 2)** A retrospective review of adult patients who were started on outpatient hemodialysis for acute kidney injury (1754 patients with acute kidney injury, 70 dialysis centers) found that 41% of those patients (725 patients) recovered kidney function and discontinued hemodialysis.(29) **(EG 2)**

Length of Stay

Analysis of national hospital discharge data for a pediatric population shows 43% of hospitalized patients with a principal diagnosis of acute renal failure were discharged in 2 days or less.(5) **(EG 3)** Analysis of national hospital discharge data shows 30% of hospitalized adult patients with a primary diagnosis of acute renal failure were discharged in 2 days or less.(5) **(EG 3)**

Rationale

Use of this MCG care guideline helps the clinician identify patient-specific complex clinical factors that make it reasonable to expect a necessity for hospital care across 2 or more midnights. The evidence-based clinical criteria assist the clinician in the decision to appropriately admit a patient to inpatient care. For Medicare enrollees, MCG care guidelines also support the clinician in the decision to appropriately admit a patient to inpatient care when there is not an expectation of 2 or more midnights of hospital care (eg, CMS' Two-Midnight Rule exceptions).

Use of these evidence-based clinical criteria to support decision making around the need for inpatient treatment is of clinical benefit to the patient and is crucial for quality patient care. Use of evidence-based clinical criteria ensures patients receive the most appropriate care for their specific condition, reduces unnecessary risks, promotes recovery, and provides a standard that can reduce disparities in care. Evidence-based clinical criteria not only help align treatment decisions with best practice, but they can also help reduce unwarranted variation in care by fostering consistency and equality in healthcare provision across regions and facilities. Appropriate use of the evidence-based clinical criteria in conjunction with the Supplemental Medicare Criteria ensures Medicare patients receive full access to Medicare benefits (eg, inpatient care), without risk of delay or decreased access, based on variables such as clinical severity of illness, length of provision of medically necessary hospital-level care, or satisfaction of specified Medicare criteria as directed by CMS. Use of these criteria will help ensure that the patient receives necessary care in the appropriate setting.

Related CMS Coverage Guidance

This guideline supplements but does not replace, modify, or supersede existing Medicare regulations or applicable National Coverage Determinations (NCDs) or Local Coverage Determinations (LCDs).

Code of Federal Regulations (CFR): 42 CFR 412.3(21); 42 CFR 419.22(61); 42 CFR 422.101(62)

Internet-Only Manual (IOM) Citations: CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 1 - Inpatient Hospital Services Covered Under Part A(63); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 6 - Hospital Services Covered Under Part B(64); CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 15 - Covered Medical and Other Health Services(65); CMS IOM Publication 100-08, Medicare Program Integrity Manual, Chapter 6, Section 6.5 - Medical Review of Inpatient Hospital Claims for Part A Payment(66)

Medicare Coverage Determinations: Medicare Coverage Database(67)

References

1. Kidney Disease: Improving Global Outcomes (KDIGO) Acute Kidney Injury Work Group. KDIGO clinical practice guideline for acute kidney injury. Kidney International. Supplement 2012;2(1):1-138. (Reaffirmed 2025 Apr) [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32]
2. Weisbord SD, Palevsky PM. Prevention and management of acute kidney injury. In: Yu AS, Chertow GM, Luyckx VA, Marsden PA, Skorecki K, Taal MW, editors. Brenner and Rector's The Kidney. 11th ed. Philadelphia, PA: Elsevier; 2020:940-977.e15. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14]
3. Acute kidney injury: prevention, detection and management. NICE Clinical Guidance NG148 [Internet] National Institute for Health and Care Excellence. 2024 Oct 16 Accessed at: <https://www.nice.org.uk/guidance>. [created 2019; accessed 2025 Jul 21] [Context Link 1, 2, 3, 4]
4. Proprietary health insurance data sources, 2024; and Medicare 5% Standard Analytical File, 2023. [Context Link 1, 2]
5. Premier PINC AI™ Healthcare Database (PHD), 01/01/2023-12/31/2024. Premier, Inc. [Context Link 1, 2, 3, 4, 5]
6. Wolfson AB. Renal failure. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:1163-1180.e2. [Context Link 1, 2]
7. Ehmann MR, et al. Epidemiology and clinical outcomes of community-acquired acute kidney injury in the emergency department: a multisite retrospective cohort study. American Journal of Kidney Diseases 2024;83(6):762-771.e1. DOI: 10.1053/j.ajkd.2023.10.009. [Context Link 1, 2, 3] View abstract...
8. Kliegman RM, et al. Renal failure. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:3242-3253. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19]
9. Gaudry S, Palevsky PM, Dreyfuss D. Extracorporeal kidney-replacement therapy for acute kidney injury. New England Journal of Medicine 2022;386(10):964-975. DOI: 10.1056/NEJMra2104090. [Context Link 1, 2, 3] View abstract...
10. Freedman SB, van de Kar NCAJ, Tarr PI. Shiga toxin-producing escherichia coli and the hemolytic-uremic syndrome. New England Journal of Medicine 2023;389(15):1402-1414. DOI: 10.1056/NEJMra2108739. [Context Link 1] View abstract...
11. Mazumder NR, Junna S, Sharma P. The diagnosis and non-pharmacological management of acute kidney injury in patients with cirrhosis. Clinical Gastroenterology and Hepatology 2023;21(10S):S11-S19. DOI: 10.1016/j.cgh.2023.04.033. [Context Link 1] View abstract...
12. Agarwal A, Barasch J. Acute kidney injury. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:770-775.e1. [Context Link 1, 2, 3]
13. Genovesi S, et al. The conundrum of the complex relationship between acute kidney injury and cardiac arrhythmias. Nephrology, Dialysis, Transplantation 2023;38(5):1097-1112. DOI: 10.1093/ndt/gfac210. [Context Link 1] View abstract...
14. Jones DW, et al. 2025 AHA/ACC/AANP/AAPA/ABC/ACCP/ACPM/AGS/AMA/ASPC/NMA/PCNA/SGIM guideline for the prevention, detection, evaluation and management of high blood pressure in adults: a report of the American College of Cardiology/American Heart Association Joint Committee on Clinical Practice Guidelines. Circulation 2025;DOI: 10.1161/CIR.0000000000001356. [Context Link 1] View abstract...
15. Flynn JT, et al. Clinical practice guideline for screening and management of high blood pressure in children and adolescents. Pediatrics 2017;140(3):e20171904. DOI: 10.1542/peds.2017-1904. (Reaffirmed 2025 Jul) [Context Link 1] View abstract...
16. Mirvis DM, Goldberger AL. Electrocardiography. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:141-174.e3. [Context Link 1]
17. Mount DB. Fluid and electrolyte disturbances. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:338-356. [Context Link 1]

18. Kreitzer N, et al. EMCREG-international multidisciplinary consensus panel on management of hyperkalemia in chronic kidney disease and heart failure. *Cardiorenal Medicine* 2025;15(1):133-152. DOI: 10.1159/000543385. [Context Link 1] View abstract...
19. Friedman KG, Jone PN. Update on the management of Kawasaki disease. *Pediatric Clinics of North America* 2020;67(5):811-819. DOI: 10.1016/j.pcl.2020.06.002. [Context Link 1] View abstract...
20. Free M, Jennette JC, Falk RJ, Jain K. Renal and systemic vasculitis. In: Johnson RJ, Floege J, Tonelli M, editors. *Comprehensive Clinical Nephrology*. 7th ed. Elsevier; 2024:294-308.e3. [Context Link 1]
21. Centers for Medicare and Medicaid Services. "Admissions." 42 CFR 412.3 Washington, DC 2025 Jul [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/current/title-42/chapter-IV/subchapter-B/part-412>. [Context Link 1, 2]
22. Co I, Gunnerson K. Emergency department management of acute kidney injury, electrolyte abnormalities, and renal replacement therapy in the critically ill. *Emergency Medicine Clinics of North America* 2019;37(3):459-471. DOI: 10.1016/j.emc.2019.04.006. [Context Link 1, 2] View abstract...
23. Kanagasundaram S, et al. Acute Kidney Injury (AKI). Clinical Practice Guideline [Internet] Renal Association. 2019 Aug Accessed at: <https://renal.org/>. [accessed 2025 Sep 16] [Context Link 1, 2, 3]
24. Perazella MA, Rosner MH. Drug-induced acute kidney injury. *Clinical Journal of the American Society of Nephrology* 2022;17(8):1220-1233. DOI: 10.2215/CJN.11290821. [Context Link 1] View abstract...
25. Babroudi S, Weiner DE. Acute kidney injury care following hospitalization: care provision and public policy for acute kidney injury survivors. *Advances in Kidney Disease and Health* 2025;32(2):205-216. DOI: 10.1053/j.akdh.2024.10.002. [Context Link 1] View abstract...
26. Wong SPY, et al. Long-term outcomes among patients with advanced kidney disease who forgo maintenance dialysis: a systematic review. *JAMA Network Open* 2022;5(3):Online. DOI: 10.1001/jamanetworkopen.2022.2255. [Context Link 1] View abstract...
27. Petrosyan R, et al. Healthcare resource utilization and goals of care discussions in patients with cirrhosis and acute kidney injury. *Digestive Diseases and Sciences* 2024;69(6):1948-1955. DOI: 10.1007/s10620-024-08317-9. [Context Link 1] View abstract...
28. Liu C, et al. Predicting successful continuous renal replacement therapy liberation in critically ill patients with acute kidney injury. *Journal of Critical Care* 2021;66:6-13. DOI: 10.1016/j.jcrc.2021.07.020. [Context Link 1, 2] View abstract...
29. McCoy IE, Weinhandl E, Hussein W, Hsu CY. Initial management and potential opportunities to deprescribe dialysis among patients with AKI-D patients after hospital discharge. *Journal of the American Society of Nephrology* 2023;34(12):1949-1951. DOI: 10.1681/ASN.0000000000000225. [Context Link 1, 2] View abstract...
30. Blankestijn PJ, et al. Effect of hemodiafiltration or hemodialysis on mortality in kidney failure. *New England Journal of Medicine* 2023;389(8):700-709. DOI: 10.1056/NEJMoa2304820. [Context Link 1] View abstract...
31. Harding JL, Li Y, Burrows NR, Bullard KM, Pavkov ME. US trends in hospitalizations for dialysis-requiring acute kidney injury in people with versus without diabetes. *American Journal of Kidney Diseases* 2020;75(6):897-907. DOI: 10.1053/j.ajkd.2019.09.012. [Context Link 1] View abstract...
32. Neyra JA, et al. Prediction of mortality and major adverse kidney events in critically ill patients with acute kidney injury. *American Journal of Kidney Diseases* 2023;81(1):36-47. DOI: 10.1053/j.ajkd.2022.06.004. [Context Link 1] View abstract...
33. Hudson T. The role of social determinates of health in discharge practices. *Nursing Clinics of North America* 2021;56(3):369-378. DOI: 10.1016/j.cnur.2021.04.004. [Context Link 1] View abstract...
34. Chovanec KA, Arsene C, Beck A, Zachrich K, Liedel B, Wolff-Elliott J. Association of discharge disposition with outcomes. *Population Health Management* 2021;24(1):116-121. DOI: 10.1089/pop.2019.0176. [Context Link 1] View abstract...
35. Centers for Medicare and Medicaid Services. "Condition of participation: Discharge planning." 42 CFR Pt. 482.43 Washington, DC 2025 Jan 01 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>. [Context Link 1, 2, 3]
36. Management of patients with kidney disorders. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:1554-1603. [Context Link 1, 2]
37. Dennison HA. Acute kidney injury and chronic kidney disease. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:1232-1264. [Context Link 1, 2]

38. The client with alterations in the kidney and urinary tract. In: Haugen N, Galura SJ, editors. *Ulrich & Canale's Nursing Care Planning Guides*, 8th Edition Revised Reprint with 2021-2023 NANDA-I Updates. 8th ed. Elsevier; 2022:633-671. [Context Link 1, 2, 3, 4]
39. Targeted care planning and care transitions. In: Perez R, editor. *CMSA's Integrated Case Management*. 2nd ed. Springer Publishing Company LLC; 2023:178-189. [Context Link 1, 2]
40. Elements of Excellence in Transitions of Care (TOC). TOC Checklist [Internet] National Transitions of Care Coalition. 2006 Accessed at: <https://www.ntocc.org>. [accessed 2025 Sep 16] [Context Link 1]
41. Medical-surgical nursing. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:33-55. [Context Link 1]
42. Transition of care. In: Gillingham DC, editor. *Foundations of Case Management*. Blue Bayou Press; 2021:137-144. [Context Link 1]
43. Becker C, et al. Interventions to improve communication at hospital discharge and rates of readmission: a systematic review and meta-analysis. *JAMA Network Open* 2021;4(8):e2119346. DOI: 10.1001/jamanetworkopen.2021.19346. [Context Link 1] View abstract...
44. The nursing process in drug therapy and patient safety. In: Tucker RG, editor. *Karch's Focus on Nursing Pharmacology*. 9th ed. Wolters Kluwer; 2023:47-57. [Context Link 1]
45. Adverse drug reactions and medication errors. In: Burchum JR, Rosenthal LD, editors. *Lehne's Pharmacology for Nursing Care*. 12th ed. Elsevier; 2025:64-75. [Context Link 1]
46. Safe medication preparation. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. *Clinical Nursing Skills and Techniques*. 11th ed. Elsevier; 2025:586-607. [Context Link 1]
47. The case manager's roles, functions, and activities. In: Gillingham DC, editor. *Foundations of Case Management*. Blue Bayou Press; 2021:9-14. [Context Link 1]
48. Anderson TS, Ayanian JZ, Herzig SJ, Souza J, Landon BE. Gaps in primary care follow-up after hospital discharge among Medicare beneficiaries. *Journal of the American Geriatrics Society* 2025;DOI: 10.1111/jgs.19496. [Context Link 1] View abstract...
49. Kotanko P, Kuhlmann MK, Chan C, Levin NW. Hemodialysis: principles and techniques. In: Johnson RJ, Floege J, Tonelli M, editors. *Comprehensive Clinical Nephrology*. 7th ed. Elsevier; 2024:1091-1099.e1. [Context Link 1]
50. Bargman JM, Auguste BL. Peritoneal dialysis: principles, techniques, and adequacy. In: Johnson RJ, Floege J, Tonelli M, editors. *Comprehensive Clinical Nephrology*. 7th ed. Elsevier; 2024:1120-1129.e2. [Context Link 1]
51. Bouchard J, Macedo E. Prevention and nondialytic management of acute kidney injury. In: Johnson RJ, Floege J, Tonelli M, editors. *Comprehensive Clinical Nephrology*. 7th ed. Elsevier; 2024:850-862.e2. [Context Link 1]
52. Ostermann M, Lumlertgul N, Jeong R, See E, Joannidis M, James M. Acute kidney injury. *Lancet* 2025;405(10474):241-256. DOI: 10.1016/S0140-6736(24)02385-7. [Context Link 1, 2] View abstract...
53. Molitoris BA. Low-flow acute kidney injury: the pathophysiology of prerenal azotemia, abdominal compartment syndrome, and obstructive uropathy. *Clinical Journal of the American Society of Nephrology* 2022;17(7):1039-1049. DOI: 10.2215/CJN.15341121. [Context Link 1] View abstract...
54. Winkelman C, Duffy EG. Kidney disease. In: Buttaro TM, Polgar-Bailey P, Sandberg-Cook J, Dick KL, Montgomery JB, editors. *Primary Care: Interprofessional Collaborative Practice*. 7th ed. Elsevier; 2025:847-856. [Context Link 1]
55. Judd E, Sanders PW, Agarwal A. Diagnosis and clinical evaluation of acute kidney injury. In: Johnson RJ, Floege J, Tonelli M, editors. *Comprehensive Clinical Nephrology*. 7th ed. Elsevier; 2024:835-844.e2. [Context Link 1]
56. Snyder JS, Sump CA. Psychosocial support for the patient. In: Snyder JS, Sump CA, editors. *Swearingen's All-in-One Nursing Care Planning Resource*. 6th ed. Elsevier; 2024:97-105. [Context Link 1, 2]
57. Bragg-Underwood T. Family health. In: Nies MA, editor. *Community/Public Health Nursing*. 8th ed. Elsevier; 2024:404-427. [Context Link 1]
58. Harding MM. Substance use disorders in acute care. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:162-178. [Context Link 1]
59. Vital signs. In: Lynn P, editor. *Taylor's Clinical Nursing Skills*. 6th ed. Wolters Kluwer; 2023:42-96. [Context Link 1]
60. Renal and urinary disorders. In: Nettina SM, Nelson-Tuttle C, editors. *Lippincott Manual of Nursing Practice*. 12th ed. WoltersKluwer; 2025:568-616. [Context Link 1]

61. Centers for Medicare and Medicaid Services. "Hospital services excluded from payment under the hospital outpatient prospective payment system." 42 CFR 419.22 Washington, DC 2023 Jul [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
62. Centers for Medicare and Medicaid Services. "Requirements relating to basic benefits." 42 CFR 422.101 Washington, DC 2025 Jun 03 [accessed 2025 Jul 23] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
63. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09] [Context Link 1]
64. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 6 - Hospital Services Covered Under Part B Rev. 12425 [Internet] Centers for Medicare and Medicaid Services. 2023 Dec Accessed at: <http://www.cms.gov/manuals/>. [accessed 2025 Sep 09] [Context Link 1]
65. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 15 - Covered Medical and Other Health Services Rev. 13108 [Internet] Centers for Medicare and Medicaid Services. 2025 Apr 11 Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09] [Context Link 1]
66. Centers for Medicare and Medicaid Services. Medicare Program Integrity Manual. Chapter 6, Section 6.5 - Medical Review of Inpatient Hospital Claims for Part A Payment Rev. 10365 [Internet] Centers for Medicare and Medicaid Services. 2020 Oct 02 Accessed at: <https://www.cms.gov/medicare/regulations-guidance/manuals/internet-only-manuals-ioms>. [accessed 2025 Sep 09] [Context Link 1]
67. Medicare Coverage Database. [Internet] Centers for Medicare and Medicaid Services. Accessed at: <https://www.cms.gov/medicare-coverage-database/search.aspx?> Updated 2025 [accessed 2025 Oct 23] [Context Link 1]

Footnotes

[A] A specialty society practice guideline defines acute kidney injury as a change in renal function that is known or presumed to have occurred over the previous 7 days.(1) [A in Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10]

[B] For purposes of these criteria, the term acute renal failure can be viewed as stage 3 acute kidney injury in the KDIGO classification system.(1) Of note, in the ICD-10 diagnostic coding system (main coding scheme used in labeling diagnoses), the relevant codes covering the clinical entity of significant acute kidney injury use the term acute kidney failure (eg, code N17.9). [B in Context Link 1, 2, 3]

[C] If the baseline serum creatinine is not known, absent known or suspected chronic kidney disease, a baseline serum creatinine can be approximated by assuming an estimated glomerular filtration rate of 75 mL/min/1.73m² (1.25 mL/sec/1.73m²). (1) [C in Context Link 1, 2, 3, 4]

[D] The use of urine output as an indicator of acute kidney injury is important, as change in urine output can occur before a noticeable rise in creatinine.(1)(2) It is also suggested that urine output can identify patients with acute kidney injury that may otherwise be missed.(1)(2) However, urine output as a metric is only valid under the appropriate conditions. For example, the patient must not be hypovolemic, there must not be urinary obstruction, and the collection and measurement of urinary output must be complete and accurate.(1)(2) The use of urinary output to define acute kidney injury can be inaccurate in obese patients due to the nonlinear relationship between body weight and urine output. For example, in a 100-kg (220-lb) patient, urine output of 45 mL/hr over 12 hours (adequate output) would qualify as stage 2 acute kidney injury.(1)(2) [D in Context Link 1, 2]

[E] ECG findings include peaked T-waves, PR interval greater than 0.20 seconds, QRS interval greater than 0.12 seconds, and arrhythmia.(16)(17) [E in Context Link 1]

[F] ICU admission is appropriate for severe life-threatening acidosis, pulmonary edema, hypertensive emergency, hypotension, severe electrolyte disturbance, if an underlying critical illness is contributing to acute renal failure, or if continuous renal replacement therapy is required.(1)(9) See Intensive, Intermediate, and Telemetry Care Guidelines  ISC for further information. [F in Context Link 1]

[G] See Clinical Indications for Admission to Inpatient Care in this guideline. [G in Context Link 1]

[H] Important abnormalities include hyperkalemia and acidosis.(1) [H in Context Link 1]

[I] Drug doses should be modified, with monitoring of levels, when appropriate.(1) [I in Context Link 1]

[J] Medications may include calcium, phosphate binders, bicarbonate, vasoactive agents, and antihypertensives.(1) [J in Context Link 1]

[K] Patient is ambulatory or near baseline activity for age and development. [K in Context Link 1]

[L] Some patients may have their hydration needs met via alternative means (eg, percutaneous endoscopic gastrostomy tube). [L in Context Link 1]

[M] The treatment plan will depend on anticipated need for ongoing dialysis and concomitant vascular access and the patient's decision regarding initiating or continuing dialysis.(1)(28)(29) [M in Context Link 1]

[N] Discharge instructions should be given in the patient's and caregiver's native language using trained language interpreters whenever possible.(33) [N in Context Link 1]

Definitions

Adult Readmission Risk Assessment

- Risk of readmission is increased by presence (upon admission or during hospital stay) of **1 or more** of the following(1):
 - Clinically significant comorbid illness that necessitates (during current hospital stay) new or intensified treatment, prolongs length of stay, or impacts discharge level of care (eg, heart failure, COPD, active malignancy, cirrhosis, dementia, psychosis)(1)(2)(3)
 - End-stage renal disease or new need for dialysis(1)
 - Malnutrition[A](1)
 - Need for parenteral nutrition or supplemental enteral tube feeding (eg, via gastrostomy tube or jejunostomy tube)(1)
 - Need for transfusion of platelets or red blood cells(1)
 - Need for supplemental oxygen after discharge (new or pre-existing)(1)
 - Potential barriers to postdischarge clinical care (eg, homelessness, lack of transportation, impaired access to care or medication, active substance abuse)(1)
 - Frailty (ie, as indicated by a frailty measurement tool)[B][C](1)(5)(6)(7)
 - Two or more emergency department visits and/or inpatient admissions within previous 6 months(1)(8)

References

1. Proprietary health insurance data sources, 2022; and Medicare 100% Standard Analytical File, 2022.
2. Barnett ML, Hsu J, McWilliams JM. Patient characteristics and differences in hospital readmission rates. *JAMA Internal Medicine* 2015;175(11):1803-12. DOI: 10.1001/jamainternmed.2015.4660.
3. Berry JG, et al. Age trends in 30 day hospital readmissions: US national retrospective analysis. *British Medical Journal* 2018;360:k497.
4. Cederholm T, Bosaeus I. Malnutrition in adults. *New England Journal of Medicine* 2024;391(2):155-165. DOI: 10.1056/NEJMra2212159.
5. Sison SDM, et al. A crosswalk of commonly used frailty scales. *Journal of the American Geriatrics Society* 2023;71(10):3189-3198. DOI: 10.1111/jgs.18453.
6. Kim DH, Rockwood K. Frailty in older adults. *New England Journal of Medicine* 2024;391(6):538-548. DOI: 10.1056/NEJMra2301292.
7. Soares MR, Mahanna Gabrielli E, Manjarrez EC. The geriatric patient: frailty, prehabilitation, and postoperative delirium. *Medical Clinics of North America* 2024;108(6):1101-1117. DOI: 10.1016/j.mcna.2024.06.001.
8. Aubert CE, et al. Simplification of the HOSPITAL score for predicting 30-day readmissions. *BMJ Quality and Safety* 2017;26(10):799-805. DOI: 10.1136/bmjqs-2016-006239.

Footnotes

- A. There are many tools used to assess for malnutrition in adults. Near universal elements include recent weight loss and low muscle mass. Most tools include criteria based on BMI. Some include criteria around etiology such as low food intake vs disease burden or inflammation.(4) The Malnutrition Universal Screening Tool (MUST) is available at www.mdcalc.com/calc/10190/malnutrition-universal-screening-tool-must. The Nutrition Risk Screening 2002 (NRS-2002) tool is available at www.mdcalc.com/calc/4012/nutrition-risk-screening-2002-nrs-2002. A document that outlines the elements and scoring of several tools can be downloaded from <https://aspenjournals.onlinelibrary.wiley.com/action/downloadSupplement?doi=10.1002%2Fjpen.1440&file=jpen1440-sup-0001-Appendix.docx>.
- B. There are numerous measurement tools used to assess for frailty. Some of the more commonly used scales (with score indicating frailty in parentheses) include the following: Frailty Phenotype (3 or higher), 40-item Frailty Index (0.25 or higher), Study of Osteoporotic Fracture Index (2 or higher), Edmonton Frail Scale (8 or higher), FRAIL Scale (3 or higher), Clinical Frailty Scale (5 or higher), and Tilburg Frailty Indicator (5 or higher).(5)
- C. Frailty may be described as a phenotype or as an accumulation of health deficits. The Fried phenotype delineates a clinical syndrome resulting from altered metabolism coupled with abnormal stress responses.(6) This model includes 5 characteristic features: exhaustion (energy level), weakness (hand grip strength), slowness (gait speed), physical inactivity (falls), and weight loss.(6)(7) The presence of 3 or more of these deficits indicates the presence of frailty.(6)(7) Frailty as deficit accumulation considers findings in multiple domains (eg, cognitive, physical, nutritional, laboratory) and high-risk comorbidities.(6)(7) In this model, the degree of frailty is quantified by the number of indicative items present.(6)

Altered mental status

- Altered mental status (ie, different from baseline), as indicated by **1 or more** of the following(1)(2)(3)(4):
 - Confusional state (eg, disorientation, difficulty following commands, deficit in attention)
 - Lethargy (eg, awake or arousable, but with drowsiness; reduced awareness of self and environment)
 - Obtundation (ie, arousable only with strong stimuli, lessened interest in environment, slowed responses to stimulation)
 - Stupor (ie, may be arousable but patient does not return to normal baseline level of awareness)
 - Coma (ie, not arousable)

References

1. Maher PJ. Confusion. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:126-131.e1.
2. Lei C, Smith C. Depressed consciousness and coma. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:117-125.e1.
3. Abraham G, Maher PJ. Delirium and dementia. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:1269-1280.e2.
4. Holler-Managan YF. Neurologic evaluation. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:3550-3560.

Anasarca

- Anasarca is generalized edema that is widespread and may be more pronounced in dependent areas (eg, back while lying down, legs while sitting up).(1)

References

1. Kleczka J, Dabbouseh NM. Evaluation of the patient with cardiovascular disease. In: Wing EJ, Schiffman FJ, editors. Cecil Essentials of Medicine. 10 th ed. 1600 John F. Kennedy Blvd.: Elsevier; 2022:10-23.

Cardiac arrhythmias of immediate concern

- Cardiac arrhythmias or findings of immediate concern, as indicated by **1 or more** of the following(1)(2)(3):
 - Heart rhythms that are inherently dangerous or unstable, as indicated by **1 or more** of the following(4)(5)(6):
 - Resuscitated ventricular fibrillation or cardiac arrest
 - Ventricular escape rhythm
 - Sustained ventricular tachycardia (30 seconds or more of ventricular rhythm at greater than 100 beats per minute)
 - Nonsustained ventricular tachycardia and **1 or more** of the following:
 - Suspected cardiac ischemia as cause or consequence of ventricular tachycardia
 - Acute myocarditis
 - Unstable cardiac conduction defects, as indicated by **1 or more** of the following(7)(8):
 - Type II second-degree atrioventricular block
 - Third-degree atrioventricular block
 - New-onset left bundle branch block with suspected myocardial ischemia(9)
 - Any heart rhythm and **1 or more** of the following(5)(8)(10)(11)(12)(13):
 - Continuous long-term ECG monitoring needed (eg, initiation of drug requiring monitoring beyond observation care)
 - Patient has automatic implantable cardioverter-defibrillator that is repeatedly firing, malfunctioning, or in need of immediate adjustment of settings beyond scope of observation care.
 - Heart rhythms of concern due to **1 or more** of the following(8):
 - Hypotension
 - Respiratory distress
 - Association with other significant symptoms (eg, syncope)(10)(11)

References

1. Curtis AB, Tomaselli GF. Approach to the patient with cardiac arrhythmias. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1145-1162.e7.
2. Dalal AS, Van Hare GF. Disturbances of rate and rhythm of the heart. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:2844-2859.e1.
3. Joglar JA, et al. 2023 HRS Expert Consensus Statement on the management of arrhythmias during pregnancy. Heart Rhythm 2023;20(10):Online. DOI: 10.1016/j.hrthm.2023.05.017.
4. Garan H. Ventricular arrhythmias. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:332-340.
5. Al-Khatib SM, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. Circulation 2018;138(13):e272-e391. DOI: 10.1161/CIR.0000000000000549. (Reaffirmed 2025 Jun)
6. Myerburg RJ, Lampert R. Cardiac arrest and life-threatening arrhythmias. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:312-317.
7. Miller JM, Ellenbogen KA. Therapy for cardiac arrhythmias. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1208-1244.
8. Patton KK, Olgin JE. Bradyarrhythmias and atrioventricular block. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine. 12th ed. Elsevier; 2022:1312-1320.
9. Tan NY, Witt CM, Oh JK, Cha YM. Left bundle branch block: current and future perspectives. Circulation. Arrhythmia and Electrophysiology 2020;13(4):e008239. DOI: 10.1161/CIRCEP.119.008239.

10. Joglar JA, et al. 2023 ACC/AHA/ACCP/HRS Guideline for the diagnosis and management of atrial fibrillation: a report of the American College of Cardiology/American Heart Association joint committee on clinical practice guidelines. *Journal of the American College of Cardiology* 2024;149(1):Online. DOI: 10.1016/j.jacc.2023.08.017. (Reaffirmed 2025 Mar)
11. Zimetbaum P, Goldman L. Supraventricular ectopy and tachyarrhythmias. In: Goldman L, Cooney KA, editors. *Goldman-Cecil Medicine*. 27th ed. Elsevier; 2024:322-332.
12. Santucci PA, Wilber DJ. Electrophysiologic procedures and surgery. In: Goldman L, Cooney KA, editors. *Goldman-Cecil Medicine*. 27th ed. Elsevier; 2024:340-347.
13. Pendharkar SS, et al. AHA telemetry guidelines improve telemetry utilization in the inpatient setting. *American Journal of Managed Care* 2020;26(11):476-481. DOI: 10.37765/ajmc.2020.88525.

Extended Stay Risk Assessment

- Risk of extended length of stay is increased by presence of **1 or more** of the following(1):
 - New need for dialysis
 - Acute heart failure (eg, pulmonary edema)

References

1. Premier PINC AI™ Healthcare Database (PHD), 01/01/2023-12/31/2024. Premier, Inc.

Hemodynamic instability

- Hemodynamic instability, as indicated by **1 or more** of the following:
 - Vital sign abnormality not corrected by appropriate treatment, as indicated by **1 or more** of the following[A]:
 - Tachycardia that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)[A]
 - Hypotension that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)[A]
 - Orthostatic hypotension that persists despite appropriate treatment (eg, treatment of underlying cause, despite observation care)[A]
 - Severe hypotension in adult patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 80 mm Hg(1)[A]
 - Shock index greater than 1.0 (shock index is the heart rate divided by systolic blood pressure)[A](2)(3)(4)
 - Mean arterial pressure[B] less than 65 mm Hg  MAP Calculator[A](1)(6)
 - IV inotropic or vasopressor medication required to maintain adequate blood pressure or perfusion(3)(6)
 - Hypoperfusion due to hypotension, as indicated by **ALL** of the following:
 - Hypotension
 - Evidence of hypoperfusion, as indicated by **1 or more** of the following:
 - Lactate of 2.0 mmol/L (18 mg/dL) or more secondary to hypotension (ie, hypoperfusion)[C](3)(6)(8)
 - Metabolic acidosis (arterial or venous[D] pH less than 7.35) not otherwise explained(3)(6)(8)
 - Significant end organ dysfunction due to hypotension (eg, Altered mental status, myocardial ischemia, 2-fold or higher acute increase in serum creatinine)(3)(6)
 - Severe hypotension in pediatric patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 80 mm Hg in child 6 to 17 years of age[A](14)
 - Systolic blood pressure less than 75 mm Hg in child 6 months to 5 years of age[A](14)

- Shock index greater than 0.9 in child 13 to 17 years of age (shock index is the heart rate divided by systolic blood pressure)[A](2)(15)
- Shock index greater than 1.0 in child 7 to 12 years of age (shock index is the heart rate divided by systolic blood pressure)[A](2)(15)
- Shock index greater than 1.2 in child 4 to 6 years of age (shock index is the heart rate divided by systolic blood pressure)[A](2)(15)
- IV inotropic or vasopressor medication required to maintain adequate blood pressure or perfusion(5)(6)
- Hypoperfusion due to hypotension, as indicated by **ALL** of the following:
 - Hypotension
 - Evidence of hypoperfusion, as indicated by **1 or more** of the following:
 - Lactate of 2.0 mmol/L (18 mg/dL) or more secondary to hypotension (ie, hypoperfusion)[C](6)(8)
 - Metabolic acidosis (arterial or venous pH less than 7.35) not otherwise explained[D](5)(6)(14)
 - Significant end organ dysfunction due to hypotension (eg, Altered mental status, myocardial ischemia, 2-fold or higher acute increase in serum creatinine)(5)(6)(14)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Koch E, Lovett S, Nghiem T, Riggs RA, Rech MA. Shock index in the emergency department: utility and limitations. Open Access Emergency Medicine 2019;11:179-199. DOI: 10.2147/OAEM.S178358.
3. Angus DC. Approach to the patient with shock. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:658-665.
4. Balhara KS, Hsieh YH, Hamade B, Circh R, Kelen GD, Bayram JD. Clinical metrics in emergency medicine: the shock index and the probability of hospital admission and inpatient mortality. Emergency Medicine Journal 2017;34(2):89-94. DOI: 10.1136/emmermed-2015-205532.
5. Weiss SL, Tissieres P, Peters MJ. Shock. In: Kliegman RM, et al., editors. Nelson Textbook of Pediatrics. 22nd ed. Elsevier; 2025:600-611.
6. Singer M, et al. The Third International Consensus definitions for sepsis and septic shock (Sepsis-3). Journal of the American Medical Association 2016;315(8):801-810. DOI: 10.1001/jama.2016.0287.
7. Wardi G, Brice J, Correia M, Liu D, Self M, Tainter C. Demystifying lactate in the emergency department. Annals of Emergency Medicine 2020;75(2):287-298. DOI: 10.1016/j.annemergmed.2019.06.027.
8. Kraut JA, Madias NE. Lactic acidosis. New England Journal of Medicine 2014;371(24):2309-2319. DOI: 10.1056/NEJMra1309483.
9. Olivieri L, Chasm R. Diabetic ketoacidosis in the pediatric emergency department. Emergency Medicine Clinics of North America 2013;31(3):755-773. DOI: 10.1016/j.emc.2013.05.004.
10. Maloney GE, Glauser JM. Diabetes mellitus and disorders of glucose homeostasis. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:1543-1558.e2.
11. Arnold TD, Miller M, van Wessel KP, Evans JA, Balogh ZJ. Base deficit from the first peripheral venous sample: a surrogate for arterial base deficit in the trauma bay. Journal of Trauma 2011;71(4):793-7; discussion 797. DOI: 10.1097/TA.0b013e31822ad694.
12. Malinoski DJ, Todd SR, Slone S, Mullins RJ, Schreiber MA. Correlation of central venous and arterial blood gas measurements in mechanically ventilated trauma patients. Archives of Surgery 2005;140(11):1122-5. DOI: 10.1001/archsurg.140.11.1122.
13. Zakrisson T, McFarlan A, Wu YY, Keshet I, Nathens A. Venous and arterial base deficits: do these agree in occult shock and in the elderly? A Bland-Altman analysis. Journal of Trauma and Acute Care Surgery 2013;74(3):936-9. DOI: 10.1097/TA.0b013e318282747a.
14. Alexander PMA, et al. Cardiovascular dysfunction criteria in critically ill children: the PODIUM consensus conference. Pediatrics 2022;149(1 Suppl 1):S39-S47. DOI: 10.1542/peds.2021-052888F.
15. Ramgopal S. The shock index among children presenting to the emergency department: analysis of nationally representative sample. Journal of Emergency Medicine 2024;67(2):e146-e156. DOI: 10.1016/j.jemermed.2024.03.022.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. The mean arterial pressure (MAP) takes into account both SBP and DBP readings.(3)(5)
- C. There are numerous causes of an elevated lactate level. The most common are cardiogenic or hypovolemic shock, severe heart failure, severe trauma, or sepsis. However, there are also other etiologies to consider such as vigorous exercise, seizures, liver disease, or medication use (eg, metformin, beta2-agonists); therefore, interpretation of elevated lactate levels requires consideration of the clinical context (eg, well-appearing post exercise vs hypotensive). The severity and persistence of the elevation can sometimes be helpful in this differentiation. In most instances of lactic acidosis, blood pH is less than 7.35, with a serum bicarbonate level of 20 mEq/L (mmol/L) or lower. However, a coexisting respiratory or metabolic alkalosis can mask these findings.(7)(8)
- D. Analyses have concluded that venous and arterial pH measurements are highly correlated, with small differences between them that are usually not clinically significant.(9)(10)(11)(12)(13) If a mixed acid-base disorder is suspected, an arterial blood gas is indicated.(10)

Hemodynamic stability

- Hemodynamic stability, as indicated by **1 or more** of the following:
 - Hemodynamic abnormalities at baseline or acceptable for next level of care
 - Patient hemodynamically stable, as indicated by **ALL** of the following(1)(2):
 - Tachycardia absent
 - Hypotension absent
 - No evidence of inadequate perfusion (eg, no myocardial ischemia)
 - No other hemodynamic abnormalities (eg, no Orthostatic hypotension)

References

- Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
- Ramgopal S, Sepanski RJ, Martin-Gill C. Empirically derived age-based vital signs for children in the out-of-hospital setting. Annals of Emergency Medicine 2023;81(4):402-412. DOI: 10.1016/j.annemergmed.2022.09.019.

Hypotension

- Hypotension, as indicated by **1 or more** of the following:
 - Hypotension in adult patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 90 mm Hg[A][B](1)
 - Mean arterial pressure[C] less than 70 mm Hg[A][B]  MAP Calculator(1)(2)
 - Decrease in baseline systolic blood pressure of 30 mm Hg or more, with significant signs or symptoms due to lower blood pressure (eg, near syncope, syncope, chest pain)[A][B](1)
 - Hypotension in pediatric patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure less than 110 mm Hg in child 13 to 17 years of age[A][D](3)
 - Systolic blood pressure less than 100 mm Hg in child 6 to 12 years of age[A][D](3)
 - Systolic blood pressure less than 95 mm Hg in child 3 to 5 years of age[A][D](3)
 - Systolic blood pressure less than 90 mm Hg in child 1 or 2 years of age[A][D](3)

- Systolic blood pressure less than 80 mm Hg in infant 6 to 11 months of age^{[A][D](3)}
- Systolic blood pressure less than 70 mm Hg in infant 3 to 5 months of age^{[A][D](3)}
- Systolic blood pressure less than 65 mm Hg in infant 1 or 2 months of age^{[A][D](3)}

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Puskarich MA, Jones AE. Shock. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:34-41.e1.
3. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of blood pressure requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline blood pressure, medications, and clinical impact. For example, an elderly patient with heart failure may be prescribed medication with the intentional goal of a reduced pressure. If the patient's baseline SBP is 92 mm Hg to 96 mm Hg, absent signs or symptoms of hypotension, an emergency department SBP of 86 mm Hg should not automatically be categorized as hypotensive. This would hold for mean arterial pressure (MAP) as well. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term hypotension. Whether a blood pressure below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.
- C. The mean arterial pressure (MAP) takes into account both SBP and DBP readings.
- D. Interpretation of blood pressure requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline blood pressure, medications, and clinical impact. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term hypotension. Whether a blood pressure below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Hypotension absent

- Hypotension absent^[A] as indicated by **1 or more** of the following:
 - Hypotension absent in adult patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure greater than or equal to 90 mm Hg^{[A](1)}
 - Mean arterial pressure^[B] greater than or equal to 70 mm Hg  MAP Calculator^{[A](1)(2)}
 - Blood pressure at patient's baseline (eg, healthy adult with low systolic blood pressure), at intentional therapeutic goal (eg, patient with heart failure), or acceptable for next level of care (eg, blood pressure stable and no significant signs or symptoms due to low blood pressure)
 - Hypotension absent in pediatric patient, as indicated by **1 or more** of the following:
 - Systolic blood pressure greater than or equal to 110 mm Hg in child 13 to 17 years of age^{[A](3)}
 - Systolic blood pressure greater than or equal to 100 mm Hg in child 6 to 12 years of age^{[A](3)}
 - Systolic blood pressure greater than or equal to 95 mm Hg in child 3 to 5 years of age^{[A](3)}
 - Systolic blood pressure greater than or equal to 90 mm Hg in child 1 or 2 years of age^{[A](3)}
 - Systolic blood pressure greater than or equal to 80 mm Hg in infant 6 to 11 months of age^{[A](3)}

- Systolic blood pressure greater than or equal to 70 mm Hg in infant 3 to 5 months of age^{[A](3)}
- Systolic blood pressure greater than or equal to 65 mm Hg in infant 1 or 2 months of age^{[A](3)}
- Blood pressure at patient's baseline (eg, healthy child with low systolic blood pressure), at intentional therapeutic goal, or acceptable for next level of care (eg, blood pressure stable and no significant signs or symptoms due to low blood pressure)

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Puskarch MA, Jones AE. Shock. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:34-41.e1.
3. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. The mean arterial pressure (MAP) takes into account both SBP and DBP readings.

Hypoxemia

- Hypoxemia, as indicated by **1 or more** of the following⁽¹⁾:
 - Patient without baseline need for supplemental oxygen with oxygen saturation (SpO₂) of less than 90% or arterial blood gas partial pressure of oxygen (PaO₂) of less than 60 mm Hg (8.0 kPa) on room air^{[A][B]}
 - Patient without baseline need for supplemental oxygen who now requires supplemental oxygen to keep SpO₂ greater than 89% or PaO₂ greater than 59 mm Hg (7.9 kPa)^[A]
 - Patient with baseline need for supplemental oxygen who now requires increased supplemental oxygen to maintain oxygenation at baseline or acceptable level

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Jamali H, et al. Racial disparity in oxygen saturation measurements by pulse oximetry: evidence and implications. Annals of the American Thoracic Society 2022;19(12):1951-1964. DOI: 10.1513/AnnalsATS.202203-270CME.
3. Rojas-Camayo J, et al. Reference values for oxygen saturation from sea level to the highest human habitation in the Andes in acclimatized persons. Thorax 2018;73(8):776-778. DOI: 10.1136/thoraxjnl-2017-210598.

Footnotes

- A. Pulse oximetry (SpO₂) may underestimate arterial saturation (SaO₂), especially in people with dark skin pigmentation, thick skin, cold body temperature, or who are wearing colored nail polish, and thus may not accurately detect hypoxemia.⁽²⁾ Where appropriate, pulse oximetry should be measured after warming fingers or removing nail polish. Pulse oximetry results should be assessed within the context of the patient's clinical presentation, and performance of an arterial blood

gas considered for a more accurate assessment of oxygenation if indicated.(2) Specific target values may require adjustment when care is delivered at high altitude.(3)

- B. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Hypoxemia absent

- Hypoxemia absent, as indicated by **1 or more** of the following(1):
 - Arterial oxygen saturation (SpO₂) of 90% or greater or arterial partial pressure of oxygen (PaO₂) of 60 mm Hg (8.0 kPa) or greater on room air[A][B]
 - Oxygen requirements are stable and arranged for at next level of care.

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Rojas-Camayo J, et al. Reference values for oxygen saturation from sea level to the highest human habitation in the Andes in acclimatised persons. Thorax 2018;73(8):776-778. DOI: 10.1136/thoraxjnl-2017-210598.

Footnotes

- A. Specific target values (ie, 90% or 60 mm Hg (8.0 kPa)) may require adjustment when care is delivered at high altitude.(2)
- B. Criteria based upon clinician-acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Observation care

- When a Clinical Indication says "...despite (or beyond) observation care," the intent is to signify that, despite an attempt to address a sign, symptom, finding, or treatment need in observation care, a continued need for hospital-based care persists beyond the observation care time frame (ie, Observation Care Discharge Criteria are not met). The time frame of observation care is determined by regulation (eg, CMS's Two-Midnight Rule) or payer-provider contractual agreement. MCG's understanding of observation care is that it must be finite and of short duration. The Two-Midnight Rule for Medicare patients (inpatient is defined as medical necessity for hospital-based care across 2 or more midnights) provides a good example of what is meant by a short duration (ie, observation care should not continue beyond the second midnight).

Orthostatic hypotension

- Orthostatic hypotension,[A][B] as indicated by **1 or more** of the following(1)(2)(3):
 - Fall in SBP of 20 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position
 - Fall in DBP of 10 mm Hg or more 1 to 3 minutes after patient sits or stands from recumbent position

References

1. Shibao C, Lipsitz LA, Biaggioni I, American Society of Hypertension Writing Group. Evaluation and treatment of orthostatic hypotension. *Journal of the American Society of Hypertension* 2013 Jul-Aug;7(4):317-324. DOI: 10.1016/j.jash.2013.04.006.
2. Dalal AS, Van Hare GF. Syncope. In: Kliegman RM, et al., editors. *Nelson Textbook of Pediatrics*. 22nd ed. Elsevier; 2025:592-599.
3. Fang JC, O'Gara PT. History and physical examination: an evidence-based approach. In: Libby P, Bonow RO, Mann DL, Tomaselli GF, Bhatt DL, Solomon SD, editors. *Braunwald's Heart Disease: A Textbook of Cardiovascular Medicine*. 12th ed. Elsevier; 2022:123-140.

Footnotes

- A. Concomitant measurements of the heart rate are important to measure to help diagnose subtypes of orthostatic hypotension (eg, the lack of a compensatory increase in heart rate is typical of autonomic failure and an exaggerated tachycardia may be reflective of volume depletion). However, the heart rate is not a component of the definition of orthostatic hypotension, which relies upon blood pressure alone.(1)(2)(3)
- B. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

Pediatric Readmission Risk Assessment

- Risk of readmission is increased by presence (upon admission or during hospital stay) of **1 or more** of the following(1):
 - Immunodeficiency (eg, neutropenia)(1)
 - Malnutrition[A](1)
 - Need for parenteral nutrition or supplemental enteral tube feeding (eg, via gastrostomy tube or jejunostomy tube)(1)
 - Need for transfusion of platelets or red blood cells(1)
 - Parenteral antineoplastic (eg, chemotherapy) administered(1)
 - Presence of complex chronic condition (eg, cerebral palsy, congenital malformations, muscular dystrophy, cystic fibrosis, autoimmune disease, malignancy, genetic disorder)[B](1)(4)(5)(6)
 - Potential barriers to postdischarge clinical care (eg, unstable home environment, ability of caregivers to adhere to treatment plan unclear, impaired access to care or medication, active substance abuse in patient or caregiver)(1)(6)
 - Inpatient admission within previous 6 months(7)
 - Two or more emergency department visits within previous 6 months(1)(6)(8)

References

1. Proprietary health insurance data sources, 2022; and Medicare 100% Standard Analytical File, 2022.
2. Diab LK, Gilley SP, Krebs NF. Malnutrition in high-resource settings. In: Kliegman RM, et al., editors. *Nelson Textbook of Pediatrics*. 22nd ed. Elsevier; 2025:433-437.e1.
3. Feinstein JA, Hall M, Davidson A, Feudtner C. Pediatric complex chronic condition system version 3. *JAMA Network Open* 2024;7(7):Online. DOI: 10.1001/jamanetworkopen.2024.20579.
4. Bucholz EM, Toomey SL, Schuster MA. Trends in pediatric hospitalizations and readmissions: 2010-2016. *Pediatrics* 2019;143(2):Online. DOI: 10.1542/peds.2018-1958.
5. Toomey SL, et al. Potentially preventable 30-day hospital readmissions at a children's hospital. *Pediatrics* 2016;138(2):Online. DOI: 10.1542/peds.2015-4182.
6. Ehwerhemuepha L, Finn S, Rothman M, Rakovski C, Feaster W. A novel model for enhanced prediction and understanding of unplanned 30-day pediatric readmission. *Hospital Pediatrics* 2018;8(9):578-587. DOI: 10.1542/hpeds.2017-0220.

7. Carroll AR, et al. Validation of 30-day pediatric hospital readmission risk prediction models. JAMA Network Open 2025;8(2):e2459684. DOI: 10.1001/jamanetworkopen.2024.59684.
8. Goodman DM, et al. Development and validation of an integrated suite of prediction models for all-cause 30-day readmissions of children and adolescents aged 0 to 18 years. JAMA Network Open 2022;5(11):Online. DOI: 10.1001/jamanetworkopen.2022.41513.

Footnotes

- A. In children, malnutrition can be assessed by inadequate weight for corrected age, failure to gain adequate weight over time (weight gain velocity), height velocity, weight-for-length or BMI depending on age, and developmental outcomes.(2) These growth and anthropometric parameters are measured longitudinally and plotted on growth charts appropriate for the child's sex and age (corrected if premature).(2) The Centers for Disease Control and Prevention (CDC) recommends that tables developed by the World Health Organization be used in children up to 2 years of age and that CDC-developed tables be used for children age 2 years or older.(2) Both sets can be obtained and downloaded from www.cdc.gov/growthcharts/who_charts.htm#The%20WHO%20Growth%20Charts.
- B. A complex chronic condition is any medical condition that is expected to last at least 12 months (unless death intervenes) and to involve either several different organ systems or one organ system severely enough to require specialty pediatric care.(3)

Respiratory distress

- Respiratory distress, as indicated by **ALL** of the following(1)(2)(3)(4):
 - Patient with **1 or more** of the following:
 - Dyspnea (difficulty breathing)
 - Tachypnea
 - Abnormal breathing pattern (eg, chest retractions)
 - Other evidence of difficulty breathing
 - Evidence of respiratory compromise, as indicated by **1 or more** of the following:
 - Hypoxemia
 - Altered mental status
 - Other evidence of respiratory compromise (eg, respiratory acidosis)

References

1. Braithwaite SA, Wessel AL. Dyspnea. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:194-201.e2.
2. Naureckas ET, Solway J. Disturbances of respiratory function. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:2133-2139.
3. Matthay MA, Ware LB. Acute respiratory failure. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:642-652.
4. Rakoczy K. Acute respiratory failure. In: Gershel JC, Rauch DA, editors. Caring for the Hospitalized Child: A Handbook of Inpatient Pediatrics. 3rd ed. Elk Grove Village, IL: American Academy of Pediatrics; 2023:749-754.

Social Determinants of Health Assessment

- Risk of poor health outcomes may be increased by the presence of **1 or more** of the following social determinants of health(1)(2)(3):
 - Housing insecurity, as indicated by **1 or more** of the following:
 - Individual or caregiver's current living situation is **1 or more** of the following(4):
 - Does not have own housing (eg, staying in a hotel, shelter, or with others)

- Has own housing (eg, house, apartment), but at risk of losing it in the future (ie, behind on rent or mortgage)
- Has own housing (eg, house, apartment), but has lived in 3 or more places in past year
- Current housing has **1 or more** of the following:
 - Electrical appliances (eg, stove, refrigerator) not working or unavailable
 - Insufficient heating or cooling
 - Insufficient ventilation
 - Lead paint or pipes
 - Mold
 - Pests (eg, bugs) or rodents
 - Smoke detectors not working or unavailable
- Food insecurity, as indicated by **1 or more** of the following(5):
 - In the past year, individual or caregiver ran out of food and did not have money to buy more food.
 - In the past year, individual or caregiver worried that they would run out of food before they received money to buy more food.
- Insufficient transportation, as indicated by **1 or more** of the following(6):
 - In the past year, individual or caregiver missed medical appointments or could not get medications due to lack of transportation.
 - In the past year, individual or caregiver missed nonmedical activities, work, or could not get things needed for daily living due to lack of transportation.
- Insufficient utilities, as indicated by **1 or more** of the following(7):
 - Utilities (eg, electricity, water, gas, or oil) are currently shut off or unavailable.
 - In the past year, electric, water, gas, or oil company threatened to shut off services.
- Personal safety risk, as indicated by **2 or more** of the following(5):
 - Individual is sometimes or frequently physically hurt by another person (including family member).
 - Individual is sometimes or frequently insulted or talked down to by another person (including family member).
 - Individual is sometimes or frequently threatened with physical harm by another person (including family member).
 - Individual is sometimes or frequently screamed or cursed at by another person (including family member).
- Insufficient dependent care, as indicated by **1 or more** of the following:
 - In the past year, individual or caregiver was unable to work due to lack of dependent care.
 - In the past year, individual or caregiver was unable to work more (additional) hours due to lack of dependent care.
 - In the past year, individual or caregiver missed medical appointments or could not get medications due to lack of dependent care.
 - In the past year, individual or caregiver missed nonmedical activities (eg, school, church, social activity) due to lack of dependent care.
- Depression risk, as indicated by **ALL** of the following(8):
 - In the past 2 weeks, individual had little interest or pleasure in normal activities on at least several days.
 - In the past 2 weeks, individual felt down, depressed, or hopeless on at least several days.

References

1. Scanlon A, Reinisch C. Social determinants of health. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:18-20.
2. Billioux A, Verlander K, Anthony S, Alley D. Standardized Screening for Health-Related Social Needs in Clinical Settings: the Accountable Health Communities Screening Tool. [Internet] National Academy of Sciences. 2017 May Accessed at: <https://nam.edu/>. [accessed 2025 Sep 12]
3. Addressing social determinants of health. In: Perez R, editor. CMSA's Integrated Case Management. 2nd ed. Springer Publishing Company LLC; 2023:62-75.
4. Sandel M, et al. Unstable housing and caregiver and child health in renter families. Pediatrics 2018;14(2):e20172199. DOI: 10.1542/peds.2017-2199.
5. Children's HealthWatch Survey. Screening Instrument [Internet] Children's HealthWatch. 2024 Feb 13 Accessed at: <https://childrenshealthwatch.org/>. [accessed 2025 Mar 31]

6. PRAPARE®: Protocol for Responding to and Assessing Patient Assets, Risks, and Experiences Screening Tool. 2.0 [Internet] Association of Asian Pacific Community Health Organizations (AAPCHO) and National Association of Community Health Centers (NACHC). 2025 Accessed at: <https://prapare.org/the-prapare-screening-tool/>. [accessed 2025 Sep 08]
7. Cook JT, et al. A brief indicator of household energy security: associations with food security, child health, and child development in US infants and toddlers. *Pediatrics* 2008;122(4):e867-75. DOI: 10.1542/peds.2008-0286.
8. Kroenke K, Spitzer RL, Williams JB. The Patient Health Questionnaire-2: validity of a two-item depression screener. *Medical Care* 2003;41(11):1284-92. DOI: 10.1097/01.MLR.0000093487.78664.3C.

Tachycardia

- Tachycardia,[A][B] as indicated by **1 or more** of the following:
 - Heart rate greater than 100 beats per minute in adult[A][B](1)
 - Heart rate greater than 85 beats per minute in child 13 to 17 years of age[A][C](2)
 - Heart rate greater than 95 beats per minute in child 6 to 12 years of age[A][C](2)
 - Heart rate greater than 110 beats per minute in child 1 to 5 years of age[A][C](2)
 - Heart rate greater than 120 beats per minute in infant 3 to 11 months of age[A][C](2)
 - Heart rate greater than 150 beats per minute in infant 1 or 2 months of age[A][C](2)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. *Goldman-Cecil Medicine*. 27th ed. Elsevier; 2024:32-35.
2. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. *The Harriet Lane Handbook: A Manual for Pediatric House Officers*. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. For example, an elderly patient on a beta-blocker medication with a baseline resting heart rate of 60 beats per minute may be clinically tachycardic at a heart rate of 94 beats per minute. Likewise, a patient who is upset, in pain, or nervous in the emergency department with a heart rate of 106 beats per minute may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as chest pain or hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.
- C. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. A patient who is upset, in pain, or nervous in the emergency department with an elevated heart rate may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Tachycardia absent

- Tachycardia[A][B] absent, as indicated by **1 or more** of the following:
 - Heart rate less than or equal to 100 beats per minute in adult[A][B](1)
 - Heart rate less than or equal to 85 beats per minute in child 13 to 17 years of age[A][B](2)
 - Heart rate less than or equal to 95 beats per minute in child 6 to 12 years of age[A][B](2)
 - Heart rate less than or equal to 110 beats per minute in child 1 to 5 years of age[A][B](2)
 - Heart rate less than or equal to 120 beats per minute in infant 3 to 11 months of age[A][B](2)
 - Heart rate less than or equal to 150 beats per minute in infant 1 or 2 months of age[A][B](2)

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of heart rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline heart rate, medications, and clinical impact. For example, an elderly patient on a beta-blocker medication with a baseline resting heart rate of 60 beats per minute may be clinically tachycardic at a heart rate of 94 beats per minute. Likewise, a patient who is upset, in pain, or nervous in the emergency department with a heart rate of 106 beats per minute may meet the technical definition of tachycardia, but this tachycardia (absent associated findings such as chest pain or hypotension) may not be clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachycardia. Whether a heart rate above or below the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment.

Tachypnea

- Tachypnea,[A][B] as indicated by **1 or more** of the following:
 - Respiratory rate greater than 20 breaths per minute in adult[A][B](1)
 - Respiratory rate greater than 18 breaths per minute in child 13 to 17 years of age[A][B](2)
 - Respiratory rate greater than 22 breaths per minute in child 6 to 12 years of age[A][B](2)
 - Respiratory rate greater than 25 breaths per minute in child 3 to 5 years of age[A][B](2)
 - Respiratory rate greater than 30 breaths per minute in child 1 or 2 years of age[A][B](2)
 - Respiratory rate greater than 40 breaths per minute in infant 6 to 11 months of age[A][B](2)
 - Respiratory rate greater than 45 breaths per minute in infant 3 to 5 months of age[A][B](2)
 - Respiratory rate greater than 55 breaths per minute in infant 1 or 2 months of age[A][B](2)

References

1. Schrager DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.

2. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.
- B. Interpretation of respiratory rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline respiratory rate and whether the respiratory rate is indicative of significant underlying respiratory or other pathology. A mildly increased respiratory rate (eg, 20 to 22 breaths per minute in an adult), absent other indications of serious illness (eg, hypoxemia, metabolic acidosis, decreased tidal volume, pain), may be transitory and not clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachypnea. Whether a respiratory rate above the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment. In addition, the measurement of the respiratory rate is subject to error. A medical textbook states that because there can be significant breath-to-breath variation in respiratory rate, the rate should be assessed for at least 30 seconds, preferably for a full minute.⁽¹⁾ This same textbook also states that new technologies designed to measure the respiratory rate have not proved to be clinically useful.⁽¹⁾

Tachypnea absent

- Tachypnea^{[A][B]} absent, as indicated by **1 or more** of the following:
 - Respiratory rate less than or equal to 20 breaths per minute in adult^{[A][B]}(1)
 - Respiratory rate less than or equal to 18 breaths per minute in child 13 to 17 years of age^{[A][B]}(2)
 - Respiratory rate less than or equal to 22 breaths per minute in child 6 to 12 years of age^{[A][B]}(2)
 - Respiratory rate less than or equal to 25 breaths per minute in child 3 to 5 years of age^{[A][B]}(2)
 - Respiratory rate less than or equal to 30 breaths per minute in child 1 or 2 years of age^{[A][B]}(2)
 - Respiratory rate less than or equal to 40 breaths per minute in infant 6 to 11 months of age^{[A][B]}(2)
 - Respiratory rate less than or equal to 45 breaths per minute in infant 3 to 5 months of age^{[A][B]}(2)
 - Respiratory rate less than or equal to 55 breaths per minute in infant 1 or 2 months of age^{[A][B]}(2)

References

1. Schriger DL. Approach to the patient with abnormal vital signs. In: Goldman L, Cooney KA, editors. Goldman-Cecil Medicine. 27th ed. Elsevier; 2024:32-35.
2. Anderson CC, Kapoor S, Mark TE. Pediatric parameters and equipment. In: Anderson CC, Kapoor S, Mark TE, editors. The Harriet Lane Handbook: A Manual for Pediatric House Officers. 23rd ed. Elsevier; 2024:i-iii.

Footnotes

- A. Criteria based upon clinician acquired numeric values (eg, vital signs, oxygen saturation) should be used if they are accurate reflections of the patient's condition. Transitory findings (eg, abnormal only upon initial emergency department intake or only one time out of multiple readings) that rapidly improve with no or minimal treatment usually do not reflect disease severity or risk for deterioration. This does not imply that an initial or one-time reading cannot ever be applicable. The goal is to separate erroneous or incidental findings from those that truly represent the patient's clinical picture.

B. Interpretation of respiratory rate requires clinical judgment and consideration of several patient-specific factors, such as the patient's baseline respiratory rate and whether the respiratory rate is indicative of significant underlying respiratory or other pathology. A mildly increased respiratory rate (eg, 20 to 22 breaths per minute in an adult), absent other indications of serious illness (eg, hypoxemia, metabolic acidosis, decreased tidal volume, pain), may be transitory and not clinically important. The numeric values included in this definition are provided to allow for consistency in terms of a technical definition of the term tachypnea. Whether a respiratory rate above the technical threshold is clinically meaningful is a matter of persistence, context, and clinical judgment. In addition, the measurement of the respiratory rate is subject to error. A medical textbook states that because there can be significant breath-to-breath variation in respiratory rate, the rate should be assessed for at least 30 seconds, preferably for a full minute.⁽¹⁾ This same textbook also states that new technologies designed to measure the respiratory rate have not proved to be clinically useful.⁽¹⁾

Codes

ICD-10 Diagnosis: E10.21, E10.22, E10.29, E11.21, E11.22, E11.29, E13.21, E13.22, E13.29, N17.0, N17.1, N17.2, N17.8, N17.9, N99.0, O03.32, O03.82, O04.82, O07.32, O08.4, O90.49, T79.5XXA [Hide]

ICD-10 Procedure: 3E1M39Z

CPT®: 90935, 90937, 90945, 90947, 90999

CPT copyright 2025 American Medical Association. All rights reserved.

MCG Health
Inpatient & Surgical Care 30th Edition
Copyright © 2026 MCG Health, LLC
All Rights Reserved

Last Update: 1/25/2026 6:08:23 AM
Build Number: 30.0.2026012500524.025256