

Recovery Facility Care > Optimal Recovery Guidelines > Inpatient Rehabilitation Facility (IRF) > Cardiology > Inpatient Rehabilitation Facility (Acute Rehabilitation): Cardiac Disorders (I-7018)

Inpatient Rehabilitation Facility (Acute Rehabilitation): Cardiac Disorders

IRF: I-7018 (RFC)

[Link to Codes](#)

MCG Health
Recovery Facility Care
30th Edition

- Clinical Indications for Admission to Inpatient Rehabilitation Facility (Acute Rehabilitation)
- Alternative Levels of Care
- Length of Stay
- Evaluation and Treatment
 - General Treatment Course
 - Extended Stay
 - Rehabilitation Treatment Plan
 - Physical Therapy Treatment Plan
 - Occupational Therapy Treatment Plan
 - Patient Education
- Discharge Planning
- Quality Measures
- Evidence Summary
 - Criteria
 - Length of Stay
 - Treatment
 - Care Coordination
 - Rationale
 - Related CMS Coverage Guidance
- References
- Footnotes
- Definitions
- Codes

Clinical Indications for Admission to Inpatient Rehabilitation Facility (Acute Rehabilitation)

- **Inpatient rehabilitation facility** (acute rehabilitation) admission may be indicated by **ALL** of the following^{[A](1)(2)(3):N}
 - No acute hospital care needs
 - Preadmission screening completed; assessment examples include^{[B][C](1):}

- Level of function prior to the event
- Condition that caused the need for rehabilitation
- Expected level of improvement
- Expected length of time to achieve level of improvement
- Risk for clinical complications
- Combination of treatments needed (PT, OT, SLP, or orthosis/prosthesis)
- Anticipated discharge destination
- Patient status supports **active** participation in, and benefit from, intensive rehabilitation Skilled care that includes **ALL** of the following[D](3)(8):
 - Intensity and complexity of nursing, medical management, and rehabilitation needs require and can be expected to benefit from an inpatient stay and interdisciplinary team approach.[I]
 - Ability to actively participate in Intensive rehabilitation therapy program
 - Patient can reasonably be expected to demonstrate Goals for measurable improvement.
 - Requires physician supervision to assess the patient both medically and functionally[E]
 - Requires an intensive and coordinated Interdisciplinary team care approach[F]
- Therapy services needed, including **ALL** of the following(10)(11):[I]
 - Skilled therapy services needed for **1 or more** of the following:
 - Cardiac rehabilitation evaluation and training(14)
 - Edema-management techniques(14)
 - Environmental and activity modification for ADL performance(15)
 - Equipment and adaptive technology use and safety
 - Fall-prevention and recovery training(16)
 - Functional activities evaluation and training(17)
 - Gait evaluation and training(18)
 - Motor control and balance evaluation and training(14)
 - Neurobehavioral and neurocognitive assessment and treatment strategies (eg, learning, memory, attention, problem-solving, abstract reasoning, and impulsivity)(19)
 - Orthosis or prosthesis evaluation and training
 - Pain-management techniques(20)
 - Positioning techniques and training
 - Range of motion evaluation and training
 - Safety awareness training related to functional mobility and ADL performance(19)
 - Strength evaluation and training(21)
 - Therapeutic exercises or activities supervision to ensure patient safety and treatment effectiveness(21)
 - Multidisciplinary therapy services needed, including **2 or more** of the following(22):
 - Physical therapy (PT)
 - Occupational therapy (OT)
 - Orthosis or prosthesis
- Nursing services needed, including **1 or more** of the following(8)(23):
 - Behavioral health condition monitoring(19)
 - Bowel and bladder management(24)(25)
 - Infusion therapy management(14)
 - Medical condition monitoring with evaluation of rehabilitation on clinical status(26)
 - Musculoskeletal management(27)

- Nutritional management(21)
- Respiratory management(14)
- Therapy skills incorporated and reinforced in nursing care
- Time-tabling and routine establishment (eg, for ADL adaptation)
- Pain management(20)
- Patient and caregiver education (eg, safety, skin care, medication)(28)
- Wound management(14)(26)

Alternative Levels of Care

- Patient may be candidate for other levels of care, including:
 - Subacute or skilled care. See:
 - Angina or Chest Pain [RFC](#)
 - Arrhythmia Disorders [RFC](#)
 - Heart Failure [RFC](#)
 - Myocardial Infarction [RFC](#)
 - Home care. For admission criteria, see:
 - Angina [HC](#)
 - Atrial Fibrillation [HC](#)
 - Heart Failure [HC](#)
 - Myocardial Infarction [HC](#)
 - Supraventricular Arrhythmia [HC](#)
 - Ventricular Arrhythmia [HC](#)

Length of Stay

Length of stay is displayed as percentiles that are based on the observed utilization of inpatient rehabilitation facility length of stay for this diagnosis. For guidelines with low claims data volume, length of stay statistics that are displayed are from combined categories (eg, major postoperative and major medical). Individual patients may require shorter or longer stays as appropriate for their clinical status and care needs. This table is designed to allow organizations to define their goals for inpatient rehabilitation utilization by choosing from the displayed range.

Inpatient Rehabilitation Facility Commercial Length of Stay					Inpatient Rehabilitation Facility Medicare Length of Stay				
10%ile	20%ile	30%ile	40%ile	50%ile	10%ile	20%ile	30%ile	40%ile	50%ile
6	7	8	10	11	7	8	9	10	11

Evaluation and Treatment

General Treatment Course

Stage	Clinical Status	Interventions
-------	-----------------	---------------

1

- **Clinical indications for admission met**
- Inpatient Rehabilitation Admission Assessment complete with intensive rehabilitation needs and services identified and evaluation of current level of functioning (eg, Inpatient Rehabilitation Facility Patient Assessment Instrument (IRF-PAI)) **QM** [G][H]
- Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (IADL) Assessment  SR complete  Provide treatments and monitoring:
 - Behavioral management(37)
 - Bladder management
 - Bowel management
 - Cardiac care(14)
 - Chronic condition management as indicated
 - Cognitive management techniques(14)
 - Fall risk management **QM** (38)
 - Fluid and electrolyte management(39)
 - Infusion (IV or subcutaneous) therapy management (eg, fluids, antibiotics, parenteral nutrition)(40)
 - Laboratory testing or monitoring, if ordered(39)
 - Medication management **QM** (41)
 - Nutrition management(42)
 - Pain management
 - Pressure injury management **QM** (43)
 - Respiratory care
 - Thromboprophylaxis
 - Tracheostomy management
 - Wound or dressing management
- for evaluation of current level of functioning (eg,  Activities of Daily Living (ADL) Scoring Tool Calculator) **QM**
- Physical ability assessment (eg, FIM Score®, Functional Independence Measure for children (WeeFIM®))
- Pressure injury assessment (ie, risk for new or presence of existing pressure injury). See Wounds Due to Surgery, Injury, or Pressure  SR for further information. **QM**
- Psychosocial status assessment. See Psychosocial Assessment  SR for further information. **QM**
- Social determinants of health screening. See Social Determinants of Health Screening Tool  SR for further information. **QM**(32)
- Hospital readmission risk factor evaluation **QM** (33)(34)
- Transition of care planning initiated with evaluation for next level of care **QM** (35)
- Interdisciplinary care plan established and implemented **QM** [I](3)(36)
- Education for self-care
- Physical therapy (PT) evaluation
- Occupational therapy (OT) evaluation
- Neuropsychological (eg, neurocognitive and neurobehavioral) evaluation, if indicated(19)
- Orthosis or prosthesis evaluation, if indicated(11)
- Social work or case management evaluation(1)(3)(44)

2

- **Intensive rehabilitation therapy program established and patient actively participating**[J]
- Unexpected clinical event resolved (eg, infection, DVT, pneumonia)[K]
- Interdisciplinary care plan continues **QM**
- Ongoing assessment of clinical needs **QM**

- Patient demonstrating measurable practical improvement in functional condition with evaluation of current level of functioning (eg,  Activities of Daily Living (ADL) Scoring Tool Calculator) **QM**(45)
 - **Physical therapy (PT) progression**
 - **Occupational therapy (OT) progression**
 - **Other care needs stable or diminished**
- ☐ Provide treatments and monitoring:
- Behavioral management
 - Bladder management
 - Bowel management
 - Cardiac care
 - Chronic condition management as indicated
 - Cognitive management techniques
 - Fall risk management **QM**
 - Fluid and electrolyte management
 - Infusion (IV or subcutaneous) therapy management (eg, fluids, antibiotics, parenteral nutrition)(40)
 - Laboratory testing or monitoring, if ordered
 - Medication management **QM** (41)
 - Nutrition management
 - Pain management
 - Pressure injury management **QM**
 - Respiratory care
 - Thromboprophylaxis
 - Tracheostomy management
 - Wound or dressing management
 - Ongoing patient or caregiver education
 - Ongoing equipment (eg, DME) training
 - Ongoing rehabilitation therapy(3)
 - Ongoing neuropsychological management, if indicated
 - Orthosis or prosthesis training, if indicated
 - Transition of care planning continues with evaluation for next level of care. **QM** (46)

3

- Adequate intensive inpatient rehabilitation completed for safe transfer to next level of care, or patient's condition prevents active participation in intensive therapy program, or patient no longer demonstrating significant functional gains (eg,  Activities of Daily Living (ADL) Scoring Tool Calculator)
- Bowel and bladder management training completed
- Communication technique training completed
- Enteral or parenteral nutrition tolerated, or oral diet
- Equipment (eg, DME) training completed
- Gastrointestinal status stable or manageable at lower level of care
- Genitourinary status stable or manageable at lower level of care
- **Transition of care planning completed QM**
- Safe to go home and transition to community or transition to alternative level of care (eg, skilled nursing facility, home care) **QM**

- **Medical status stable or manageable at lower level of care (eg, vital signs, laboratory values)**
- **Neurobehavioral and neurocognitive status manageable at lower level of care**
- **Neurovascular status stable or manageable at lower level of care**
- **Orthosis or prosthesis training completed or manageable at lower level of care**
- **Pain controlled or manageable at lower level of care**
- Patient and caregiver education completed or manageable at lower level of care
- Patient transitioned to another level of care due to unexpected clinical event (eg, DVT, infection)
- **Respiratory equipment and care properly demonstrated**
- **Respiratory status stable or manageable at lower level of care**
- **Wound(s) or dressing changes manageable at lower level of care**
- **Transition plans and education understood**



Recovery Milestones are indicated in **bold**.

Extended Stay

Extended recovery facility stay may be: brief (1 to 3 days), moderate (4 to 7 days), or prolonged (more than 7 days).

- Extended stay beyond goal length of stay may be indicated when **ALL** of the following exist:
 - No acute hospital care needs
 - Patient able to make measurable improvement in functional condition in predetermined and reasonable period of time
 - Intensity of medical and rehabilitation management necessitates inpatient stay.
 - Coordinated interdisciplinary team approach required to manage intensity of medical and rehabilitation needs
- ☐ Therapy services required, including **2 or more** of the following:
 - Physical therapy (PT)
 - Occupational therapy (OT)
 - Orthosis or prosthesis
- Patient has had an unexpected clinical event and Skilled care needed on daily basis for **1 or more** of the following^{[D][K]}:
 - ☐ Deep venous thrombosis (DVT) or pulmonary embolus (PE) management; examples include(48):
 - Pharmacoprophylaxis initiation(49)
 - Hemodynamic measurements outside normal limits
 - Laboratory values outside normal limits.
 - Signs of active bleeding
 - Expect brief stay extension.
 - ☐ Fall management; examples include(50)(51):
 - Functional status impaired, impacting ability to actively participate in intensive rehabilitation program
 - Evaluation and treatment of fall injury, injury regimen established within 3 days of fall, and intensive rehabilitation program resumed(3)
 - Expect brief stay extension.
 - ☐ Fever management; examples include(52):
 - Temperature status outside prescribed limits

- Laboratory values outside normal limits(53)
 - Hemodynamic measurements outside normal limits
 - Behavioral symptoms (eg, agitation, somnolence, or inappropriate behavior) that interfere with medical care
 - Expect brief stay extension.
- ☐ Infusion therapy; examples include(3)(40):
- Administration of agent (eg, chemotherapy, new intravenous medication) that requires clinical management and may require brief break in intensive therapy rehabilitation program(54)
 - Expect brief stay extension.
- ☐ Malnutrition, dehydration, or electrolyte management; examples include:
- Oral nutrition and hydration not tolerated and enteral or parenteral nutrition implemented
 - Enteral or parenteral nutrition complications present(55)
 - Transition from enteral or parenteral nutrition to oral nutrition(56)
 - Hemodynamic measurements outside normal limits
 - Laboratory values outside normal limits(57)
 - Urinary output below prescribed parameters(57)
 - Expect brief stay extension.
- ☐ Mental status changes; examples include(58):
- Danger to self or others
 - Behavior crisis management (eg, physical or chemical restraints) needed(59)
 - Behavioral symptoms (eg, agitation, somnolence, or inappropriate behavior) that interfere with medical care
 - Sudden disturbances in consciousness, changes in cognition (eg, disorientation, memory loss, or language impairment), or disturbances in thought that prevent intensive therapy services
 - Laboratory values outside normal limits
 - Expect brief stay extension.
- ☐ Pain evaluation and management; examples include(60)(61)(62):
- Frequent pain assessment using validated diagnostic tool[L]
 - Early referral and treatment of pain
 - Ongoing education and training of pain treatment plan (eg, multimodal pain management)[M][N]
 - Pain medication side effects or risk factors for adverse effects (eg, constipation, sedation, falls)
 - Expect brief to moderate stay extension.
- ☐ Respiratory management; examples include(64)(65):
- Breathing abnormalities and oxygen regimen not established
 - Laboratory values outside normal limits
 - Respiratory status assessment to evaluate patient's response to treatment (eg, suctioning or pulmonary hygiene that requires skilled intervention or monitoring)
 - Assessment or care unmanageable at lower level of care
 - Expect brief stay extension.
- ☐ Seizure management(66)(67)
- Seizure management with assessment of clinical status, monitoring for complications, administering medication and evaluating effect, airway management, and injury prevention
 - Expect brief stay extension.

- ☐ Significant comorbid disease exacerbation with need for skilled clinical intervention and monitoring; examples include(3)(68):
 - Hyperglycemia or hypoglycemia
 - Hypertension or hypotension
 - Inadequate pain management
 - Condition assessment or care unmanageable at lower level of care
 - Expect brief extension stay.
- ☐ Tracheostomy evaluation and equipment management; examples include(69)(70)(71):
 - Tracheostomy stoma with signs and symptoms of complications (eg, infection)
 - Tracheostomy care (eg, suctioning, pulmonary toilet, assessment for instability) requires skilled intervention or monitoring.
 - Frequent suctioning needed (eg, increased secretions)
 - Laboratory values outside normal limits; examples include:
 - Oxygen saturation below 90% (or below baseline)
 - Cultures positive or identification of infective source pending, and treatment regimen not established
 - Ongoing education and training on tracheostomy management needed
 - Assessment or care unmanageable at lower level of care
 - Expect brief stay extension.
- ☐ Urinary management; examples include(72)(73):
 - Urinary catheter or percutaneous suprapubic tube complications
 - Urinary output inadequate for medical condition(74)
 - Laboratory values outside normal limits(39)(75)
 - Expect brief stay extension.
- ☐ Ventilator evaluation and equipment management; examples include(53):
 - Mechanical ventilation unstable; examples include:
 - Breathing abnormalities exist and oxygen regimen not established
 - Respiratory status unstable for 48 hours after discontinuation of mechanical ventilation
 - Laboratory values outside normal limits; examples include:
 - Oxygen saturation below 90% (or below baseline)(64)
 - Cultures positive or identification of infective source pending, and treatment regimen not established(76)
 - Respiratory status assessment to evaluate patient's response to treatment (eg, suctioning, ventilator setting changes)(64)
 - Continued education and training for home ventilator and respiratory management programs needed(77)
 - Assessment or care unmanageable at lower level of care
 - Expect brief stay extension.
- ☐ Wound management; examples include(78):
 - Wound status changed with new type of dressing or procedure
 - New wound developed (eg, pressure injury)
 - Healing impeded due to comorbidities (eg, diabetes, malignancy)
 - Nutrition or hydration inadequate for normal duration of healing
 - Expect brief stay extension.

Rehabilitation Treatment Plan

Physical Therapy Treatment Plan

- **Physical therapy** treatment plan includes(79):

[-] Goals of physical therapy

- Activity tolerance and functional endurance improved
- ADL/IADL performance, independence, and safety optimized
- ADL/IADL restrictions, precautions, or activity guidelines understood
- Awareness of community resources optimized
- Chronic condition modifiable risk factor management strategies maximized(80)
- Durable medical equipment usage optimized
- Energy conservation techniques and proper body mechanics maximized
- Exercise tolerance and functional endurance improved
- Fall-risk-reduction strategies maximized
- Functional independence and safety with mobility-related tasks maximized
- Home rehabilitation program and recommendations understood
- Orthosis or prosthesis management and independence maximized
- Pain-management techniques optimized
- Potential personality or emotional changes understood
- Pressure injury and contracture risk prevention maximized
- Respiratory status and breathing techniques improved
- Risk and disability associated with disease, injury, or surgical procedure understood
- Self-management of health, injury, surgical procedure, or disease process optimized
- Self-reported quality of life improved
- Strength and range of motion improved
- Postdischarge complication risk factors minimized

[-] Rehabilitation techniques and training

- ADL training
- Cardiac rehabilitation(81)(82)
- Edema prevention or management(80)
- Education of therapeutic program with patient and caregiver(82)
- Energy conservation and work simplification(83)
- Equipment and device use training
- Exercise program setup and progression within precautions, which may include(10)(81)(84):
 - Aerobic exercises or endurance conditioning
 - Balance training
 - Coordination exercises
 - Range-of-motion techniques
 - Respiratory exercises and training (eg, respiratory muscle training, coughing techniques, deep breathing exercises, incentive spirometry exercises, pursed lip and diaphragmatic breathing)
 - Strengthening exercises
 - Stretching exercises
- Fall-prevention and recovery training **QM** (85)
- Functional skills training with applicable durable medical equipment (eg, walker, transfer board)
 - Bed mobility training
 - Body weight-supported gait training (eg, robot-assisted gait training)

- Gait training on level and uneven surfaces
- Stair management
- Transfer training
- Wheelchair mobility and management(84)(86)
- Lifting as prescribed for diagnosis (eg, medical, surgical)(82)
- Neuromuscular re-education (eg, proprioception training)
- Pain-management strategies and techniques(87)
- Posture, positioning, and body mechanics as prescribed for diagnosis or surgical approach(82)
- Pressure injury prevention or management(88)
- Safety awareness education(89)
- Self-management of health, injury, surgical procedure, or disease process(90)(91)
- Splint, brace, cast, prosthesis, or orthosis management(92)
- Therapeutic physical modality use and training, as indicated(93)(94)
 - Cold modalities or cryotherapy with precautions related to circulatory deficiency, areas of desensitization, open wounds, or fractures
 - Functional or neuromuscular electrical stimulation
 - Thermal modality with precautions related to circulatory deficiency, areas of desensitization, open wounds, or fractures
 - Transcutaneous electrical nerve stimulation (TENS)
 - Ultrasound therapeutic modality
 - Vasopneumatic compression
- Vestibular rehabilitation or vertigo management(95)(96)(97)(98)
- Weight-bearing as prescribed for diagnosis and/or surgical approach(92)
- ☐ Discharge recommendations, which may include(81):
 - Activity modifications
 - Environmental adaptation
 - Home exercise program as prescribed
 - Home modifications

Occupational Therapy Treatment Plan

- **Occupational therapy** treatment plan includes(99)(100)(101):
 - ☐ Goals of occupational therapy
 - Activity tolerance and functional endurance improved
 - ADL/IADL performance, independence, and safety optimized(102)
 - ADL/IADL restrictions, precautions, or activity guidelines understood
 - Balance and body alignment for ADL/IADL improved
 - Cognitive skills for ADL/IADL improved(103)(104)
 - Fall-risk-reduction strategies maximized(38)
 - Home rehabilitation program and recommendations understood(103)
 - Pain-management techniques optimized
 - Participation arousal within environment improved
 - Potential personality or emotional changes understood
 - Proper posture for sitting, ADL, and transfers maximized
 - Upper body range of motion, strength, and functional use for ADL or IADL maximized(102)
 - Postdischarge complication risk factors minimized

- [-] Rehabilitation techniques and training
 - ADL or IADL training(104)
 - Behavior-management strategies and techniques (eg, cognitive behavioral therapy, cognitive rehabilitation techniques)(105)
 - Compensatory strategies for completion of ADL/IADL(103)
 - Cognition impairment management(106)
 - Energy conservation
 - Perceptual deficits management
 - Sensory impairment management
 - Work simplification
 - Compression bandage or garment management(107)
 - Contracture prevention or management(108)
 - Edema prevention or management
 - Education of therapeutic program with patient and caregiver(103)
 - Environmental and activity modification for ADL/IADL performance(104)
 - Equipment and device use training
 - Exercise program setup and progression within precautions, which may include(103):
 - Balance training
 - Coordination exercises
 - Range-of-motion techniques
 - Strengthening exercises
 - Stretching exercises
 - Fall-prevention and recovery training **QM** (85)
 - Functional skills training with applicable durable medical equipment (eg, shower chair, commode)(104)
 - Bed mobility training
 - Transfer training
 - Joint-protection techniques and mobilization
 - Lifting as prescribed for diagnosis (eg, medical, surgical)
 - Manual therapy (eg, soft tissue mobilization techniques, scar management, massage)(109)
 - Neuromuscular re-education (eg, proprioception training)
 - Pain-management strategies and techniques
 - Posture, positioning, and body mechanics as prescribed for diagnosis or surgical approach(103)
 - Safety awareness education(102)
 - Sensory stimulation and integration techniques
 - Stress-management techniques(110)
 - Therapeutic physical modality (eg, heat or ice packs) use and training, as indicated
- [-] Discharge recommendations, which may include(102)(111):
 - Activity modifications
 - Environmental adaptation
 - Home exercise program
 - Home modifications
 - Pre-driver assessment and automobile modification
 - Vocational rehabilitation(110)

Patient Education

- Patient and caregiver education. See:
 - Angina: Patient Education for Clinicians [SR](#)
 - Atrial Fibrillation: Patient Education for Clinicians [SR](#)
 - Heart Failure: Patient Education for Clinicians [SR](#)
 - Myocardial Infarction: Patient Education for Clinicians [SR](#)
 - Supraventricular Arrhythmia: Patient Education for Clinicians [SR](#)
 - Ventricular Arrhythmia: Patient Education for Clinicians [SR](#)

Discharge Planning

- Transition of care planning may include(11):[NN](#)
 - ☐ Assessment and early establishment of anticipated discharge destination and plans for care, including(117):
 - Treatment plan development involving multiple providers
 - Premorbid functioning
 - Patient and caregiver preferences and abilities
 - Evaluation for skilled services at next level of care as appropriate for patient's continued needs
 - Psychosocial status. See Psychosocial Assessment [SR](#) for further information. **QM**(118)
 - Social determinants of health screening. See Social Determinants of Health Screening Tool [SR](#) for further information. **QM**(32)
 - Acceptance for care for patient at next level of care, as appropriate
 - Transition of care plan complete, which may include:
 - Patient and caregiver education complete. See Teach-Back Tool [SR](#) for further information.
 - ☐ Medication reconciliation completion includes **QM** (119)(120):
 - Compare patient's discharge list of medications (prescribed and over-the-counter) against provider's admission or transfer orders.
 - Assess each medication for correlation to disease state or medical condition.
 - Report medication discrepancies to prescribing provider, attending physician, and primary care provider, and ensure accurate medication order is identified.
 - Provide reconciled medication list to all treating providers.
 - Confirm that patient or caregiver can acquire medication.
 - Educate patient and caregiver.
 - Provide complete medication list to patient and caregiver.
 - Importance of presenting personal medication list to all providers at each care transition, including all provider appointments
 - Reason, dosage, and timing of medication (eg, use "teach-back" techniques)(121)(122)
 - Encourage communication between patient, caregiver, and pharmacy for obtaining prescriptions, setting up home medication delivery, and reviewing for drug-drug interactions.
 - See Medication Reconciliation Tool [SR](#) for further information.
 - ☐ Appointments planned or scheduled(46):
 - Primary care provider
 - Cardiologist

- Dietitian
 - Outpatient imaging, tests, or procedures
 - Physiatrist or rehabilitation provider
 - Rehabilitation program (eg, cardiac, pulmonary) at next level of care
 - Rehabilitation therapy (eg, PT) at next level of care(46)
 - Respiratory therapy
 - Specialists for management of comorbidities as needed
 - Surgeon
 - Transplant center or team
 - Wound care center
 - Wound, ostomy, and continence nurse (WOCN)
 - Other
- ☐ Referrals made for assistance or support, which may include:
- Alcohol and other drug abuse or dependence treatment, as needed **QM**
 - Behavioral health support (eg, counseling)
 - Community services(123)(124)
 - Educational program (eg, chronic condition management, self-management)(121)
 - End-stage disease planning
 - Financial, for follow-up care, medication, and transportation
 - Home architectural modifications
 - Home healthcare(125)
 - Local electric or power company regarding need for life support if ventilator needed at home
 - Protective services or legal authorities, if indicated
 - Self-help or support groups
 - Social services (eg, social programs, advance directives)
 - Tobacco use treatment, as needed **QM** (126)(127)
 - Vocational rehabilitation(110)
 - Other
- ☐ Medical equipment and supplies coordinated (ie, delivered or delivery confirmed), as indicated(128):
- ADL-supportive equipment
 - Cardiac care equipment and supplies (eg, compression stockings)
 - Drain care supplies
 - Medication equipment and supplies
 - Nutritional care equipment and supplies
 - Pediatric and neonatal equipment and supplies
 - Respiratory equipment and supplies
 - Urinary care equipment and supplies
 - Wound care equipment and supplies
 - Other
- Transition plan communicated to all members of patient's care team **QM**

Quality Measures

- HEDIS® Measures include(35):
 - Medication reconciliation is completed.
 - Transition of care needs are assessed and addressed.
 - Psychosocial status (eg, depression screening) is assessed and addressed.
 - Fall risk is assessed and addressed.
 - Tobacco use is assessed and addressed.
 - Immunization status is assessed and addressed.
 - Readmission risk is assessed and addressed.
 - Alcohol and substance misuse is assessed and addressed.
 - Social needs screening and intervention is completed.
- Inpatient Rehabilitation Facility Compare Quality Measures include(129):
 - Pressure injury risk is assessed and addressed.
 - ADL status is assessed for improvement.
 - Fall risk is assessed and addressed.
 - Transfer of health information to provider and patient is completed.
 - Healthcare-associated infection risk is reduced
 - Medication reconciliation is completed.
 - Discharge to the community is completed.
 - Readmission risk is assessed and addressed.
- Inpatient Rehabilitation Facility Quality Reporting Program measures include(129):
 - Pressure injury risk is assessed and addressed.
 - ADL status is assessed for improvement.
 - Fall risk is assessed and addressed.
 - Transfer of health information to provider and patient is completed.
 - Readmission risk is assessed and addressed.
 - Medication reconciliation is completed.
 - Immunization status is assessed and addressed.
 - Discharge to the community is completed.

Evidence Summary

Criteria

The evidence for the clinical indications found in this guideline includes 6 published peer reviewed articles, 1 specialty society or other evidence-based guideline, and 15 book sections.

A retrospective review of over 1.9 million inpatient rehabilitation patients discharged between January 2016 and December 2019 showed that 70% of patients had at least one ICD-10 pain diagnosis; pain diagnosis was common among all impairment groups, affecting half of patients in each group, with the highest prevalence in the arthritis and pain syndromes impairment group.(4) **(EG 2)** A retrospective cohort study of 3219 patients with heart failure compared inpatient cardiac rehabilitation (CR) with no CR and found that CR was associated with a 20% reduction in long-term mortality and significant improvements in functional capacity, as indicated by the mean 6-minute walking distance increasing by 63 meters (206 feet).(5) **(EG 2)**

Length of Stay

A retrospective cohort study of 116 heart and/or lung transplant recipients (49 orthotopic heart transplants, 65 lung transplants, and 2 combined heart-lung transplants) admitted to a multidisciplinary inpatient rehabilitation program from 2009 to 2016 found that the average length of stay was 26.9 days.(47) **(EG 2)**

Treatment

A retrospective study of 479 Medicare patients who received rehabilitation in a nursing home or inpatient rehabilitation facility (IRF) setting found that 1 in 3 patients reported no functional improvement during rehabilitation. Factors associated with greater odds of reporting no improvement included high school education or less, impaired instrumental activities of daily living, less than 1 month total duration of rehabilitation, and no outpatient rehabilitation services.(9) **(EG 2)**

A claims-based study of 1.4 million Medicare beneficiaries discharged from an acute care hospital to a post-acute care setting, including an inpatient rehabilitation facility (IRF), a skilled nursing facility (SNF), or home care (HC), found that more hours of occupational therapy and physical therapy services were associated with greater functional improvements, and fewer hours of occupational therapy and physical therapy services were associated with a higher risk for 30-day hospital readmission across all post-acute care settings.(12) **(EG 2)** A retrospective study of 3212 patients who were discharged from an IRF found that increased therapy hours were associated with increased functional gains among all types of patients.(13) **(EG 2)**

Care Coordination

An observational review of 567,850 inpatient rehabilitation facility (IRF) Medicare inpatient stays in 1166 facilities during 2013 to 2014 found that adjusted hospital readmission rate 30 days post-IRF discharge was in the range of 12.4% to 13.1%, but almost 75% of facilities demonstrated readmission rates that were significantly different than this average. The mean number of days to readmission was 13.0, which varied significantly by rehabilitation diagnosis.(112) **(EG 2)** A retrospective cohort study of a Medicare claims database involving 371,846 patients discharged from inpatient rehabilitation from 2013 to 2014 found that functional independence at discharge was associated with lower rates of potentially preventable hospital readmissions. Readmission rates for highest vs lowest quartiles within each functional domain were 3.4% vs 6.9% for self-care, 3.3% vs 7.2% for mobility, and 3.5% vs 6.2% for cognition; 44.1% of readmissions were for infection-related conditions, 31.2% were for inadequate management of chronic conditions, and 24.7% were for inadequate management of other unplanned events.(113) **(EG 2)** A retrospective cohort analysis of 539 geriatric patients discharged from acute care to geriatric rehabilitation found that predictors of acute readmission include severe infection, fracture secondary to falls, intra-abdominal complications, cardiac complications, and acute neurologic events.(114) **(EG 2)** An evidence-based improvement study of 79 patients discharged from a 28-bed IRF showed implementing the teach-back intervention from the Agency for Healthcare Research and Quality (AHRQ) Health Literacy Universal Precautions Toolkit into the discharge process resulted in a mean 45% fewer 30-day readmissions.(115) **(EG 2)**

A longitudinal observational study of 43 adult inpatient rehabilitation facility patients who were not safe to return home at discharge and required a skilled rehabilitation facility placement found that 58.2% of patients returned to baseline independence, and 63.4% of patients who lived at home prior to acute hospitalization returned home.(116) **(EG 2)**

Rationale

Inpatient rehabilitation facilities are designed to provide intensive rehabilitation therapy in a resource intensive inpatient hospital environment. Use of this MCG care guideline helps in the decision to appropriately admit a patient to an acute inpatient rehabilitation facility and assess progress through an interdisciplinary plan of care that incorporates interventions for skilled nursing (eg, wound care, pain management, infusion medication) and rehabilitation therapy (eg, physical, occupational, speech-language) in alignment with CMS requirements.

Use of these evidence-based clinical criteria benefits patients by assisting in the identification of patients who, due to the complexity of their nursing, medical management, and rehabilitation needs, require and can reasonably be expected to benefit from an inpatient stay and an interdisciplinary team approach to the delivery of rehabilitation care. This evidence-based guidance ensures that Medicare and Medicare Advantage patients receive full access to Medicare benefits without risk of delay or decreased access. Additional evidence-based guidance benefits patients by outlining complicated factors (eg, pain exacerbations, infection, comorbid disease exacerbation) that could extend the need for a rehabilitation facility stay. Criteria for transitioning a patient to the next appropriate level of care (eg,

acute care, skilled nursing facility, home care) are also provided to assist providers and patients in identifying next steps for the patient's care. This support can increase consistency in treatment thresholds, leading to less variation in care and promoting equitable treatment among patients.

Related CMS Coverage Guidance

This guideline supplements but does not replace, modify, or supersede existing Medicare regulations or applicable National Coverage Determinations (NCDs) or Local Coverage Determinations (LCDs).

Code of Federal Regulations (CFR): 42 CFR 410.38(128); 42 CFR 412.29(2); 42 CFR 412.606(30); 42 CFR 412.610(31); 42 CFR Pt. 412.622(1)

Internet-Only Manual (IOM) Citations: CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 1 - Inpatient Hospital Services Covered Under Part A(3); CMS IOM Publication 100-08, Medicare Program Integrity Manual, Chapter 6 - Medical Review of Inpatient Hospital Claims for Part A(130)

Medicare Coverage Determinations: Medicare Coverage Database(131)

Other CMS Guidance: Centers for Medicare and Medicaid Services, "Inpatient Rehabilitation Facility Patient Assessment Instrument (IRF-PAI) and IRF-PAI Manual" Version 4.2(29)

References

- Centers for Medicare and Medicaid Services. "Basis of payment." 42 CFR Pt. 412.622 Washington, DC 2025 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9]
- Centers for Medicare and Medicaid Services. "Classification criteria for payment under the inpatient rehabilitation facility prospective payment system." 42 CFR Pt. 412.29 Washington, DC 2020 Aug 10 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/current/title-42/section-412.29>. [Context Link 1, 2, 3]
- Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09] [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13]
- Curry ZA, et al. Examination of pain comorbid diagnoses in the inpatient rehabilitation population across all impairment groups. American Journal of Physical Medicine and Rehabilitation 2024;Online. DOI: 10.1097/PHM.0000000000002512. [Context Link 1] View abstract...
- Scrutinio D, et al. Characteristics, Outcomes, and Long-Term Survival of Patients With Heart Failure Undergoing Inpatient Cardiac Rehabilitation. Archives of Physical Medicine and Rehabilitation 2022;103(5):891-898.e4. DOI: 10.1016/j.apmr.2021.10.014. [Context Link 1] View abstract...
- Centers for Medicare and Medicaid Services. "Criteria for skilled services and the need for skilled service." 42 CFR Pt. 409.32 Washington, DC 2023 Oct 02 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
- Centers for Medicare and Medicaid Services. "Skilled services requirements." 42 CFR Pt. 409.44 Washington, DC 2023 Oct 02 [accessed 2025 Jul 23] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
- Pryor J, O'Reilly K, Bonser M, Garret G, McKenchnie D. Rehabilitation for the individual and family. In: Chang E, Johnson A, editors. Living With Chronic Illness and Disability. 4th ed. Chatswood NSW 2067: Elsevier; 2022:161-182. [Context Link 1, 2]
- Simning A, Caprio TV, Seplaki CL, Temkin-Greener H, Szanton SL, Conwell Y. Patient-reported outcomes in functioning following nursing home or inpatient rehabilitation. Journal of the American Medical Directors Association 2018;19(10):864-870. DOI: 10.1016/j.jamda.2018.06.014. [Context Link 1] View abstract...
- Tan Dy RC, Nguyen N. Therapeutic exercise. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:145-154. [Context Link 1, 2]
- Medical-surgical nursing. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:33-55. [Context Link 1, 2, 3]

12. Therapy Outcomes in Post-Acute Care Settings (TOPS) Study Chartbook. [Internet] American Occupational Therapy Association, Inc and American Physical Therapy Association. 2021 Apr Accessed at: <https://www.aota.org/>. [accessed 2025 Sep 05] [Context Link 1]
13. Roberts PS, et al. Retrospective study demonstrating therapy time impact on inpatient rehabilitation functional gains. *Disability and Rehabilitation* 2022;44(17):4639-4647. DOI: 10.1080/09638288.2021.1912836. [Context Link 1] View abstract...
14. Dolansky M, Mola A. CR in inpatient and transitional settings. In: American Association of Cardiovascular and Pulmonary Rehabilitation, editor. *Guidelines for Cardiac Rehabilitation Programs*. 6th ed. Human Kinetics; 2021:15-32. [Context Link 1, 2, 3, 4, 5, 6, 7, 8]
15. Parcetch KM, Schmitz TJ. Examination and modification of the environment. In: Fulk G, Chui K, editors. *O'Sullivan & Schmitz's Physical Rehabilitation*. 8th ed. USA: F.A. Davis Company; 2024:292-331. [Context Link 1]
16. Kilma DW, Schmitz TJ, O'Sullivan SB. Examination of coordination and balance. In: Fulk G, Chui K, editors. *O'Sullivan & Schmitz's Physical Rehabilitation*. 8th ed. USA: F.A. Davis Company; 2024:180-214. [Context Link 1]
17. Myers J. Physical activity and exercise. In: American Association of Cardiovascular and Pulmonary Rehabilitation, editor. *Guidelines for Cardiac Rehabilitation Programs*. 6th ed. Human Kinetics; 2021:61-68. [Context Link 1]
18. Gait and Posture Analysis. In: Dutton M, editor. *Dutton's Orthopaedic Examination, Evaluation, and Intervention*. 6th ed. McGraw-Hill Education; 2023:272-318. [Context Link 1]
19. Goughwin PM, Cameron ID. Psychological assessment and intervention in rehabilitation. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:29-36. [Context Link 1, 2, 3, 4]
20. Pain management. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. *Clinical Nursing Skills and Techniques*. 11th ed. Elsevier; 2025:443-497. [Context Link 1, 2]
21. Dias KJ. Heart disease. In: Fulk G, Chui K, editors. *O'Sullivan & Schmitz's Physical Rehabilitation*. 8th ed. USA: F.A. Davis Company; 2024:415-468. [Context Link 1, 2, 3]
22. Makam AN, Grabowski DC. Policy in clinical practice: choosing post-acute care in the new decade. *Journal of Hospital Medicine* 2021;16(3):171-174. DOI: 10.12788/jhm.3577. [Context Link 1] View abstract...
23. Kilgore C. Nursing patients through rehabilitation. In: O'Hanlon S, Smith M, editors. *Comprehensive Guide to Rehabilitation of the Older Patient*. 4th ed. Elsevier; 2021:69-72. [Context Link 1]
24. Varma AK, Mishra S. Neurogenic lower urinary tract dysfunction. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:191-200. [Context Link 1]
25. Oey NE, Lim PA. Neurogenic bowel: Dysfunction and rehabilitation. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:201-210. [Context Link 1]
26. Eickmeyer SM, Barker KD, Sayyad A, Rydberg L. The rehabilitation of patients with advanced heart failure after left ventricular assist device placement: a narrative review. *PM & R: The Journal of Injury, Function, and Rehabilitation* 2019;11(1):64-75. DOI: 10.1016/j.pmrj.2018.05.027. [Context Link 1, 2] View abstract...
27. Management of patients with musculoskeletal disorders. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:1113-1151. [Context Link 1]
28. Brown TM, et al. Core Components of Cardiac Rehabilitation Programs: 2024 Update: A Scientific Statement From the American Heart Association and the American Association of Cardiovascular and Pulmonary Rehabilitation. *Circulation* 2024;Online. DOI: 10.1161/CIR.0000000000001289. (Reaffirmed 2025 Jul) [Context Link 1] View abstract...
29. Centers for Medicare and Medicaid Services. "Inpatient Rehabilitation Facility Patient Assessment Instrument (IRF-PAI) and IRF-PAI Manual." Version 4.2 Washington, DC 2024 Feb 01 [accessed 2025 Jul 23] Accessed at: <https://www.cms.gov/medicare/quality/inpatient-rehabilitation-facility/irf-pai-and-irf-grp-manual>. [Context Link 1, 2]
30. Centers for Medicare and Medicaid Services. "Patient assessments." 42 CFR Pt. 412.606 Washington, DC 2023 Aug [accessed 2025 Jul 23] Accessed at: <https://www.ecfr.gov/>. [Context Link 1, 2]
31. Centers for Medicare and Medicaid Services. "Assessment schedule." 42 CFR Pt. 412.610 Washington, DC 2025 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>. [Context Link 1, 2]
32. Imran A, Rawal MD, Botre N, Patil A. Improving and promoting social determinants of Health at a system level. *Joint Commission Journal on Quality and Patient Safety* 2022;48(8):376-384. DOI: 10.1016/j.jcjq.2022.06.004. [Context Link 1, 2] View abstract...

33. Shea CA, et al. Variation in 30-day readmission rates from inpatient rehabilitation facilities to acute care hospitals. *Journal of the American Medical Directors Association* 2021;22(12):2461-2467. DOI: 10.1016/j.jamda.2021.03.033. [Context Link 1] View abstract...
34. Wan CS, Reijnierse EM, Maier AB. Risk factors of readmissions in geriatric rehabilitation patients: RESORT. *Archives of Physical Medicine and Rehabilitation* 2021;102(8):1524-1532. DOI: 10.1016/j.apmr.2021.01.082. [Context Link 1] View abstract...
35. HEDIS® quality measures 2026. [Internet] NCQA. Accessed at: <https://www.ncqa.org/hedis/measures/>. Updated 2026 [accessed 2026 Apr 02] [Context Link 1, 2]
36. Wong R. Care planning meetings. In: O'Hanlon S, Smith M, editors. *Comprehensive Guide to Rehabilitation of the Older Patient*. 4th ed. Elsevier; 2021:395-398. [Context Link 1]
37. Galeema D. Behavior modification for risk factor reduction. In: American Association of Cardiovascular and Pulmonary Rehabilitation, editor. *Guidelines for Cardiac Rehabilitation Programs*. 6th ed. Human Kinetics; 2021:87-96. [Context Link 1]
38. Falls: assessment and prevention in older people and in people 50 and over at higher risk. NICE Guideline NG249 [Internet] National Institute for Health and Care Excellence. 2025 Apr Accessed at: <https://www.nice.org.uk/>. [accessed 2025 Aug 18] [Context Link 1, 2]
39. Fluid and electrolytes. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:224-272. [Context Link 1, 2, 3]
40. Intravenous and vascular access therapy. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. *Clinical Nursing Skills and Techniques*. 11th ed. Elsevier; 2025:842-894. [Context Link 1, 2, 3]
41. Medicare Coverage Database Local Coverage Determination documents. [Internet] Centers for Medicare and Medicaid Services. Accessed at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Updated 2024 [accessed 2025 Jan 08] [Context Link 1, 2]
42. Aberegg ES. Nutritional guidelines. In: American Association of Cardiovascular and Pulmonary Rehabilitation, editor. *Guidelines for Cardiac Rehabilitation Programs*. 6th ed. Human Kinetics; 2021:69-86. [Context Link 1]
43. European Pressure Ulcer Advisory Panel, National Pressure Injury Advisory Panel, and Pan Pacific Pressure Injury Alliance. Prevention and Treatment of Pressure Ulcers/Injuries: Clinical Practice Guideline. The International Guideline Online Living Guideline 4th Edition [Internet] National Pressure Injury Advisory Panel. 2025 Feb Accessed at: <https://npiap.com/page/InternationalGuidelines>. [accessed 2025 Sep 16] [Context Link 1]
44. de Wilde C. Social work. In: O'Hanlon S, Smith M, editors. *Comprehensive Guide to Rehabilitation of the Older Patient*. 4th ed. Elsevier; 2021:91-94. [Context Link 1]
45. Kee YYK. Measuring progress with rehabilitation. In: O'Hanlon S, Smith M, editors. *Comprehensive Guide to Rehabilitation of the Older Patient*. 4th ed. Elsevier; 2021:111-115. [Context Link 1]
46. Admitting, transfer, and discharge. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. *Clinical Nursing Skills and Techniques*. 11th ed. Elsevier; 2025:40-51. [Context Link 1, 2, 3]
47. Shiner CT, Woodbridge G, Skalicky DA, Faux SG. Multidisciplinary inpatient rehabilitation following heart and/or lung transplantation-examining cohort characteristics and clinical outcomes. *PM & R: The Journal of Injury, Function, and Rehabilitation* 2019;11(8):849-857. DOI: 10.1002/pmrj.12057. [Context Link 1] View abstract...
48. Assessment and management of patients with vascular disorders and problems of peripheral circulation. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:817-864. [Context Link 1]
49. Stevens SM, et al. Antithrombotic therapy for VTE disease: second update of the CHEST guideline and expert panel report. *Chest* 2021;160(6):e545-e608. DOI: 10.1016/j.chest.2021.07.055. (Reaffirmed 2025 Jan) [Context Link 1] View abstract...
50. Bourke R, Rivasi G, Kenny RA. Falls, Dizziness and Funny Turns. In: O'Hanlon S, Smith M, editors. *Comprehensive Guide to Rehabilitation of the Older Patient*. 4th ed. Elsevier; 2021. [Context Link 1]
51. Oh-Park M, Gruber L. Geriatrics. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:289-296. [Context Link 1]
52. Blum FC, Biros MH. Fever in the adult patient. In: Walls RM, editor. *Rosen's Emergency Medicine: Concepts and Clinical Practice*. 10th ed. Elsevier; 2023:90-95.e1. [Context Link 1]
53. Assessment of respiratory function. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:462-493. [Context Link 1, 2]

54. Management of patients with oncologic disorders. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:301-366. [Context Link 1]
55. Complications of parenteral nutrition. In: Ayers P, Bobo ES, Hurt RT, Mays AA, Worthington PH, editors. A.S.P.E.N. Parenteral Nutrition Handbook. 3rd ed. Silver Spring, MD: American Society for Parenteral and Enteral Nutrition; 2020:173-198. [Context Link 1]
56. Parenteral nutrition administration and monitoring. In: Ayers P, Bobo ES, Hurt RT, Mays AA, Worthington PH, editors. A.S.P.E.N. Parenteral Nutrition Handbook. 3rd ed. Silver Spring, MD: American Society for Parenteral and Enteral Nutrition; 2020:141-172. [Context Link 1]
57. Rateau MR. Fluid, electrolyte, and acid-base imbalances. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:302-334. [Context Link 1, 2]
58. Safety. In: Lynn P, editor. Taylor's Clinical Nursing Skills. 6th ed. Wolters Kluwer; 2023:153-195. [Context Link 1]
59. Fosbre CD. Anger, aggression, and violence. In: Fosbre CD, Varcarolis EM, Chiappetta L, editors. Varcarolis' Essentials of Psychiatric-Mental Health Nursing. 5th ed. Elsevier; 2023:388-401. [Context Link 1]
60. Pain management. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:196-223. [Context Link 1, 2, 3]
61. Opioid analgesics, opioid antagonists, and nonopioid centrally acting analgesics. In: Burchum JR, Rosenthal LD, editors. Lehne's Pharmacology for Nursing Care. 12th ed. Elsevier; 2025:306-326. [Context Link 1]
62. Ezidiegwu D, Kim SY. Pain management principles. In: Gonzalez-Fernandez M, Schaaf S, Zumsteg JM, Perret D, Kennedy DJ, editors. Handbook of Physical Medicine and Rehabilitation. Springer Publishing Company; 2022:375-384. [Context Link 1]
63. Dowell D, Ragan KR, Jones CM, Baldwin GT, Chou R. CDC Clinical Practice Guideline for Prescribing Opioids for Pain - United States, 2022. MMWR - Recommendations and Reports 2022;71(3):1-95. DOI: 10.15585/mmwr.r7103a1. [Context Link 1] View abstract...
64. Vital signs. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. Clinical Nursing Skills and Techniques. 11th ed. Elsevier; 2025:70-109. [Context Link 1, 2, 3]
65. Management of patients with chest and lower respiratory tract disorders. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:526-601. [Context Link 1]
66. Management of patients with neurologic dysfunction. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:1992-2030. [Context Link 1]
67. Amerson C. Chronic neurologic problems. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:1538-1572. [Context Link 1]
68. Ainger E, Ninan S. Exercise tolerance. In: O'Hanlon S, Smith M, editors. Comprehensive Guide to Rehabilitation of the Older Patient. 4th ed. Elsevier; 2021:171-176. [Context Link 1]
69. Management of patients with upper respiratory tract disorders. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:494-525. [Context Link 1]
70. Sarasnick JA. Upper respiratory problems. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:577-595. [Context Link 1]
71. Airway management. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. Clinical Nursing Skills and Techniques. 11th ed. Elsevier; 2025:753-788. [Context Link 1]
72. Management of patients with urinary disorders. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:1604-1637. [Context Link 1]
73. Doiron RC, Nickel JC. Interstitial cystitis/bladder pain syndrome. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:325-331. [Context Link 1]
74. Mount DB. Azotemia and urinary abnormalities. In: Loscalzo J, Fauci A, Kasper D, Hauser S, Longo D, Jameson JL, editors. Harrison's Principles of Internal Medicine. 21st ed. McGraw Hill Education; 2022:331-338. [Context Link 1]

75. Boswell B, Thomas AA. Pediatric genitourinary and renal tract disorders. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:2169-2183.e2. [Context Link 1]
76. Waxman MA, Moran GJ. Pneumonia. In: Walls RM, editor. Rosen's Emergency Medicine: Concepts and Clinical Practice. 10th ed. Elsevier; 2023:829-838.e2. [Context Link 1]
77. Mondor EE. Supporting ventilation. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:539-576. [Context Link 1]
78. Ratliff CR. Inflammation and healing. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:179-198. [Context Link 1]
79. Nelson F. Physical medicine and rehabilitation: physical therapy and occupational therapy. In: Nelson F, editor. A Manual of Orthopaedic Terminology. 9th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:386-396. [Context Link 1]
80. Exercise testing and prescription for populations with other chronic diseases and health conditions. In: American College of Sports Medicine, editor. ACSM's Guidelines for Exercise Testing and Prescription. 12th ed. Wolters Kluwer; 2025:389-480. [Context Link 1, 2]
81. Hussein N. Acute medical conditions. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:253-264. [Context Link 1, 2, 3]
82. Snyder JS, Sump CA. Cardiac surgery. In: Snyder JS, Sump CA, editors. Swearingen's All-in-One Nursing Care Planning Resource. 6th ed. Elsevier; 2024:156-158. [Context Link 1, 2, 3, 4]
83. Cardiac and vascular care plans. In: Gulanick M, Myers JL, Bowman-Woodall C, editors. Nursing Care Plans. 11th ed. Elsevier; 2025:308-394. [Context Link 1]
84. Stoeckmann TM, O'Sullivan SB. Strategies to improve motor function. In: Fulk G, Chui K, editors. O'Sullivan & Schmitz's Physical Rehabilitation. 8th ed. USA: F.A. Davis Company; 2024:333-367. [Context Link 1, 2]
85. Ritchey KC, Olney AL, Campen SJ. Falls. In: Gonzalez-Fernandez M, Schaaf S, Zumsteg JM, Perret D, Kennedy DJ, editors. Handbook of Physical Medicine and Rehabilitation. Springer Publishing Company; 2022:337-342. [Context Link 1, 2]
86. Michael E, Farrell E, Linden M, Cabahug PG, Sadowsky CL, Recio AC. Wheelchairs. In: Gonzalez-Fernandez M, Schaaf S, Zumsteg JM, Perret D, Kennedy DJ, editors. Handbook of Physical Medicine and Rehabilitation. Springer Publishing Company; 2022:543-548. [Context Link 1]
87. Russek LN. Chronic pain. In: Fulk G, Chui K, editors. O'Sullivan & Schmitz's Physical Rehabilitation. 8th ed. USA: F.A. Davis Company; 2024:966-1010. [Context Link 1]
88. Parish LC, Karadag AS, Parish JL. Decubitus (pressure) ulcers. In: Lebowitz MG, Heymann WR, Coulson IH, Murrell DF, editors. Treatment of Skin Disease. 6th ed. Elsevier; 2022:185-192. [Context Link 1]
89. Patient safety. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. Clinical Nursing Skills and Techniques. 11th ed. Elsevier; 2025:382-416. [Context Link 1]
90. Bezner JR, Maxwell J. Promoting health and wellness. In: Fulk G, Chui K, editors. O'Sullivan & Schmitz's Physical Rehabilitation. 8th ed. USA: F.A. Davis Company; 2024:1118-1148. [Context Link 1]
91. Effgen SK. The educational environment. In: Palisano RJ, Orlin M, Schreiber J, editors. Campbell's Physical Therapy for Children. 6th ed. Elsevier; 2023:768-795. [Context Link 1]
92. Tissue behavior, injury, healing, and treatment. In: Dutton M, editor. Dutton's Orthopaedic Examination, Evaluation, and Intervention. 6th ed. McGraw-Hill Education; 2023:26-56. [Context Link 1, 2]
93. Wu CH. Physical agent modalities. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:165-172. [Context Link 1]
94. Hsieh LF. Rheumatologic rehabilitation. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:297-310. [Context Link 1]
95. Liu DH. Auditory, vestibular, and visual impairments. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:519-526. [Context Link 1]
96. Fritz NE, McIsaac TL. Multiple sclerosis. In: Fulk G, Chui K, editors. O'Sullivan & Schmitz's Physical Rehabilitation. 8th ed. USA: F.A. Davis Company; 2024:588-634. [Context Link 1]
97. McIsaac TL, Bezkor EW. Parkinson disease. In: Fulk G, Chui K, editors. O'Sullivan & Schmitz's Physical Rehabilitation. 8th ed. USA: F.A. Davis Company; 2024:679-725. [Context Link 1]

98. Fulk GD, Nirider C, Williams G, DeBlois A. Traumatic brain injury. In: Fulk G, Chui K, editors. O'Sullivan & Schmitz's Physical Rehabilitation. 8th ed. USA: F.A. Davis Company; 2024:726-759. [Context Link 1]
99. Coppola S, Gillen, Boyt Schell BA. Contemporary occupational therapy practice and future directions. In: Gillen G, Brown C, editors. Willard & Spackman's Occupational Therapy. 14th ed. Wolters Kluwer Health; 2023:54-70. [Context Link 1]
100. Nilsen DM, Gillen G. Motor function and occupational performance. In: Gillen G, Brown C, editors. Willard & Spackman's Occupational Therapy. 14th ed. Wolters Kluwer Health; 2023:911-941. [Context Link 1]
101. Toglia JP, Goverover Y. Cognitive, perception, and occupational performance. In: Gillen G, Brown C, editors. Willard & Spackman's Occupational Therapy. 14th ed. Wolters Kluwer Health; 2023:942-991. [Context Link 1]
102. Hobbs M. Cardiopulmonary disorders. In: Atchison BJ, Dirette DP, editors. Conditions in Occupational Therapy: Effect on Occupational Performance. 6th ed. Wolters Kluwer; 2023:388-406. [Context Link 1, 2, 3, 4]
103. Takatani S. Cardiac and pulmonary disease. In: Pendleton HM, Schultz-Krohn W, editors. Pedretti's Occupational Therapy. 9th ed. Elsevier; 2024:1185-1204. [Context Link 1, 2, 3, 4, 5, 6]
104. Miller A. General conditioning. In: Atchison BJ, Dirette DP, editors. Conditions in Occupational Therapy: Effect on Occupational Performance. 6th ed. Wolters Kluwer; 2023:726-745. [Context Link 1, 2, 3, 4]
105. Pitonyak JS, Wilbur KL. Activities of daily living and instrumental activities of daily living. In: Gillen G, Brown C, editors. Willard & Spackman's Occupational Therapy. 14th ed. Wolters Kluwer Health; 2023:706-743. [Context Link 1]
106. Dobkin BH. Neurological rehabilitation. In: Jankovic J, Mazziotta JC, Pomeroy SL, Newman NJ, editors. Bradley and Daroff's Neurology in Clinical Practice. 8th ed. Elsevier; 2022:819-849.e2. [Context Link 1]
107. Eapen BC, Oshiro RL. Vascular diseases. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:235-242. [Context Link 1]
108. Dicks MA, Verduzco-Gutierrez M. Spasticity and contracture. In: Gonzalez-Fernandez M, Schaaf S, Zumsteg JM, Perret D, Kennedy DJ, editors. Handbook of Physical Medicine and Rehabilitation. Springer Publishing Company; 2022:427-434. [Context Link 1]
109. Manual techniques. In: Dutton M, editor. Dutton's Orthopaedic Examination, Evaluation, and Intervention. 6th ed. McGraw-Hill Education; 2023:382-404. [Context Link 1]
110. Ramiro RPT. Occupational medicine and vocational rehabilitation. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:43-54. [Context Link 1, 2, 3]
111. Schultz-Krohn W, Pendleton HM. Application of the occupational therapy practice framework to physical dysfunction. In: Pendleton HM, Schultz-Krohn W, editors. Pedretti's Occupational Therapy. 9th ed. Elsevier; 2024:29a-53. [Context Link 1]
112. Daras LC, et al. Evaluating hospital readmission rates after discharge from inpatient rehabilitation. Archives of Physical Medicine and Rehabilitation 2018;99(6):1049-1059. DOI: 10.1016/j.apmr.2017.07.008. [Context Link 1] View abstract...
113. Middleton A, Graham JE, Ottenbacher KJ. Functional status is associated with 30-day potentially preventable hospital readmissions after inpatient rehabilitation among aged Medicare Fee-for-Service beneficiaries. Archives of Physical Medicine and Rehabilitation 2018;99(6):1067-1076. DOI: 10.1016/j.apmr.2017.05.001. [Context Link 1] View abstract...
114. Osuafor CN, Sahimi SNM, Enduluri S, McCarthy F. Incidence and outcome of interrupted geriatric rehabilitation requiring acute hospital transfer. Irish Journal of Medical Science 2019;188(4):1451-1454. DOI: 10.1007/s11845-019-01974-0. [Context Link 1] View abstract...
115. Biddlespacher K, Mulkey DC. The Effect of Teach-Back on Readmission Rates in Rehabilitation Patients. Rehabilitation Nursing 2024 Mar-Apr 01;49(2):65-72. DOI: 10.1097/RNJ.0000000000000452. [Context Link 1] View abstract...
116. Philippe AM, Farid R. From inpatient rehabilitation facility to skilled nursing facility to home: A retrospective study of functional recovery. PM & R: The Journal of Injury, Function, and Rehabilitation 2024;16(3):226-230. DOI: 10.1002/pmrj.13065. [Context Link 1] View abstract...
117. Program standards. In: CARF, International, editors. Medical Rehabilitation Standards Manual. 2025 ed. Tucson, AZ: Commission on Accreditation of Rehabilitation Facilities; 2025:189-272. [Context Link 1]

118. Adult health and physical, nutritional, and cultural assessment. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:71-96. [Context Link 1]
119. The nursing process in drug therapy and patient safety. In: Tucker RG, editor. Karch's Focus on Nursing Pharmacology. 9th ed. Wolters Kluwer; 2023:47-57. [Context Link 1]
120. Adverse drug reactions and medication errors. In: Burchum JR, Rosenthal LD, editors. Lehne's Pharmacology for Nursing Care. 12th ed. Elsevier; 2025:64-75. [Context Link 1]
121. Health education and health promotion. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:56-70. [Context Link 1, 2]
122. Safe medication preparation. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. Clinical Nursing Skills and Techniques. 11th ed. Elsevier; 2025:586-607. [Context Link 1]
123. Angelelli J, et al. Effect of social determinants of health interventions on adults living with disabilities: a scoping review. Archives of Physical Medicine and Rehabilitation 2022;103(5):1023-1033.e11. DOI: 10.1016/j.apmr.2021.06.021. [Context Link 1] View abstract...
124. Ezell JM, Hamdi S, Borrero N. Approaches to addressing nonmedical services and care coordination needs for older adults. Research on Aging 2022 Mar-Apr;44(3-4):323-333. DOI: 10.1177/01640275211033929. [Context Link 1] View abstract...
125. Martinez-Kratz M. Guide to planning care. In: Makic MB, Martinez-Kratz MR, editors. Ackley and Ladwig's Nursing Diagnosis Handbook. 13th ed. Elsevier; 2023:125-1051. [Context Link 1]
126. Bassil NK, Ohanian MLK, Bou Saba TG. Nicotine use disorder in older adults. Clinics in Geriatric Medicine 2022;38(1):119-131. DOI: 10.1016/j.cger.2021.07.008. [Context Link 1] View abstract...
127. Harding MM. Substance use disorders in acute care. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:162-178. [Context Link 1]
128. Centers for Medicare and Medicaid Services. "Durable medical equipment, prosthetics, orthotics and supplies (DMEPOS): scope and conditions." 42 CFR Pt. 410.38 Washington, DC 2024 Jan 01 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/current/title-42/>. [Context Link 1, 2]
129. Care and Compare. Find and Compare Nursing Homes, Hospitals, and Other Providers Near You [Internet] Medicare.gov. Accessed at: <https://www.medicare.gov/care-compare/>. [accessed 2025 Apr 02] [Context Link 1, 2]
130. Centers for Medicare and Medicaid Services. Medicare Program Integrity Manual. Chapter 6 - Medical Review of Inpatient Hospital Claims for Part A Payment Rev. 10365 [Internet] Centers for Medicare and Medicaid Services. 2020 Oct 02 Accessed at: <https://www.cms.gov/medicare/regulations-guidance/manuals/internet-only-manuals-ioms>. [accessed 2025 Sep 09] [Context Link 1]
131. Medicare Coverage Database. [Internet] Centers for Medicare and Medicaid Services. Accessed at: <https://www.cms.gov/medicare-coverage-database/search.aspx>? Updated 2025 [accessed 2025 Oct 23] [Context Link 1]

Footnotes

[A] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) requires specific considerations for admission to this level of care for certain diagnoses, which are outlined in the electronic Code of Federal Regulations part 412.(1)(2) [A in Context Link 1]

[B] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) requires a preadmission screening to be conducted within 48 hours preceding the inpatient rehabilitation admission and is to be completed by a licensed or certified clinician or physician either in person or by telephone through review of the referring hospital medical records.(1) [B in Context Link 1]

[C] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) required elements of the preadmission screening include: level of function prior to the event, condition that caused the need for rehabilitation, expected level of improvement and expected length of time to achieve that level of improvement, risk for clinical complications, combinations of treatments needed (PT, OT, SLP, or orthosis), and anticipated discharge destination.(1) [C in Context Link 1]

[D] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) defines reasonable and necessary skilled care as a service that is "so inherently complex that it can only be safely and effectively performed by, or under the supervision of, professional or technical personnel" based on the documented assessment of the patient's individual care needs. In some cases, a service that would originally be considered unskilled may be considered skilled if the patient's underlying conditions or complications require the involvement of licensed personnel to ensure that essential nonskilled care is achieving its purpose.(6)(7) [D in Context Link 1, 2]

[E] **National guidelines:** According to the Centers for Medicare and Medicaid Services (CMS), the requirement for medical supervision means that a licensed physician who is determined by the inpatient rehabilitation facility to have specialized training and experience in inpatient rehabilitation must conduct face-to-face encounters with the patient at least 3 days per week throughout the patient's stay to assess the patient, as well as to modify the course of treatment as needed to maximize the patient's ability to benefit from rehabilitation. After the first week of physician encounters, CMS will allow a non-physician practitioner to independently conduct one of the three required visits.(3) [E in Context Link 1]

[F] **National guidelines:** According to the Centers for Medicare and Medicaid Services (CMS), the interdisciplinary care team must meet weekly throughout the patient's stay and consist of a registered nurse with specialized training or experience in rehabilitation, a social worker or case manager (or both), a licensed or certified therapist from each therapy discipline involved with the patient's care, and be led by a rehabilitation physician.(3) [F in Context Link 1]

[G] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) uses a data collection system to establish best practice, understand costs, and identify cost-effective systems and procedures. The current data collection system for eligible patients is the Inpatient Rehabilitation Facility Patient Assessment Instrument (IRF-PAI). IRF-PAI is an assessment tool used to identify patient problems, strengths, and preferences, and to create an individualized care plan.(29) [G in Context Link 1]

[H] **National guidelines:** Starting October 1, 2024, the Centers for Medicare and Medicaid Services (CMS) requires inpatient rehabilitation facilities to use the Inpatient Rehabilitation Facility Patient Assessment Instrument (IRF-PAI) to assess all inpatients, regardless of payer. The admission IRF-PAI should cover calendar days 1 through 3 of the patient's current admission; the patient assessment schedule is further detailed in the electronic Code of Federal Regulations part 412.610.(30)(31) [H in Context Link 1]

[I] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) requires that an individualized overall plan of care for the patient is developed by a rehabilitation physician with input from the interdisciplinary team within 4 days of the patient's admission.(1) [I in Context Link 1]

[J] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) states that therapy treatments should begin within 36 hours of admission.(1)(3) [J in Context Link 1]

[K] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) recommends that if an unexpected clinical event occurs that limits a patient's ability to participate in an intensive therapy program for a brief period, not to exceed 3 days, the specific reasons for the break should be documented in the patient's medical record.(3) [K in Context Link 1, 2]

[L] Pain assessment should include how pain affects patient's physical and psychosocial function and progress toward treatment goals.(60) [L in Context Link 1]

[M] The patient's agreed-upon written treatment plan should include measurable and realistic functional or quality-of-life goals, pain-management education, treatment options, and safe use of opioid and nonopioid medications as prescribed.(60) [M in Context Link 1]

[N] **National guidelines:** The Centers for Disease Control and Prevention (CDC) recommends nonpharmacologic and nonopioid therapy before opioids for management of chronic pain (not due to malignancy). Considerations for opioids should include pain-management goal setting and determining if benefits outweigh risks. When opioids are prescribed, they should be provided in conjunction with nonpharmacologic pain-management interventions and planned monitoring for misuse or use escalation, including naloxone education.(63) [N in Context Link 1]

Definitions

Custodial care

- Custodial care is personal unskilled care to support the patient's care of activities of daily living (ADL), such as bathing, dressing, and eating.(1)

References

1. Mauk KL. Post-acute care. In: Larsen PD, editor. Lubkin's Chronic Illness: Impact and Intervention. 11th ed. Jones & Bartlett Learning; 2022:477-520.

FIM Score

- The Functional Independence Measure (FIM®) is an instrument that classifies a patient's ability to carry out activities and the need for assistance from a person or device. It is a 7-level scale measurement of gradations in behavior (from dependence to independence) that can be used to monitor patient progress throughout rehabilitation and help with care plan development and defining discharge needs.(1)

References

1. Carter WE III, Peretiakko S. Quality and outcome measures for medical rehabilitation. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:55-62.

Goals for measurable improvement

- Goals for measurable improvement include identification of **ALL** of the following(1)(2):
 - Nature and degree of improvement
 - Time expected for achievement
 - Functional improvement anticipated to be practical, ongoing, and sustainable

References

1. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09]
2. Pryor J, O'Reilly K, Bonser M, Garret G, McKenchnie D. Rehabilitation for the individual and family. In: Chang E, Johnson A, editors. Living With Chronic Illness and Disability. 4th ed. Chatswood NSW 2067: Elsevier; 2022:161-182.

Intensive rehabilitation therapy program

- Intensive rehabilitation therapy program that includes **1 or more** of the following[A](1)(2):
 - Therapy at least 3 hours per day for 5 days per week
 - Therapy at least 15 hours per week (in a 7 consecutive calendar day period starting from the date of admission)
 - Other intensive therapy to improve functional capacity at patient's maximum participation level without compromising safety or exceeding ability or tolerance(3)

References

1. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09]

2. Centers for Medicare and Medicaid Services. "Basis of payment." 42 CFR Pt. 412.622 Washington, DC 2025 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>.
3. Pryor J, O'Reilly K, Bonser M, Garret G, McKenchnie D. Rehabilitation for the individual and family. In: Chang E, Johnson A, editors. Living With Chronic Illness and Disability. 4th ed. Chatswood NSW 2067: Elsevier; 2022:161-182.

Footnotes

- A. **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) Medicare Benefit Policy Manual states that admissions that are intended to assess if a patient would benefit and tolerate treatment in an inpatient rehabilitation facility (IRF) are no longer considered reasonable and necessary.(1)

Interdisciplinary team care

- Interdisciplinary care team meetings held weekly and discipline examples include(1)(2)(3)(4):
 - A rehabilitation physician designated by the Inpatient Rehabilitation Facility to have specialized training and experience in inpatient rehabilitation services
 - A registered nurse with specialized training or experience in rehabilitation
 - A social worker or case manager (or both)
 - A licensed or certified therapist from each discipline involved in evaluating and treating the patient

References

1. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09]
2. Centers for Medicare and Medicaid Services. "Basis of payment." 42 CFR Pt. 412.622 Washington, DC 2025 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>.
3. Pryor J, O'Reilly K, Bonser M, Garret G, McKenchnie D. Rehabilitation for the individual and family. In: Chang E, Johnson A, editors. Living With Chronic Illness and Disability. 4th ed. Chatswood NSW 2067: Elsevier; 2022:161-182.
4. Centers for Medicare and Medicaid Services. Medicare Program Integrity Manual. Chapter 6 - Medical Review of Inpatient Hospital Claims for Part A Payment Rev. 10365 [Internet] Centers for Medicare and Medicaid Services. 2020 Oct 02 Accessed at: <https://www.cms.gov/medicare/regulations-guidance/manuals/internet-only-manuals-ioms>. [accessed 2025 Sep 09]

Occupational therapy (OT) progression

- OT progression includes **ALL** of the following(1)(2):
 - Ongoing evaluation of treatment plan with measurable short-term and long-term goals
 - Documentation of effectiveness and efficiency toward achieving established goals
 - Patient demonstrates resolving of barriers to transition of care and **1 or more** of the following:
 - Objective measurements document progress toward goals in reasonable and predictable time frame based on treatment plan.[A](4)
 - Minimal progress due to unexpected clinical event, but progress expected to resume toward goals within 3 days as condition improves(5)

References

1. Bennett K. Setting rehabilitation goals. In: O'Hanlon S, Smith M, editors. Comprehensive Guide to Rehabilitation of the Older Patient. 4th ed. Elsevier; 2021:49-52.

2. Pryor J, O'Reilly K, Bonser M, Garret G, McKenchnie D. Rehabilitation for the individual and family. In: Chang E, Johnson A, editors. Living With Chronic Illness and Disability. 4th ed. Chatswood NSW 2067: Elsevier; 2022:161-182.
3. Carter WE III, Peretiatko S. Quality and outcome measures for medical rehabilitation. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:55-62.
4. Kee YYK. Measuring progress with rehabilitation. In: O'Hanlon S, Smith M, editors. Comprehensive Guide to Rehabilitation of the Older Patient. 4th ed. Elsevier; 2021:111-115.
5. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09]

Footnotes

- A. Improvements may be measured by accuracy (percentage or number of correct trials), increased number of repetitions, decreased assistance level (maximum, moderate, or minimal assistance), decreased pain level, increased ROM/strength, increased balance scores, decreased level of cues or prompts required (in conjunction with level of assistance), or improvement in standardized functional outcome measures (eg, Continuity Assessment Record and Evaluation (CARE), WeeFIM®, Tinetti, Berg Balance Scale).(3)(4) Generally, it is expected that measured improvements should be demonstrable in targeted areas over a 1-week to 2-week period of skilled therapy. The speed of recovery is variable and may be linked to intensity of treatment, patient condition, age, comorbidities, and other factors.

Physical therapy (PT) progression

- PT progression includes **ALL** of the following(1)(2)(3)(4):
 - Ongoing evaluation of treatment plan with measurable short-term and long-term goals
 - Documentation of effectiveness and efficiency toward achieving established goals
 - Patient demonstrates resolving of barriers to transition of care and **1 or more** of the following:
 - Objective measurements document progress toward goals in reasonable and predictable time frame based on treatment plan.[A]
 - Minimal progress due to unexpected clinical event, but progress expected to resume toward goals within 3 days as condition improves

References

1. Centers for Medicare and Medicaid Services. "Skilled services requirements." 42 CFR Pt. 409.44 Washington, DC 2023 Oct 02 [accessed 2025 Jul 23] Accessed at: <https://www.ecfr.gov/>.
2. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09]
3. Hussein N. Acute medical conditions. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:253-264.
4. Carter WE III, Peretiatko S. Quality and outcome measures for medical rehabilitation. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:55-62.
5. Scalzitti DA. Examination of function. In: Fulk G, Chui K, editors. O'Sullivan & Schmitz's Physical Rehabilitation. 8th ed. USA: F.A. Davis Company; 2024:270-291.

Footnotes

- A. Improvements may be measured by accuracy (percentage or number of correct trials), increased number of repetitions, decreased assistance level (maximum, moderate, or minimal assistance), decreased pain level, increased ROM/strength, increased balance scores, decreased level of cues or prompts required (in

conjunction with level of assistance), or improvement in standardized functional outcome measures (eg, Continuity Assessment Record and Evaluation (CARE), WeeFIM®, Tinetti, Berg Balance Scale). Generally, it is expected that measured improvements should be demonstrable in targeted areas over a 1-week to 2-week period of skilled therapy. The speed of recovery is variable and may be linked to intensity of treatment, patient condition, age, comorbidities, and other factors.(4)(5)

Safe to go home

- Patient safe to go home if **ALL** of the following exist(1)(2)(3):
 - Medical condition manageable at home
 - Functional status manageable at home
 - Mental status stable for patient's condition
 - Medication availability confirmed and reconciliation complete
 - Patient/caregiver education complete and written discharge instructions provided
 - Caregiver and community resources identified (as needed)
 - Community resources identified and referrals made, as needed
 - Home care arranged, if indicated
 - Necessary medical equipment delivery arranged or available in home, if indicated
 - Necessary medical supplies ordered, or patient/caregiver can obtain, if indicated

References

1. Elements of Excellence in Transitions of Care (TOC). TOC Checklist [Internet] National Transitions of Care Coalition. 2006 Accessed at: <https://www.ntocc.org>. [accessed 2025 Sep 16]
2. Medical-surgical nursing. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:33-55.
3. Admitting, transfer, and discharge. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. Clinical Nursing Skills and Techniques. 11th ed. Elsevier; 2025:40-51.

Skilled care

- Skilled care may be direct (eg, in-person assessment and administration of treatments) or indirect (eg, coordinating care between providers, supervising care aides); examples include(1)(2)(3)(4):
 - Performed or supervised by professional or technical personnel
 - Exceed scope of Custodial care
 - Service is considered skilled as indicated by either of the following:
 - Service that is "so inherently complex that it can only be safely and effectively performed by, or under the supervision of, professional or technical personnel" based on the documented assessment of the patient's individual care needs.
 - Service that would originally be considered unskilled may be considered skilled if the patient's underlying conditions or complications require the involvement of licensed personnel to ensure that essential nonskilled care is achieving its purpose.

References

1. Centers for Medicare and Medicaid Services. "Criteria for skilled services and the need for skilled service." 42 CFR Pt. 409.32 Washington, DC 2023 Oct 02 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>.

2. Harding MM. Professional nursing. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:1-18.
3. Medical-surgical nursing. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:33-55.
4. Centers for Medicare and Medicaid Services. Medicare Program Integrity Manual. Chapter 6 - Medical Review of Inpatient Hospital Claims for Part A Payment Rev. 10365 [Internet] Centers for Medicare and Medicaid Services. 2020 Oct 02 Accessed at: <https://www.cms.gov/medicare/regulations-guidance/manuals/internet-only-manuals-ioms>. [accessed 2025 Sep 09]

Transition of care planning completed

- Transition of care planning completed; including **ALL** of the following(1)(2):
 - Transition plan communicated to all members of patient's healthcare team
 - Summary of inpatient rehabilitation facility admission, current list of medications, and discharge plan communicated to primary care provider at next level of care
 - Medication reconciliation complete
 - Education on condition management and complications to report provided to patient or caregiver
 - Follow-up appointments scheduled
 - Referrals to continue rehabilitation goals (eg, outpatient or home therapy) arranged, if needed
 - Services (eg, equipment, environment modifications, transportation) required for transition to next level of care arranged, if needed
 - Psychosocial issues addressed, or plan for management at next level of care arranged, if needed

References

1. Transition of care. In: Gillingham DC, editor. Foundations of Case Management. Blue Bayou Press; 2021:137-144.
2. Admitting, transfer, and discharge. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. Clinical Nursing Skills and Techniques. 11th ed. Elsevier; 2025:40-51.

Codes

ICD-10 Diagnosis: I09.81, I11.0, I13.0, I13.2, I20.0, I20.1, I20.2, I20.81, I20.89, I20.9, I21.01, I21.02, I21.09, I21.11, I21.19, I21.21, I21.29, I21.3, I21.4, I21.9, I21.A1, I21.A9, I21.B, I22.0, I22.1, I22.2, I22.8, I22.9, I23.3, I23.7, I24.0, I24.89, I24.9, I25.110, I25.111, I25.112, I25.118, I25.119, I25.700, I25.701, I25.702, I25.708, I25.709, I25.710, I25.711, I25.712, I25.718, I25.719, I25.720, I25.721, I25.722, I25.728, I25.729, I25.730, I25.731, I25.732, I25.738, I25.739, I25.750, I25.751, I25.752, I25.758, I25.759, I25.760, I25.761, I25.762, I25.768, I25.769, I25.790, I25.791, I25.792, I25.798, I25.799, I45.6, I46.2, I46.8, I46.9, I47.0, I47.10, I47.19, I47.20, I47.21, I47.29, I47.9, I48.0, I48.11, I48.19, I48.20, I48.21, I48.3, I48.4, I48.91, I48.92, I49.01, I49.02, I49.1, I49.2, I49.3, I49.40, I49.49, I49.5, I49.8, I49.9, I50.1, I50.20, I50.21, I50.22, I50.23, I50.30, I50.31, I50.32, I50.33, I50.40, I50.41, I50.42, I50.43, I50.814, I50.82, I50.83, I50.84, I50.89, I50.9, T82.855A, T82.855D, T82.855S, T86.22 [Hide]

ICD-10 Procedure: 5A2204Z

CPT®: 0659T, 92941, 92960

CPT copyright 2025 American Medical Association. All rights reserved.

Copyright © 2026 MCG Health, LLC
All Rights Reserved

Last Update: 1/25/2026 12:58:13 AM
Build Number: 30.0.2026012500524.025256