

Recovery Facility Care > Optimal Recovery Guidelines > Inpatient Rehabilitation Facility (IRF) > Problem Oriented Guidelines > Inpatient Rehabilitation Facility (Acute Rehabilitation): Musculoskeletal (I-7004)

Inpatient Rehabilitation Facility (Acute Rehabilitation): Musculoskeletal

IRF: I-7004 (RFC)

[Link to Codes](#)

MCG Health
Recovery Facility Care
30th Edition

- Clinical Indications for Admission to Inpatient Rehabilitation Facility (Acute Rehabilitation)
- Evaluation and Treatment
 - General Treatment Course
 - Rehabilitation Treatment Plan
 - Physical Therapy Treatment Plan
 - Occupational Therapy Treatment Plan
 - Patient Education
- Discharge Planning
- Quality Measures
- Evidence Summary
 - Criteria
 - Treatment
 - Care Coordination
 - Rationale
 - Related CMS Coverage Guidance
- References
- Footnotes
- Definitions
- Codes

Clinical Indications for Admission to Inpatient Rehabilitation Facility (Acute Rehabilitation)

- **Inpatient rehabilitation facility** (acute rehabilitation) admission may be indicated by **ALL** of the following[A](1)(3):[N](#)
 - No acute hospital care needs
 - Preadmission screening completed; assessment examples include[B][C](1):
 - Level of function prior to the event
 - Condition that caused the need for rehabilitation
 - Expected level of improvement
 - Expected length of time to achieve level of improvement

- Risk for clinical complications
- Combination of treatments needed (PT, OT, SLP, or orthosis/prosthesis)
- Anticipated discharge destination
- Patient status supports **active** participation in, and benefit from, intensive rehabilitation Skilled care that includes **ALL** of the following^{[D](3)(7)}:
 - Intensity and complexity of nursing, medical management, and rehabilitation needs require and can be expected to benefit from an inpatient stay and interdisciplinary team approach.^[N]
 - Ability to actively participate in Intensive rehabilitation therapy program
 - Patient can reasonably be expected to demonstrate Goals for measurable improvement.
 - Requires physician supervision to assess the patient both medically and functionally^[E]
 - Requires an intensive and coordinated Interdisciplinary team care approach^[F]
- Therapy services needed, including **ALL** of the following^{(15)(16):[N]}
 - Skilled therapy services needed for **1 or more** of the following:
 - Augmentative and alternative communication device training
 - Edema-management techniques⁽¹⁹⁾
 - Environmental and activity modification for ADL performance⁽²⁰⁾
 - Equipment and adaptive technology use and safety⁽¹⁹⁾
 - Fall-prevention and recovery training⁽²¹⁾
 - Functional activities evaluation and training
 - Gait evaluation and training⁽²²⁾
 - Motor control and balance evaluation and training
 - Neurobehavioral and neurocognitive assessment and treatment strategies (eg, learning, memory, attention, problem-solving, abstract reasoning, and impulsivity)⁽²³⁾
 - Orthosis or prosthesis evaluation and training⁽²⁴⁾⁽²⁵⁾
 - Pain-management techniques⁽¹⁹⁾⁽²⁶⁾
 - Positioning techniques and training⁽²⁷⁾
 - Range of motion evaluation and training⁽¹⁹⁾
 - Restoration of speech or swallowing with services of speech-language pathologist (SLP)⁽²⁸⁾
 - Safety awareness training related to functional mobility and ADL performance⁽²³⁾
 - Strength evaluation and training
 - Therapeutic exercises or activities supervision to ensure patient safety and treatment effectiveness
 - Multidisciplinary therapy services needed, including **2 or more** of the following⁽²⁹⁾:
 - Physical therapy (PT)
 - Occupational therapy (OT)
 - Other therapy, including **1 or more** of the following:
 - Speech-language pathology (SLP) therapy
 - Orthosis or prosthesis
- Nursing services needed, including **1 or more** of the following⁽⁷⁾⁽³⁰⁾:
 - Behavioral health condition monitoring⁽³¹⁾
 - Bowel and bladder management⁽³²⁾⁽³³⁾⁽³⁴⁾
 - Coordination of care between interdisciplinary team members⁽¹⁹⁾
 - Infusion therapy management
 - Medical condition monitoring with evaluation of rehabilitation on clinical status
 - Musculoskeletal management⁽³⁵⁾

- Nutritional management(36)
- Respiratory management
- Therapy skills incorporated and reinforced in nursing care(27)
- Time-tabling and routine establishment (eg, for ADL adaptation)
- Pain management(26)
- Patient and caregiver education (eg, safety, skin care, medication)(27)
- Wound management

Evaluation and Treatment

General Treatment Course

Stage	Clinical Status	Interventions
1	• Clinical indications for admission met • Inpatient Rehabilitation Admission Assessment complete with intensive rehabilitation needs and services identified and evaluation of current level of functioning (eg, Inpatient Rehabilitation Facility Patient Assessment Instrument (IRF-PAI)) QM [G][H] • Activities of Daily Living (ADL) and Instrumental Activities of Daily Living (IADL) Assessment  SR complete for evaluation of current level of functioning (eg,  Activities of Daily Living (ADL) Scoring Tool Calculator) QM • Physical ability assessment (eg, FIM Score®, Functional Independence Measure for children (WeeFIM®)) • Pressure injury assessment (ie, risk for new or presence of existing pressure injury). See Wounds Due to Surgery, Injury, or Pressure  SR for further information. QM • Psychosocial status assessment. See Psychosocial Assessment  SR for further information. QM • Social determinants of health screening. See Social Determinants of Health Screening Tool  SR for further information. QM(40) • Hospital readmission risk factor evaluation QM (41)(42)	• Transition of care planning initiated with evaluation for next level of care QM (43) • Interdisciplinary care plan established and implemented QM [I](3)(44)  Provide treatments and monitoring: ◦ Chronic condition management as indicated ◦ Fall risk management QM (45) ◦ Fluid and electrolyte management(46) ◦ Infusion (IV or subcutaneous) therapy management (eg, fluids, antibiotics, parenteral nutrition)(47) ◦ Laboratory testing or monitoring, if ordered(46) ◦ Medication management QM (48) ◦ Nutrition management(49) ◦ Pain management ◦ Pressure injury management QM (50) ◦ Thromboprophylaxis(51)(52) ◦ Wound or dressing management(53) • Education for self-care(54) • Physical therapy (PT) evaluation • Occupational therapy (OT) evaluation • Speech-language pathology (SLP) evaluation • Neuropsychological (eg, neurocognitive and neurobehavioral) evaluation, if indicated(23) • Orthosis or prosthesis evaluation, if indicated(16)

- Social work or case management evaluation(1)(3)(55)

2

- **Intensive rehabilitation therapy program established and patient actively participating[J]**
- Unexpected clinical event resolved (eg, infection, DVT, pneumonia)[K]
- Patient demonstrating measurable practical improvement in functional condition with evaluation of current level of functioning (eg,  Activities of Daily Living (ADL) Scoring Tool Calculator) **QM**(56)
- **Physical therapy (PT) progression**
- **Occupational therapy (OT) progression**
- **Speech-language pathology (SLP) progression, if indicated**
- **Other care needs stable or diminished**



- Interdisciplinary care plan continues **QM**
- Ongoing assessment of clinical needs **QM**
- Provide treatments and monitoring:
 - Chronic condition management as indicated
 - Fall risk management **QM**
 - Fluid and electrolyte management
 - Infusion (IV or subcutaneous) therapy management (eg, fluids, antibiotics, parenteral nutrition)
 - Laboratory testing or monitoring, if ordered
 - Medication management **QM**
 - Nutrition management
 - Pain management
 - Pressure injury management **QM**
 - Thromboprophylaxis
 - Wound or dressing management
- Ongoing patient or caregiver education
- Ongoing equipment (eg, DME) training
- Ongoing rehabilitation therapy(3)
- Ongoing neuropsychological management, if indicated
- Orthosis or prosthesis training, if indicated
- Transition of care planning continues with evaluation for next level of care **QM** (57)

3

- **Adequate intensive inpatient rehabilitation completed for safe transfer to next level of care, or patient's condition prevents active participation in intensive therapy program, or patient no longer demonstrating significant functional gains (eg,  Activities of Daily Living (ADL) Scoring Tool Calculator)**
- **Equipment (eg, DME) training completed**
- **Gastrointestinal status stable or manageable at lower level of care**
- **Medical status stable or manageable at lower level of care (eg, vital signs, laboratory values)**
- **Neurobehavioral and neurocognitive status manageable at lower level of care**
- **Neurovascular status stable or manageable at lower level of care**
- **Orthosis or prosthesis training completed or manageable at lower level of care**
- **Pain controlled or manageable at lower level of care**

- **Transition of care planning completed **QM****
- Safe to go home and transition to community or transition to alternative level of care (eg, skilled nursing facility, home care) **QM**

- Patient and caregiver education completed or manageable at lower level of care
- Patient transitioned to another level of care due to unexpected clinical event (eg, DVT, infection)
- **Wound(s) or dressing changes manageable at lower level of care**
- **Transition plans and education understood**

Recovery Milestones are indicated in **bold**.

Rehabilitation Treatment Plan

Physical Therapy Treatment Plan

- **Physical therapy** treatment plan includes(58):

☐ Goals of physical therapy(59)

- Activity tolerance and functional endurance improved
- ADL/IADL performance, independence, and safety optimized
- ADL/IADL restrictions, precautions, or activity guidelines understood
- Awareness of community resources optimized
- Chronic condition modifiable risk factor management strategies maximized(60)
- Durable medical equipment usage optimized
- Energy conservation techniques and proper body mechanics maximized
- Exercise tolerance and functional endurance improved
- Fall-risk-reduction strategies maximized
- Functional independence and safety with mobility-related tasks maximized
- Home rehabilitation program and recommendations understood
- Orthosis or prosthesis management and independence maximized(61)(62)(63)(64)(65)
- Pain-management techniques optimized
- Potential personality or emotional changes understood
- Pressure injury and contracture risk prevention maximized
- Respiratory status and breathing techniques improved(66)
- Risk and disability associated with disease, injury, or surgical procedure understood
- Self-management of health, injury, surgical procedure, or disease process optimized
- Self-reported quality of life improved
- Strength and range of motion improved
- Postdischarge complication risk factors minimized

☐ Rehabilitation techniques and training

- ADL training
- Age-appropriate functional strategies and techniques in alignment with developmental progression(67)
- Amputation recovery, including residual limb care and preparation for prosthesis(68)
- Compression bandage or garment management(69)
- Contracture prevention or management(70)
- Edema prevention or management(71)

- Education of therapeutic program with patient and caregiver(72)
- Energy conservation and work simplification(73)
- Equipment and device use training(62)(74)
- Exercise program setup and progression within precautions, which may include(15)(75)(76):
 - Aerobic exercises or endurance conditioning
 - Balance training
 - Coordination exercises
 - Range-of-motion techniques
 - Respiratory exercises and training (eg, respiratory muscle training, coughing techniques, deep breathing exercises, incentive spirometry exercises, pursed lip and diaphragmatic breathing)(66)(77)
 - Strengthening exercises(60)
 - Stretching exercises
- Fall-prevention and recovery training **QM** (78)(79)
- Functional skills training with applicable durable medical equipment (eg, walker, transfer board)(80)
 - Bed mobility training
 - Gait training on level and uneven surfaces
 - Stair management
 - Transfer training
 - Wheelchair mobility and management(81)(82)(83)
- Joint-protection techniques and mobilization(84)
- Lifting as prescribed for diagnosis (eg, medical, surgical)(77)
- Manual therapy (eg, soft tissue mobilization techniques, scar management, massage)(85)
- Neuromuscular re-education (eg, proprioception training)
- Pain-management strategies and techniques(86)(87)
- Posture, positioning, and body mechanics as prescribed for diagnosis or surgical approach(88)
- Pressure injury prevention or management(89)
- Safety awareness education(90)
- Self-management of health, injury, surgical procedure, or disease process(72)(91)
- Sensory stimulation and integration techniques
- Splint, brace, cast, prosthesis, or orthosis management(92)(93)
- Therapeutic physical modality use and training, as indicated(88)(94)
 - Cold modalities or cryotherapy with precautions related to circulatory deficiency, areas of desensitization, open wounds, or fractures
 - Functional or neuromuscular electrical stimulation(95)
 - Mechanical traction(96)
 - Thermal modality with precautions related to circulatory deficiency, areas of desensitization, open wounds, or fractures
 - Transcutaneous electrical nerve stimulation (TENS)(66)
- Weight-bearing as prescribed for diagnosis and/or surgical approach(84)(92)

☐ Discharge recommendations, which may include:

- Activity modifications
- Environmental adaptation
- Home exercise program as prescribed
- Home modifications

Occupational Therapy Treatment Plan

- **Occupational therapy** treatment plan includes(97)(98)(99):

☐ Goals of occupational therapy(100)

- Activity tolerance and functional endurance improved
- ADL/IADL performance, independence, and safety optimized
- ADL/IADL restrictions, precautions, or activity guidelines understood
- Balance and body alignment for ADL/IADL improved
- Fall-risk-reduction strategies maximized
- Home rehabilitation program and recommendations understood
- Pain-management techniques optimized
- Proper posture for sitting, ADL, and transfers maximized
- Potential personality or emotional changes understood
- Upper body range of motion, strength, and functional use for ADL or IADL maximized
- Upper extremity joint range of motion for functional tasks maximized
- Upper extremity motor control improved
- Postdischarge complication risk factors minimized

☐ Rehabilitation techniques and training(101)

- ADL or IADL training
- Behavior-management strategies and techniques (eg, cognitive behavioral therapy, cognitive rehabilitation techniques)(102)
- Compensatory strategies for completion of ADL/IADL
 - Cognition impairment management(103)
 - Energy conservation(24)
 - One-handed techniques
 - Perceptual deficits management
 - Sensory impairment management
 - Visual impairment management
 - Work simplification
- Compression garment application and use
- Contracture prevention or management(70)
- Edema prevention or management
- Education of therapeutic program with patient and caregiver
- Environmental and activity modification for ADL/IADL performance(104)
- Equipment and device use training
- Exercise program setup and progression within precautions, which may include:
 - Balance training
 - Coordination exercises
 - Range-of-motion techniques
 - Strengthening exercises
 - Stretching exercises
- Fall-prevention and recovery training **QM** (78)
- Functional skills training with applicable durable medical equipment (eg, shower chair, commode)(105)
 - Bed mobility training

- Transfer training
- Wheelchair mobility and management(81)
- Joint-protection techniques and mobilization
- Lifting as prescribed for diagnosis (eg, medical, surgical)
- Manual therapy (eg, passive soft tissue mobilization techniques, scar management, massage)(85)
- Neuromuscular re-education (eg, proprioception training)(106)
- Pain-management strategies and techniques
- Posture, positioning, and body mechanics as prescribed for diagnosis or surgical approach
- Safety awareness education
- Sensory stimulation and integration techniques
- Splint, brace, cast, prosthesis, or orthosis management(24)
- Stress-management techniques(107)
- Therapeutic physical modality (eg, heat or ice packs) use and training, as indicated(108)
- [-] Discharge recommendations, which may include(109):
 - Activity modifications
 - Environmental adaptation
 - Home exercise program
 - Home modifications
 - Pre-driver assessment and automobile modification
 - Vocational rehabilitation(107)

Patient Education

- Patient and caregiver education. See Inpatient Rehabilitation Facility (Acute Rehabilitation): Musculoskeletal: Patient Education for Clinicians [SR](#).

Discharge Planning

- Transition of care planning may include(16):[NN](#)
 - [-] Assessment and early establishment of anticipated discharge destination and plans for care, including(117):
 - Treatment plan development involving multiple providers
 - Premorbid functioning
 - Patient and caregiver preferences and abilities
 - Evaluation for skilled services at next level of care as appropriate for patient's continued needs
 - Psychosocial status. See Psychosocial Assessment [SR](#) for further information. **QM**(118)
 - Social determinants of health screening. See Social Determinants of Health Screening Tool [SR](#) for further information. **QM**(40)
 - Acceptance for care for patient at next level of care, as appropriate
 - o Transition of care plan complete, which may include:
 - Patient and caregiver education complete. See Teach-Back Tool [SR](#) for further information.
 - [-] Medication reconciliation completion includes **QM** (119)(120):
 - Compare patient's discharge list of medications (prescribed and over-the-counter) against provider's admission or transfer orders.
 - Assess each medication for correlation to disease state or medical condition.

- Report medication discrepancies to prescribing provider, attending physician, and primary care provider, and ensure accurate medication order is identified.
 - Provide reconciled medication list to all treating providers.
 - Confirm that patient or caregiver can acquire medication.
 - Educate patient and caregiver.
 - Provide complete medication list to patient and caregiver.
 - Importance of presenting personal medication list to all providers at each care transition, including all provider appointments
 - Reason, dosage, and timing of medication (eg, use "teach-back" techniques)(121)(122)
 - Encourage communication between patient, caregiver, and pharmacy for obtaining prescriptions, setting up home medication delivery, and reviewing for drug-drug interactions.
 - See Medication Reconciliation Tool [SR](#) for further information.
- ☐ Appointments planned or scheduled(57):
- Primary care provider
 - Physiatrist or rehabilitation provider
 - Prostheses or orthoses provider
 - Rehabilitation therapy (eg, PT) at next level of care(57)
 - Specialists for management of comorbidities as needed
 - Surgeon
 - Wound, ostomy, and continence nurse (WOCN)
 - Other
- ☐ Referrals made for assistance or support, which may include:
- Alcohol and other drug abuse or dependence treatment, as needed **QM**
 - Behavioral health support (eg, counseling)
 - Community services(123)(124)
 - Educational program (eg, chronic condition management, self-management)(121)
 - Financial, for follow-up care, medication, and transportation
 - Home architectural modifications
 - Home healthcare(125)
 - Local electric or power company regarding need for life support if ventilator needed at home
 - Protective services or legal authorities, if indicated
 - Self-help or support groups
 - Social services (eg, social programs, advance directives)
 - Tobacco use treatment, as needed **QM** (126)(127)
 - Vocational rehabilitation(107)
 - Other
- ☐ Medical equipment and supplies coordinated (ie, delivered or delivery confirmed), as indicated(128):
- ADL-supportive equipment
 - Cardiac care equipment and supplies (eg, compression stockings)
 - Medication equipment and supplies
 - Nutritional care equipment and supplies
 - Orthopedic equipment
 - Respiratory equipment and supplies

- Wound care equipment and supplies
- Other
- Transition plan communicated to all members of patient's care team **QM**

Quality Measures

- HEDIS® Measures include(43):
 - Medication reconciliation is completed.
 - Transition of care needs are assessed and addressed.
 - Psychosocial status (eg, depression screening) is assessed and addressed.
 - Fall risk is assessed and addressed.
 - Tobacco use is assessed and addressed.
 - Immunization status is assessed and addressed.
 - Readmission risk is assessed and addressed.
 - Alcohol and substance misuse is assessed and addressed.
 - Social needs screening and intervention is completed.
- Inpatient Rehabilitation Facility Compare Quality Measures include(129):
 - Pressure injury risk is assessed and addressed.
 - ADL status is assessed for improvement.
 - Fall risk is assessed and addressed.
 - Transfer of health information to provider and patient is completed.
 - Healthcare-associated infection risk is reduced
 - Medication reconciliation is completed.
 - Discharge to the community is completed.
 - Readmission risk is assessed and addressed.
- Inpatient Rehabilitation Facility Quality Reporting Program measures include(129):
 - Pressure injury risk is assessed and addressed.
 - ADL status is assessed for improvement.
 - Fall risk is assessed and addressed.
 - Transfer of health information to provider and patient is completed.
 - Readmission risk is assessed and addressed.
 - Medication reconciliation is completed.
 - Immunization status is assessed and addressed.
 - Discharge to the community is completed.

Evidence Summary

Criteria

The evidence for the clinical indications found in this guideline includes 10 published peer reviewed articles and 20 book sections.

A retrospective review of over 1.9 million inpatient rehabilitation patients discharged between January 2016 and December 2019 showed that 70% of patients had at least one ICD-10 pain diagnosis; pain diagnosis was common among all impairment groups, affecting half of patients in each group, with the highest prevalence in

the arthritis and pain syndromes impairment group.(4) **(EG 2)**

Treatment

A retrospective study of 479 Medicare patients who received rehabilitation in a nursing home or inpatient rehabilitation facility (IRF) setting found that 1 in 3 patients reported no functional improvement during rehabilitation. Factors associated with greater odds of reporting no improvement included high school education or less, impaired instrumental activities of daily living, less than 1 month total duration of rehabilitation, and no outpatient rehabilitation services.(8) **(EG 2)** A retrospective review of 343,193 Medicare patients discharged from acute inpatient care to an IRF found that those who received vertically integrated care in a hospital-based IRF had shorter inpatient and IRF lengths of stay with maintained or improved health outcomes, fewer readmissions, and were more likely to be discharged to home with home health services when compared with patients who received care in a freestanding IRF (ie, IRF with hospitals within the same health system but not hospital-based).(9) **(EG 2)** A systematic review of 24 articles including adult patients in post-acute care (eg, IRFs, long-term care hospitals, skilled nursing facilities, or home health settings) found that occupational therapy interventions may reduce or manage preventable adverse events. Strong strength of evidence was found to support an exercise program to reduce risk of falls in an IRF. Moderate strength of evidence was found to support facility-wide pressure injury reduction programs to reduce pressure injuries; dysphagia management through multidisciplinary care and strengthening exercises for swallowing; and a multidisciplinary, multicomponent falls program.(10) **(EG 1)** A retrospective analysis of 5,033,820 Medicare patients discharged from acute care to post-acute care found that those discharged to an IRF were least likely to demonstrate cognitive impairment, use a bowel appliance, or be on a therapeutic diet, and had the shortest average length of stay when compared with other post-acute settings.(11) **(EG 2)** A retrospective cohort study of 270 medically complex adult patients, including those with cardiac, pulmonary, and orthopedic conditions who did not have a Centers for Medicare and Medicaid Services-compliant condition were shown to greatly benefit from inpatient rehabilitation. A total of 71.9% were discharged to home, 15.9% were discharged to a SNF, and 12.2% for readmission to acute care, with a mean rehabilitation length of stay of 12.6 days, mean total Functional Independence Measure (FIM®) gain of 26, and mean FIM® efficiency of 2.3.(12) **(EG 2)** A retrospective cohort study of 270 medically complex inpatient rehabilitation patients reported that patients with ICU-acquired weakness (ICUAW) had significantly greater Functional Independence Measure (FIM®) motor gain than medically complex patients without ICUAW and were 81% more likely to be discharged to home vs to a skilled nursing facility.(13) **(EG 2)** A retrospective observational study performed at a 178-bed inpatient rehabilitation hospital reported that of the 1008 patients who received a psychiatric consultation during their inpatient rehabilitation hospital stay, the most common conditions associated with a psychiatric consult were amputations (38%), burns (35%), neurologic disorder (28%), deconditioning (14%), and musculoskeletal injury (7%). Although 20% of patients did not meet criteria for psychiatric diagnosis, the most common diagnoses included mood disorder, adjustment disorder, neurocognitive disorder, and delirium.(14) **(EG 2)**

A claims-based study of 1.4 million Medicare beneficiaries discharged from an acute care hospital to a post-acute care setting, including an inpatient rehabilitation facility (IRF), a skilled nursing facility (SNF), or home care (HC), found that more hours of occupational therapy and physical therapy services were associated with greater functional improvements, and fewer hours of occupational therapy and physical therapy services were associated with a higher risk for 30-day hospital readmission across all post-acute care settings.(17) **(EG 2)** A retrospective study of 3212 patients who were discharged from an IRF found that increased therapy hours were associated with increased functional gains among all types of patients.(18) **(EG 2)**

Care Coordination

An observational review of 567,850 inpatient rehabilitation facility (IRF) Medicare inpatient stays in 1166 facilities during 2013 to 2014 found that adjusted hospital readmission rate 30 days post-IRF discharge was in the range of 12.4% to 13.1%, but almost 75% of facilities demonstrated readmission rates that were significantly different than this average. Mean number of days to readmission was 13.0, which varied significantly by rehabilitation diagnosis.(110) **(EG 2)** A retrospective cohort study of a Medicare claims database involving 371,846 patients discharged from inpatient rehabilitation from 2013 to 2014 found that functional independence at discharge was associated with lower rates of potentially preventable hospital readmissions. Readmission rates for highest vs lowest quartiles within each functional domain were 3.4% vs 6.9% for self-care, 3.3% vs 7.2% for mobility, and 3.5% vs 6.2% for cognition; 44.1% of readmissions were for infection-related conditions, 31.2% were for inadequate management of chronic conditions, and 24.7% were for inadequate management of other unplanned events.(111) **(EG 2)** A retrospective cohort analysis of 539 geriatric patients discharged from acute care to geriatric rehabilitation found that predictors of acute readmission include severe infection, fracture secondary to falls, intra-abdominal complications, cardiac complications, and acute neurologic events.(112) **(EG 2)** An observational cohort study of 682 geriatric patients in an IRF found that risk factors for 30-day and 90-day readmissions included a high fear of falling and higher

Charlson Comorbidity Index and Cumulative Illness Rating Scale (CIRS) scores.(42) **(EG 2)** A retrospective analysis of 1,102,785 patient discharges from 944 IRFs found that patient factors associated with 30-day readmissions to acute care included older age, male sex, White race, married, Medicare insured, longer duration of impairment, lower admission FIM® cognitive and motor scores, more comorbidities, dysphagia, pneumonia, and weekend admission to the IRF.(41) **(EG 2)** A retrospective analysis of 3294 patients age 55 years or older discharged from inpatient acute care to either home without home healthcare, home with home healthcare, or an IRF found that most readmissions and emergency department/observation visits within 24 hours came from IRFs. The study also found that patient characteristics associated with readmission from an IRF included younger age, less chronic illness burden, and greater and more recent functional decline. (113) **(EG 2)** A retrospective study of 75 patients discharged from an acute IRF between 2013 to 2018 who experienced a potentially preventable readmission (PPR) found that individuals were more likely to be readmitted with a PPR if they had a greater number of comorbidities, were admitted with a spinal cord injury, or had lower admission or discharge FIM® motor scores. The most common PPR diagnoses were sepsis, renal failure, respiratory problems, and urinary tract infection. (114) **(EG 2)** An evidence-based improvement study of 79 patients discharged from a 28-bed IRF showed implementing the teach-back intervention from the Agency for Healthcare Research and Quality (AHRQ) Health Literacy Universal Precautions Toolkit into the discharge process resulted in a mean 45% fewer 30-day readmissions.(115) **(EG 2)**

A longitudinal observational study of 43 adult inpatient rehabilitation facility patients who were not safe to return home at discharge and required a skilled rehabilitation facility placement found that 58.2% of patients returned to baseline independence, and 63.4% of patients who lived at home prior to acute hospitalization returned home.(116) **(EG 2)**

Rationale

Inpatient rehabilitation facilities are designed to provide intensive rehabilitation therapy in a resource intensive inpatient hospital environment. Use of this MCG care guideline helps in the decision to appropriately admit a patient to an acute inpatient rehabilitation facility and assess progress through an interdisciplinary plan of care that incorporates interventions for skilled nursing (eg, wound care, pain management, infusion medication) and rehabilitation therapy (eg, physical, occupational, speech-language) in alignment with CMS requirements.

Use of these evidence-based clinical criteria benefits patients by assisting in the identification of patients who, due to the complexity of their nursing, medical management, and rehabilitation needs, require and can reasonably be expected to benefit from an inpatient stay and an interdisciplinary team approach to the delivery of rehabilitation care. This evidence-based guidance ensures that Medicare and Medicare Advantage patients receive full access to Medicare benefits without risk of delay or decreased access. Additional evidence-based guidance benefits patients by outlining complicated factors (eg, pain exacerbations, infection, comorbid disease exacerbation) that could extend the need for a rehabilitation facility stay. Criteria for transitioning a patient to the next appropriate level of care (eg, acute care, skilled nursing facility, home care) are also provided to assist providers and patients in identifying next steps for the patient's care. This support can increase consistency in treatment thresholds, leading to less variation in care and promoting equitable treatment among patients.

Related CMS Coverage Guidance

This guideline supplements but does not replace, modify, or supersede existing Medicare regulations or applicable National Coverage Determinations (NCDs) or Local Coverage Determinations (LCDs).

Code of Federal Regulations (CFR): 42 CFR 410.38(128); 42 CFR 412.29(2); 42 CFR 412.606(38); 42 CFR 412.610(39); 42 CFR Pt. 412.622(1)

Internet-Only Manual (IOM) Citations: CMS IOM Publication 100-02, Medicare Benefit Policy Manual, Chapter 1 - Inpatient Hospital Services Covered Under Part A(3); CMS IOM Publication 100-08, Medicare Program Integrity Manual, Chapter 6 - Medical Review of Inpatient Hospital Claims for Part A(130)

Medicare Coverage Determinations: Medicare Coverage Database(131)

Other CMS Guidance: Centers for Medicare and Medicaid Services, "Inpatient Rehabilitation Facility Patient Assessment Instrument (IRF-PAI) and IRF-PAI Manual" Version 4.2(37)

References

1. Centers for Medicare and Medicaid Services. "Basis of payment." 42 CFR Pt. 412.622 Washington, DC 2025 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>. [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9]
2. Centers for Medicare and Medicaid Services. "Classification criteria for payment under the inpatient rehabilitation facility prospective payment system." 42 CFR Pt. 412.29 Washington, DC 2020 Aug 10 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/current/title-42/section-412.29>. [Context Link 1, 2]
3. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09] [Context Link 1, 2, 3, 4, 5, 6, 7, 8, 9, 10]
4. Curry ZA, et al. Examination of pain comorbid diagnoses in the inpatient rehabilitation population across all impairment groups. *American Journal of Physical Medicine and Rehabilitation* 2024;Online. DOI: 10.1097/PHM.0000000000002512. [Context Link 1] View abstract...
5. Centers for Medicare and Medicaid Services. "Criteria for skilled services and the need for skilled service." 42 CFR Pt. 409.32 Washington, DC 2023 Oct 02 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
6. Centers for Medicare and Medicaid Services. "Skilled services requirements." 42 CFR Pt. 409.44 Washington, DC 2023 Oct 02 [accessed 2025 Jul 23] Accessed at: <https://www.ecfr.gov/>. [Context Link 1]
7. Pryor J, O'Reilly K, Bonser M, Garret G, McKenchnie D. Rehabilitation for the individual and family. In: Chang E, Johnson A, editors. *Living With Chronic Illness and Disability*. 4th ed. Chatswood NSW 2067: Elsevier; 2022:161-182. [Context Link 1, 2]
8. Simning A, Caprio TV, Seplaki CL, Temkin-Greener H, Szanton SL, Conwell Y. Patient-reported outcomes in functioning following nursing home or inpatient rehabilitation. *Journal of the American Medical Directors Association* 2018;19(10):864-870. DOI: 10.1016/j.jamda.2018.06.014. [Context Link 1] View abstract...
9. Sood N, Shier V, Huckfeldt PJ, Weissblum L, Escarce JJ. The effects of vertically integrated care on health care use and outcomes in inpatient rehabilitation facilities. *Health Services Research* 2021;56(5):828-838. DOI: 10.1111/1475-6773.13667. [Context Link 1] View abstract...
10. Hunter EG, Rhodus E. Interventions within the scope of occupational therapy to address preventable adverse events in inpatient and home health postacute care settings: a systematic review. *American Journal of Occupational Therapy* 2022;76(1):7601180060. DOI: 10.5014/ajot.2022.047589. [Context Link 1] View abstract...
11. Rodriguez A, et al. Who receives post-acute care? Characteristics of national population and field test sample. *Journal of the American Geriatrics Society* 2022;70(4):991-1000. DOI: 10.1111/jgs.17649. [Context Link 1] View abstract...
12. Zhang S, Lin D, Wright ME, Swallow N. Acute inpatient rehabilitation improves function independent of comorbidities in medically complex patients. *Archives of Rehabilitation Research and Clinical Translation* 2022;4(2):Online. DOI: 10.1016/j.arrct.2022.100178. [Context Link 1] View abstract...
13. Rudra RT, Lin D, Miller B, Du P, Zhang S. Investigating inpatient rehabilitation outcomes of patients with intensive care unit-acquired weakness, and identifying comorbidities associated with unfavorable outcomes. *PM & R: The Journal of Injury, Function, and Rehabilitation* 2022;14(2):190-197. DOI: 10.1002/pmrj.12565. [Context Link 1] View abstract...
14. Daly N, Steinberg R, Boyle MT, Brenkel M, Robinson L. An overview of the rehabilitation and psychiatric diagnoses of patients referred to a psychiatry consult liaison service at an inpatient rehabilitation hospital. *PM & R: The Journal of Injury, Function, and Rehabilitation* 2023;15(10):1273-1279. DOI: 10.1002/pmrj.12943. [Context Link 1] View abstract...
15. Tan Dy RC, Nguyen N. Therapeutic exercise. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:145-154. [Context Link 1, 2]
16. Medical-surgical nursing. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:33-55. [Context Link 1, 2, 3]
17. Therapy Outcomes in Post-Acute Care Settings (TOPS) Study Chartbook. [Internet] American Occupational Therapy Association, Inc and American Physical Therapy Association. 2021 Apr Accessed at: <https://www.aota.org/>. [accessed 2025 Sep 05] [Context Link 1]

18. Roberts PS, et al. Retrospective study demonstrating therapy time impact on inpatient rehabilitation functional gains. *Disability and Rehabilitation* 2022;44(17):4639-4647. DOI: 10.1080/09638288.2021.1912836. [Context Link 1] View abstract...
19. Musculoskeletal care plans. In: Gulanick M, Myers JL, Bowman-Woodall C, editors. *Nursing Care Plans*. 11th ed. Elsevier; 2025:595-640. [Context Link 1, 2, 3, 4, 5]
20. Parcetch KM, Schmitz TJ. Examination and modification of the environment. In: Fulk G, Chui K, editors. *O'Sullivan & Schmitz's Physical Rehabilitation*. 8th ed. USA: F.A. Davis Company; 2024:292-331. [Context Link 1]
21. Kilma DW, Schmitz TJ, O'Sullivan SB. Examination of coordination and balance. In: Fulk G, Chui K, editors. *O'Sullivan & Schmitz's Physical Rehabilitation*. 8th ed. USA: F.A. Davis Company; 2024:180-214. [Context Link 1]
22. Gait and Posture Analysis. In: Dutton M, editor. *Dutton's Orthopaedic Examination, Evaluation, and Intervention*. 6th ed. McGraw-Hill Education; 2023:272-318. [Context Link 1]
23. Gaughwin PM, Cameron ID. Psychological assessment and intervention in rehabilitation. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:29-36. [Context Link 1, 2, 3]
24. McMillan IR. The biomechanical frame of reference. In: Duncan EAS, editor. *Foundations for Practice in Occupational Therapy*. 6th ed. Elsevier; 2021:152-164. [Context Link 1, 2, 3]
25. Wong CK, Lee DJ. Orthotics. In: Fulk G, Chui K, editors. *O'Sullivan & Schmitz's Physical Rehabilitation*. 8th ed. USA: F.A. Davis Company; 2024:1149-1178. [Context Link 1]
26. Pain management. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. *Clinical Nursing Skills and Techniques*. 11th ed. Elsevier; 2025:443-497. [Context Link 1, 2]
27. Snyder JS, Sump CA. Fractures. In: Snyder JS, Sump CA, editors. *Swearingen's All-in-One Nursing Care Planning Resource*. 6th ed. Elsevier; 2024:495-502. [Context Link 1, 2, 3]
28. Hsiao MY, Wang TG. Rehabilitation of swallowing disorders. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:23-28. [Context Link 1]
29. Makam AN, Grabowski DC. Policy in clinical practice: choosing post-acute care in the new decade. *Journal of Hospital Medicine* 2021;16(3):171-174. DOI: 10.12788/jhm.3577. [Context Link 1] View abstract...
30. Kilgore C. Nursing patients through rehabilitation. In: O'Hanlon S, Smith M, editors. *Comprehensive Guide to Rehabilitation of the Older Patient*. 4th ed. Elsevier; 2021:69-72. [Context Link 1]
31. Preston C. Psychological input. In: O'Hanlon S, Smith M, editors. *Comprehensive Guide to Rehabilitation of the Older Patient*. 4th ed. Elsevier; 2021:80-84. [Context Link 1]
32. Varma AK, Mishra S. Neurogenic lower urinary tract dysfunction. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:191-200. [Context Link 1]
33. Oey NE, Lim PA. Neurogenic bowel: Dysfunction and rehabilitation. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:201-210. [Context Link 1]
34. Lee R, Gibson W, Wagg A. Continence and elimination. In: O'Hanlon S, Smith M, editors. *Comprehensive Guide to Rehabilitation of the Older Patient*. 4th ed. Elsevier; 2021:195-202. [Context Link 1]
35. Management of patients with musculoskeletal disorders. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:1113-1151. [Context Link 1]
36. Nutrition assessment and patient care. In: Grodner M, Escott-Stump S, Dorner S, editors. *Nutritional Foundations and Clinical Applications*. 8th ed. Elsevier; 2023:217-247. [Context Link 1]
37. Centers for Medicare and Medicaid Services. "Inpatient Rehabilitation Facility Patient Assessment Instrument (IRF-PAI) and IRF-PAI Manual." Version 4.2 Washington, DC 2024 Feb 01 [accessed 2025 Jul 23] Accessed at: <https://www.cms.gov/medicare/quality/inpatient-rehabilitation-facility/irf-pai-and-irf-qrp-manual>. [Context Link 1, 2]
38. Centers for Medicare and Medicaid Services. "Patient assessments." 42 CFR Pt. 412.606 Washington, DC 2023 Aug [accessed 2025 Jul 23] Accessed at: <https://www.ecfr.gov/>. [Context Link 1, 2]
39. Centers for Medicare and Medicaid Services. "Assessment schedule." 42 CFR Pt. 412.610 Washington, DC 2025 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>. [Context Link 1, 2]

40. Imran A, Rawal MD, Botre N, Patil A. Improving and promoting social determinants of Health at a system level. *Joint Commission Journal on Quality and Patient Safety* 2022;48(8):376-384. DOI: 10.1016/j.jcjq.2022.06.004. [Context Link 1, 2] View abstract...
41. Shea CA, et al. Variation in 30-day readmission rates from inpatient rehabilitation facilities to acute care hospitals. *Journal of the American Medical Directors Association* 2021;22(12):2461-2467. DOI: 10.1016/j.jamda.2021.03.033. [Context Link 1, 2] View abstract...
42. Wan CS, Reijnierse EM, Maier AB. Risk factors of readmissions in geriatric rehabilitation patients: RESORT. *Archives of Physical Medicine and Rehabilitation* 2021;102(8):1524-1532. DOI: 10.1016/j.apmr.2021.01.082. [Context Link 1, 2] View abstract...
43. HEDIS® quality measures 2026. [Internet] NCQA. Accessed at: <https://www.ncqa.org/hedis/measures/>. Updated 2026 [accessed 2026 Apr 02] [Context Link 1, 2]
44. Wong R. Care planning meetings. In: O'Hanlon S, Smith M, editors. *Comprehensive Guide to Rehabilitation of the Older Patient*. 4th ed. Elsevier; 2021:395-398. [Context Link 1]
45. Falls: assessment and prevention in older people and in people 50 and over at higher risk. NICE Guideline NG249 [Internet] National Institute for Health and Care Excellence. 2025 Apr Accessed at: <https://www.nice.org.uk/>. [accessed 2025 Aug 18] [Context Link 1]
46. Fluid and electrolytes. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:224-272. [Context Link 1, 2]
47. Intravenous and vascular access therapy. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. *Clinical Nursing Skills and Techniques*. 11th ed. Elsevier; 2025:842-894. [Context Link 1]
48. Medicare Coverage Database Local Coverage Determination documents. [Internet] Centers for Medicare and Medicaid Services. Accessed at: <https://www.cms.gov/medicare-coverage-database/search.aspx>. Updated 2024 [accessed 2025 Jan 08] [Context Link 1]
49. The client with alterations in musculoskeletal function. In: Haugen N, Galura SJ, editors. *Ulrich & Canale's Nursing Care Planning Guides, 8th Edition Revised Reprint with 2021-2023 NANDA-I Updates*. 8th ed. Elsevier; 2022:672-714. [Context Link 1]
50. European Pressure Ulcer Advisory Panel, National Pressure Injury Advisory Panel, and Pan Pacific Pressure Injury Alliance. Prevention and Treatment of Pressure Ulcers/Injuries: Clinical Practice Guideline. The International Guideline Online Living Guideline 4th Edition [Internet] National Pressure Injury Advisory Panel. 2025 Feb Accessed at: <https://npiap.com/page/InternationalGuidelines>. [accessed 2025 Sep 16] [Context Link 1]
51. Management of patients with musculoskeletal trauma. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:1152-1205. [Context Link 1]
52. Snyder JS, Sump CA. Venous thrombosis/thrombophlebitis. In: Snyder JS, Sump CA, editors. *Swearingen's All-in-One Nursing Care Planning Resource*. 6th ed. Elsevier; 2024:191-195. [Context Link 1]
53. Skin integrity and wound care. In: Lynn P, editor. *Taylor's Clinical Nursing Skills*. 6th ed. Wolters Kluwer; 2023:452-540. [Context Link 1]
54. Self-care management, self-advocacy, and client adherence. In: Gillingham DC, editor. *Foundations of Case Management*. Blue Bayou Press; 2021:105-112. [Context Link 1]
55. de Wilde C. Social work. In: O'Hanlon S, Smith M, editors. *Comprehensive Guide to Rehabilitation of the Older Patient*. 4th ed. Elsevier; 2021:91-94. [Context Link 1]
56. Kee YYK. Measuring progress with rehabilitation. In: O'Hanlon S, Smith M, editors. *Comprehensive Guide to Rehabilitation of the Older Patient*. 4th ed. Elsevier; 2021:111-115. [Context Link 1]
57. Admitting, transfer, and discharge. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. *Clinical Nursing Skills and Techniques*. 11th ed. Elsevier; 2025:40-51. [Context Link 1, 2, 3]
58. Nelson F. Physical medicine and rehabilitation: physical therapy and occupational therapy. In: Nelson F, editor. *A Manual of Orthopaedic Terminology*. 9th ed. Philadelphia, PA 19103-2899: Elsevier; 2023:386-396. [Context Link 1]
59. Carter WE III, Peretiatko S. Quality and outcome measures for medical rehabilitation. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:55-62. [Context Link 1]
60. Exercise testing and prescription for populations with other chronic diseases and health conditions. In: American College of Sports Medicine, editor. *ACSM's Guidelines for Exercise Testing and Prescription*. 12th ed. Wolters Kluwer; 2025:389-480. [Context Link 1, 2]

61. Liu M. Upper limb orthosis. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:95-108. [Context Link 1]
62. Mukaino M Saitoh E. Lower limb orthoses. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:109-118. [Context Link 1, 2]
63. Ozcakar L, Ata AM. Spinal orthoses. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:119-128. [Context Link 1]
64. Murphy DP, Hermansen J. Rehabilitation and prosthetic restoration in upper limb amputation. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:75-80. [Context Link 1]
65. McLaughlin MJ. Lower limb amputation and gait. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:81-94. [Context Link 1]
66. Musculoskeletal disorders. In: Nettina SM, Nelson-Tuttle C, editors. Lippincott Manual of Nursing Practice. 12th ed. WoltersKluwer; 2025:841-879. [Context Link 1, 2, 3]
67. Gordon AM, Magill RA, Paleg G. Motor learning: application of principles to pediatric rehabilitation. In: Palisano RJ, Orlin M, Schreiber J, editors. Campbell's Physical Therapy for Children. 6th ed. Elsevier; 2023:91-108. [Context Link 1]
68. Chen T, Chui KK, Yen SC, Parcetich KM, Lockard MA. Amputation. In: Fulk G, Chui K, editors. O'Sullivan & Schmitz's Physical Rehabilitation. 8th ed. USA: F.A. Davis Company; 2024:849-888. [Context Link 1]
69. Eapen BC, Oshiro RL. Vascular diseases. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:235-242. [Context Link 1]
70. Dicks MA, Verduzco-Gutierrez M. Spasticity and contracture. In: Gonzalez-Fernandez M, Schaaf S, Zumsteg JM, Perret D, Kennedy DJ, editors. Handbook of Physical Medicine and Rehabilitation. Springer Publishing Company; 2022:427-434. [Context Link 1, 2]
71. Connective tissue disorders. In: Nettina SM, Nelson-Tuttle C, editors. Lippincott Manual of Nursing Practice. 12th ed. WoltersKluwer; 2025:804-824. [Context Link 1]
72. Bezner JR, Maxwell J. Promoting health and wellness. In: Fulk G, Chui K, editors. O'Sullivan & Schmitz's Physical Rehabilitation. 8th ed. USA: F.A. Davis Company; 2024:1118-1148. [Context Link 1, 2]
73. von Geldern G, Alschuler K. Fatigue. In: Gonzalez-Fernandez M, Schaaf S, Zumsteg JM, Perret D, Kennedy DJ, editors. Handbook of Physical Medicine and Rehabilitation. Springer Publishing Company; 2022:343-346. [Context Link 1]
74. Williams K, Annaswamy T. Gait deviations. In: Gonzalez-Fernandez M, Schaaf S, Zumsteg JM, Perret D, Kennedy DJ, editors. Handbook of Physical Medicine and Rehabilitation. Springer Publishing Company; 2022:347-354. [Context Link 1]
75. Stoeckmann TM, O'Sullivan SB. Strategies to improve motor function. In: Fulk G, Chui K, editors. O'Sullivan & Schmitz's Physical Rehabilitation. 8th ed. USA: F.A. Davis Company; 2024:333-367. [Context Link 1]
76. Exercise prescription for healthy populations with special considerations. In: American College of Sports Medicine, editor. ACSM's Guidelines for Exercise Testing and Prescription. 12th ed. Wolters Kluwer; 2025:201-249. [Context Link 1]
77. Rudolphi DM. Postoperative care. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:365-384. [Context Link 1, 2]
78. Ritchey KC, Olney AL, Campen SJ. Falls. In: Gonzalez-Fernandez M, Schaaf S, Zumsteg JM, Perret D, Kennedy DJ, editors. Handbook of Physical Medicine and Rehabilitation. Springer Publishing Company; 2022:337-342. [Context Link 1, 2]
79. Rice J, et al. Gait speed modifies efficacy of home-based exercise for falls in older adults with a previous fall: secondary analysis of a randomized controlled trial. Physical Therapy 2025;DOI: 10.1093/ptj/pzaf008. [Context Link 1] View abstract...
80. Hussein N. Acute medical conditions. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:253-264. [Context Link 1]
81. Michael E, Farrell E, Linden M, Cabahug PG, Sadowsky CL, Recio AC. Wheelchairs. In: Gonzalez-Fernandez M, Schaaf S, Zumsteg JM, Perret D, Kennedy DJ, editors. Handbook of Physical Medicine and Rehabilitation. Springer Publishing Company; 2022:543-548. [Context Link 1, 2]
82. Hasnan N. Wheelchair and seating systems. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:129-144. [Context Link 1]
83. O'Shea RK, Bonfiglio BS. Assistive technology. In: Palisano RJ, Orlin M, Schreiber J, editors. Campbell's Physical Therapy for Children. 6th ed. Elsevier; 2023:819-844. [Context Link 1]

84. Filson R. Musculoskeletal trauma and orthopedic surgery. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:1638-1672. [Context Link 1, 2]
85. Manual techniques. In: Dutton M, editor. *Dutton's Orthopaedic Examination, Evaluation, and Intervention*. 6th ed. McGraw-Hill Education; 2023:382-404. [Context Link 1, 2]
86. Goldberg JI, Rosa WE. Pain. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:119-145. [Context Link 1]
87. Russek LN. Chronic pain. In: Fulk G, Chui K, editors. *O'Sullivan & Schmitz's Physical Rehabilitation*. 8th ed. USA: F.A. Davis Company; 2024:966-1010. [Context Link 1]
88. Hsieh LF. Rheumatologic rehabilitation. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:297-310. [Context Link 1, 2]
89. Parish LC, Karadag AS, Parish JL. Decubitus (pressure) ulcers. In: Lebowitz MG, Heymann WR, Coulson IH, Murrell DF, editors. *Treatment of Skin Disease*. 6th ed. Elsevier; 2022:185-192. [Context Link 1]
90. Patient safety. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. *Clinical Nursing Skills and Techniques*. 11th ed. Elsevier; 2025:382-416. [Context Link 1]
91. Effgen SK. The educational environment. In: Palisano RJ, Orlin M, Schreiber J, editors. *Campbell's Physical Therapy for Children*. 6th ed. Elsevier; 2023:768-795. [Context Link 1]
92. Tissue behavior, injury, healing, and treatment. In: Dutton M, editor. *Dutton's Orthopaedic Examination, Evaluation, and Intervention*. 6th ed. McGraw-Hill Education; 2023:26-56. [Context Link 1, 2]
93. Hopkins MS, Gonzalez-Fernandez M. Orthoses. In: Gonzalez-Fernandez M, Schaaf S, Zumsteg JM, Perret D, Kennedy DJ, editors. *Handbook of Physical Medicine and Rehabilitation*. Springer Publishing Company; 2022:521-528. [Context Link 1]
94. Wu CH. Physical agent modalities. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:165-172. [Context Link 1]
95. The intervention. In: Dutton M, editor. *Dutton's Orthopaedic Examination, Evaluation, and Intervention*. 6th ed. McGraw-Hill Education; 2023:343-368. [Context Link 1]
96. Costello KJ. Spinal cord and peripheral nerve problems. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:1594-1620. [Context Link 1]
97. Coppola S, Gillen, Boyt Schell BA. Contemporary occupational therapy practice and future directions. In: Gillen G, Brown C, editors. *Willard & Spackman's Occupational Therapy*. 14th ed. Wolters Kluwer Health; 2023:54-70. [Context Link 1]
98. Nilsen DM, Gillen G. Motor function and occupational performance. In: Gillen G, Brown C, editors. *Willard & Spackman's Occupational Therapy*. 14th ed. Wolters Kluwer Health; 2023:911-941. [Context Link 1]
99. Togliola JP, Goverover Y. Cognitive, perception, and occupational performance. In: Gillen G, Brown C, editors. *Willard & Spackman's Occupational Therapy*. 14th ed. Wolters Kluwer Health; 2023:942-991. [Context Link 1]
100. Hock N. Orthopedics. In: Atchison BJ, Dirette DP, editors. *Conditions in Occupational Therapy: Effect on Occupational Performance*. 6th ed. Wolters Kluwer; 2023:551-570. [Context Link 1, 2]
101. Maher C, Mendonca RJ. Orthopedic conditions. In: Dirette DP, Gutman SA, editors. *Occupational Therapy for Physical Dysfunction*. 8th ed. Wolters Kluwer; 2021:857-876. [Context Link 1]
102. Pitonyak JS, Wilbur KL. Activities of daily living and instrumental activities of daily living. In: Gillen G, Brown C, editors. *Willard & Spackman's Occupational Therapy*. 14th ed. Wolters Kluwer Health; 2023:706-743. [Context Link 1]
103. Dobkin BH. Neurological rehabilitation. In: Jankovic J, Mazziotta JC, Pomeroy SL, Newman NJ, editors. *Bradley and Daroff's Neurology in Clinical Practice*. 8th ed. Elsevier; 2022:819-849.e2. [Context Link 1]
104. Van Oss T, Sanders MJ, Hewitt P. Environmental assessment and modification: home, work, and community. In: Dirette DP, Gutman SA, editors. *Occupational Therapy for Physical Dysfunction*. 8th ed. Wolters Kluwer; 2021. [Context Link 1]
105. Perr A. Restoring transfers and functional mobility. In: Dirette DP, Gutman SA, editors. *Occupational Therapy for Physical Dysfunction*. 8th ed. Wolters Kluwer; 2021:641-656. [Context Link 1]
106. Miller AC. Balance assessment intervention. In: Dirette DP, Gutman SA, editors. *Occupational Therapy for Physical Dysfunction*. 8th ed. Wolters Kluwer; 2021. [Context Link 1]

107. Ramiro RPT. Occupational medicine and vocational rehabilitation. In: Cifu DX, Lew HL, editors. *Braddom's Rehabilitation Care: A Clinical Handbook*. 2nd ed. Elsevier; 2025:43-54. [Context Link 1, 2, 3]
108. Bracciano AG. Physical agent modalities and biofeedback. In: Dirette DP, Gutman SA, editors. *Occupational Therapy for Physical Dysfunction*. 8th ed. Wolters Kluwer; 2021:487-510. [Context Link 1]
109. Schultz-Krohn W, Pendleton HM. Application of the occupational therapy practice framework to physical dysfunction. In: Pendleton HM, Schultz-Krohn W, editors. *Pedretti's Occupational Therapy*. 9th ed. Elsevier; 2024:29a-53. [Context Link 1]
110. Daras LC, et al. Evaluating hospital readmission rates after discharge from inpatient rehabilitation. *Archives of Physical Medicine and Rehabilitation* 2018;99(6):1049-1059. DOI: 10.1016/j.apmr.2017.07.008. [Context Link 1] View abstract...
111. Middleton A, Graham JE, Ottenbacher KJ. Functional status is associated with 30-day potentially preventable hospital readmissions after inpatient rehabilitation among aged Medicare Fee-for-Service beneficiaries. *Archives of Physical Medicine and Rehabilitation* 2018;99(6):1067-1076. DOI: 10.1016/j.apmr.2017.05.001. [Context Link 1] View abstract...
112. Osuafor CN, Sahimi SNM, Enduluri S, McCarthy F. Incidence and outcome of interrupted geriatric rehabilitation requiring acute hospital transfer. *Irish Journal of Medical Science* 2019;188(4):1451-1454. DOI: 10.1007/s11845-019-01974-0. [Context Link 1] View abstract...
113. Keim SK, Ratcliffe SJ, Naylor MD, Bowles KH. Patient factors linked with return acute healthcare use in older adults by discharge disposition. *Journal of the American Geriatrics Society* 2020;68(10):2279-2287. DOI: 10.1111/jgs.16645. [Context Link 1] View abstract...
114. Herrmann AA, et al. Potentially preventable readmissions after acute inpatient rehabilitation. *American Journal of Physical Medicine and Rehabilitation* 2023;102(11):1014-1019. DOI: 10.1097/PHM.0000000000002269. [Context Link 1] View abstract...
115. Bidlespacher K, Mulkey DC. The Effect of Teach-Back on Readmission Rates in Rehabilitation Patients. *Rehabilitation Nursing* 2024 Mar-Apr 01;49(2):65-72. DOI: 10.1097/RNJ.0000000000000452. [Context Link 1] View abstract...
116. Philippe AM, Farid R. From inpatient rehabilitation facility to skilled nursing facility to home: A retrospective study of functional recovery. *PM & R: The Journal of Injury, Function, and Rehabilitation* 2024;16(3):226-230. DOI: 10.1002/pmrj.13065. [Context Link 1] View abstract...
117. Program standards. In: CARF, International, editors. *Medical Rehabilitation Standards Manual*. 2025 ed. Tucson, AZ: Commission on Accreditation of Rehabilitation Facilities; 2025:189-272. [Context Link 1]
118. Adult health and physical, nutritional, and cultural assessment. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:71-96. [Context Link 1]
119. The nursing process in drug therapy and patient safety. In: Tucker RG, editor. *Karch's Focus on Nursing Pharmacology*. 9th ed. Wolters Kluwer; 2023:47-57. [Context Link 1]
120. Adverse drug reactions and medication errors. In: Burchum JR, Rosenthal LD, editors. *Lehne's Pharmacology for Nursing Care*. 12th ed. Elsevier; 2025:64-75. [Context Link 1]
121. Health education and health promotion. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. *Brunner & Suddarth's Textbook of Medical-Surgical Nursing*. 15th ed. Wolters Kluwer; 2022:56-70. [Context Link 1, 2]
122. Safe medication preparation. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. *Clinical Nursing Skills and Techniques*. 11th ed. Elsevier; 2025:586-607. [Context Link 1]
123. Angelelli J, et al. Effect of social determinants of health interventions on adults living with disabilities: a scoping review. *Archives of Physical Medicine and Rehabilitation* 2022;103(5):1023-1033.e11. DOI: 10.1016/j.apmr.2021.06.021. [Context Link 1] View abstract...
124. Ezell JM, Hamdi S, Borrero N. Approaches to addressing nonmedical services and care coordination needs for older adults. *Research on Aging* 2022 Mar-Apr;44(3-4):323-333. DOI: 10.1177/01640275211033929. [Context Link 1] View abstract...
125. Martinez-Kratz M. Guide to planning care. In: Makic MB, Martinez-Kratz MR, editors. *Ackley and Ladwig's Nursing Diagnosis Handbook*. 13th ed. Elsevier; 2023:125-1051. [Context Link 1]
126. Harding MM. Substance use disorders in acute care. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. *Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems*. 12th ed. St. Louis, MO: Mosby; 2023:162-178. [Context Link 1]

127. Bassil NK, Ohanian MLK, Bou Saba TG. Nicotine use disorder in older adults. Clinics in Geriatric Medicine 2022;38(1):119-131. DOI: 10.1016/j.cger.2021.07.008. [Context Link 1] View abstract...
 128. Centers for Medicare and Medicaid Services. "Durable medical equipment, prosthetics, orthotics and supplies (DMEPOS): scope and conditions." 42 CFR Pt. 410.38 Washington, DC 2024 Jan 01 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/current/title-42/>. [Context Link 1, 2]
 129. Care and Compare. Find and Compare Nursing Homes, Hospitals, and Other Providers Near You [Internet] Medicare.gov. Accessed at: <https://www.medicare.gov/care-compare/>. [accessed 2025 Apr 02] [Context Link 1, 2]
 130. Centers for Medicare and Medicaid Services. Medicare Program Integrity Manual. Chapter 6 - Medical Review of Inpatient Hospital Claims for Part A Payment Rev. 10365 [Internet] Centers for Medicare and Medicaid Services. 2020 Oct 02 Accessed at: <https://www.cms.gov/medicare/regulations-guidance/manuals/internet-only-manuals-ioms>. [accessed 2025 Sep 09] [Context Link 1]
 131. Medicare Coverage Database. [Internet] Centers for Medicare and Medicaid Services. Accessed at: <https://www.cms.gov/medicare-coverage-database/search.aspx>? Updated 2025 [accessed 2025 Oct 23] [Context Link 1]
-

Footnotes

[A] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) requires specific considerations for admission to this level of care for certain diagnoses, which are outlined in the electronic Code of Federal Regulations part 412.(1)(2) [A in Context Link 1]

[B] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) requires a preadmission screening to be conducted within 48 hours preceding the inpatient rehabilitation admission and is to be completed by a licensed or certified clinician or physician either in person or by telephone through review of the referring hospital medical records.(1) [B in Context Link 1]

[C] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) required elements of the preadmission screening include: level of function prior to the event, condition that caused the need for rehabilitation, expected level of improvement and expected length of time to achieve that level of improvement, risk for clinical complications, combinations of treatments needed (PT, OT, SLP, or orthosis), and anticipated discharge destination.(1) [C in Context Link 1]

[D] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) defines reasonable and necessary skilled care as a service that is "so inherently complex that it can only be safely and effectively performed by, or under the supervision of, professional or technical personnel" based on the documented assessment of the patient's individual care needs. In some cases, a service that would originally be considered unskilled may be considered skilled if the patient's underlying conditions or complications require the involvement of licensed personnel to ensure that essential nonskilled care is achieving its purpose.(5)(6) [D in Context Link 1]

[E] **National guidelines:** According to the Centers for Medicare and Medicaid Services (CMS), the requirement for medical supervision means that a licensed physician who is determined by the inpatient rehabilitation facility to have specialized training and experience in inpatient rehabilitation must conduct face-to-face encounters with the patient at least 3 days per week throughout the patient's stay to assess the patient, as well as to modify the course of treatment as needed to maximize the patient's ability to benefit from rehabilitation. After the first week of physician encounters, CMS will allow a non-physician practitioner to independently conduct one of the three required visits.(3) [E in Context Link 1]

[F] **National guidelines:** According to the Centers for Medicare and Medicaid Services (CMS), the interdisciplinary care team must meet weekly throughout the patient's stay and consist of a registered nurse with specialized training or experience in rehabilitation, a social worker or case manager (or both), a licensed or certified therapist from each therapy discipline involved with the patient's care, and be led by a rehabilitation physician.(3) [F in Context Link 1]

[G] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) uses a data collection system to establish best practice, understand costs, and identify cost-effective systems and procedures. The current data collection system for eligible patients is the Inpatient Rehabilitation Facility Patient Assessment

Instrument (IRF-PAI). IRF-PAI is an assessment tool used to identify patient problems, strengths, and preferences, and to create an individualized care plan.(37) [G in Context Link 1]

[H] **National guidelines:** Starting October 1, 2024, the Centers for Medicare and Medicaid Services (CMS) requires inpatient rehabilitation facilities to use the Inpatient Rehabilitation Facility Patient Assessment Instrument (IRF-PAI) to assess all inpatients, regardless of payer. The admission IRF-PAI should cover calendar days 1 through 3 of the patient's current admission; the patient assessment schedule is further detailed in the electronic Code of Federal Regulations part 412.610.(38)(39) [H in Context Link 1]

[I] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) requires that an individualized overall plan of care for the patient is developed by a rehabilitation physician with input from the interdisciplinary team within 4 days of the patient's admission.(1) [I in Context Link 1]

[J] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) states that therapy treatments should begin within 36 hours of admission.(1)(3) [J in Context Link 1]

[K] **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) recommends that if an unexpected clinical event occurs that limits a patient's ability to participate in an intensive therapy program for a brief period not to exceed 3 days, the specific reasons for the break should be documented in the patient's medical record.(3) [K in Context Link 1]

Definitions

Custodial care

- Custodial care is personal unskilled care to support the patient's care of activities of daily living (ADL), such as bathing, dressing, and eating.(1)

References

1. Mauk KL. Post-acute care. In: Larsen PD, editor. Lubkin's Chronic Illness: Impact and Intervention. 11th ed. Jones & Bartlett Learning; 2022:477-520.

FIM Score

- The Functional Independence Measure (FIM®) is an instrument that classifies a patient's ability to carry out activities and the need for assistance from a person or device. It is a 7-level scale measurement of gradations in behavior (from dependence to independence) that can be used to monitor patient progress throughout rehabilitation and help with care plan development and defining discharge needs.(1)

References

1. Carter WE III, Peretiatko S. Quality and outcome measures for medical rehabilitation. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:55-62.

Goals for measurable improvement

- Goals for measurable improvement include identification of **ALL** of the following(1)(2):
 - Nature and degree of improvement
 - Time expected for achievement
 - Functional improvement anticipated to be practical, ongoing, and sustainable

References

- Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09]
- Pryor J, O'Reilly K, Bonser M, Garret G, McKenchnie D. Rehabilitation for the individual and family. In: Chang E, Johnson A, editors. Living With Chronic Illness and Disability. 4th ed. Chatswood NSW 2067: Elsevier; 2022:161-182.

Intensive rehabilitation therapy program

- Intensive rehabilitation therapy program that includes **1 or more** of the following^{[A](1)(2)}:
 - Therapy at least 3 hours per day for 5 days per week
 - Therapy at least 15 hours per week (in a 7 consecutive calendar day period starting from the date of admission)
 - Other intensive therapy to improve functional capacity at patient's maximum participation level without compromising safety or exceeding ability or tolerance⁽³⁾

References

- Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09]
- Centers for Medicare and Medicaid Services. "Basis of payment." 42 CFR Pt. 412.622 Washington, DC 2025 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>.
- Pryor J, O'Reilly K, Bonser M, Garret G, McKenchnie D. Rehabilitation for the individual and family. In: Chang E, Johnson A, editors. Living With Chronic Illness and Disability. 4th ed. Chatswood NSW 2067: Elsevier; 2022:161-182.

Footnotes

- A. **National guidelines:** The Centers for Medicare and Medicaid Services (CMS) Medicare Benefit Policy Manual states that admissions that are intended to assess if a patient would benefit and tolerate treatment in an inpatient rehabilitation facility (IRF) are no longer considered reasonable and necessary.⁽¹⁾

Interdisciplinary team care

- Interdisciplinary care team meetings held weekly and discipline examples include⁽¹⁾⁽²⁾⁽³⁾⁽⁴⁾:
 - A rehabilitation physician designated by the Inpatient Rehabilitation Facility to have specialized training and experience in inpatient rehabilitation services
 - A registered nurse with specialized training or experience in rehabilitation
 - A social worker or case manager (or both)
 - A licensed or certified therapist from each discipline involved in evaluating and treating the patient

References

- Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09]
- Centers for Medicare and Medicaid Services. "Basis of payment." 42 CFR Pt. 412.622 Washington, DC 2025 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>.
- Pryor J, O'Reilly K, Bonser M, Garret G, McKenchnie D. Rehabilitation for the individual and family. In: Chang E, Johnson A, editors. Living With Chronic Illness and Disability. 4th ed. Chatswood NSW 2067: Elsevier; 2022:161-182.

4. Centers for Medicare and Medicaid Services. Medicare Program Integrity Manual. Chapter 6 - Medical Review of Inpatient Hospital Claims for Part A Payment Rev. 10365 [Internet] Centers for Medicare and Medicaid Services. 2020 Oct 02 Accessed at: <https://www.cms.gov/medicare/regulations-guidance/manuals/internet-only-manuals-ioms>. [accessed 2025 Sep 09]

Occupational therapy (OT) progression

- OT progression includes **ALL** of the following(1)(2):
 - Ongoing evaluation of treatment plan with measurable short-term and long-term goals
 - Documentation of effectiveness and efficiency toward achieving established goals
 - Patient demonstrates resolving of barriers to transition of care and **1 or more** of the following:
 - Objective measurements document progress toward goals in reasonable and predictable time frame based on treatment plan.[A](4)
 - Minimal progress due to unexpected clinical event, but progress expected to resume toward goals within 3 days as condition improves(5)

References

1. Bennett K. Setting rehabilitation goals. In: O'Hanlon S, Smith M, editors. Comprehensive Guide to Rehabilitation of the Older Patient. 4th ed. Elsevier; 2021:49-52.
2. Pryor J, O'Reilly K, Bonser M, Garret G, McKenchnie D. Rehabilitation for the individual and family. In: Chang E, Johnson A, editors. Living With Chronic Illness and Disability. 4th ed. Chatswood NSW 2067: Elsevier; 2022:161-182.
3. Carter WE III, Peretiatko S. Quality and outcome measures for medical rehabilitation. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:55-62.
4. Kee YYK. Measuring progress with rehabilitation. In: O'Hanlon S, Smith M, editors. Comprehensive Guide to Rehabilitation of the Older Patient. 4th ed. Elsevier; 2021:111-115.
5. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09]

Footnotes

- A. Improvements may be measured by accuracy (percentage or number of correct trials), increased number of repetitions, decreased assistance level (maximum, moderate, or minimal assistance), decreased pain level, increased ROM/strength, increased balance scores, decreased level of cues or prompts required (in conjunction with level of assistance), or improvement in standardized functional outcome measures (eg, Continuity Assessment Record and Evaluation (CARE), WeeFIM®, Tinetti, Berg Balance Scale).(3)(4) Generally, it is expected that measured improvements should be demonstrable in targeted areas over a 1-week to 2-week period of skilled therapy. The speed of recovery is variable and may be linked to intensity of treatment, patient condition, age, comorbidities, and other factors.

Physical therapy (PT) progression

- PT progression includes **ALL** of the following(1)(2)(3)(4):
 - Ongoing evaluation of treatment plan with measurable short-term and long-term goals
 - Documentation of effectiveness and efficiency toward achieving established goals
 - Patient demonstrates resolving of barriers to transition of care and **1 or more** of the following:
 - Objective measurements document progress toward goals in reasonable and predictable time frame based on treatment plan.[A]
 - Minimal progress due to unexpected clinical event, but progress expected to resume toward goals within 3 days as condition improves

References

1. Centers for Medicare and Medicaid Services. "Skilled services requirements." 42 CFR Pt. 409.44 Washington, DC 2023 Oct 02 [accessed 2025 Jul 23]
Accessed at: <https://www.ecfr.gov/>.
2. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09]
3. Hussein N. Acute medical conditions. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:253-264.
4. Carter WE III, Peretiatko S. Quality and outcome measures for medical rehabilitation. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:55-62.
5. Scalzitti DA. Examination of function. In: Fulk G, Chui K, editors. O'Sullivan & Schmitz's Physical Rehabilitation. 8th ed. USA: F.A. Davis Company; 2024:270-291.

Footnotes

- A. Improvements may be measured by accuracy (percentage or number of correct trials), increased number of repetitions, decreased assistance level (maximum, moderate, or minimal assistance), decreased pain level, increased ROM/strength, increased balance scores, decreased level of cues or prompts required (in conjunction with level of assistance), or improvement in standardized functional outcome measures (eg, Continuity Assessment Record and Evaluation (CARE), WeeFIM®, Tinetti, Berg Balance Scale). Generally, it is expected that measured improvements should be demonstrable in targeted areas over a 1-week to 2-week period of skilled therapy. The speed of recovery is variable and may be linked to intensity of treatment, patient condition, age, comorbidities, and other factors.(4)(5)

Safe to go home

- Patient safe to go home if **ALL** of the following exist(1)(2)(3):
 - Medical condition manageable at home
 - Functional status manageable at home
 - Mental status stable for patient's condition
 - Medication availability confirmed and reconciliation complete
 - Patient/caregiver education complete and written discharge instructions provided
 - Caregiver and community resources identified (as needed)
 - Community resources identified and referrals made, as needed
 - Home care arranged, if indicated
 - Necessary medical equipment delivery arranged or available in home, if indicated
 - Necessary medical supplies ordered, or patient/caregiver can obtain, if indicated

References

1. Elements of Excellence in Transitions of Care (TOC). TOC Checklist [Internet] National Transitions of Care Coalition. 2006 Accessed at: <https://www.ntocc.org/>. [accessed 2025 Sep 16]
2. Medical-surgical nursing. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:33-55.
3. Admitting, transfer, and discharge. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. Clinical Nursing Skills and Techniques. 11th ed. Elsevier; 2025:40-51.

Skilled care

- Skilled care may be direct (eg, in-person assessment and administration of treatments) or indirect (eg, coordinating care between providers, supervising care aides); examples include(1)(2)(3)(4):
 - Performed or supervised by professional or technical personnel
 - Exceed scope of Custodial care
 - Service is considered skilled as indicated by either of the following:
 - Service that is "so inherently complex that it can only be safely and effectively performed by, or under the supervision of, professional or technical personnel" based on the documented assessment of the patient's individual care needs.
 - Service that would originally be considered unskilled may be considered skilled if the patient's underlying conditions or complications require the involvement of licensed personnel to ensure that essential nonskilled care is achieving its purpose.

References

1. Centers for Medicare and Medicaid Services. "Criteria for skilled services and the need for skilled service." 42 CFR Pt. 409.32 Washington, DC 2023 Oct 02 [accessed 2025 Jul 22] Accessed at: <https://www.ecfr.gov/>.
2. Harding MM. Professional nursing. In: Harding MM, Kwong J, Hagler D, Reinisch C, editors. Lewis's Medical-Surgical Nursing: Assessment and Management of Clinical Problems. 12th ed. St. Louis, MO: Mosby; 2023:1-18.
3. Medical-surgical nursing. In: Hinkle JL, Cheever KH, Overbaugh KJ, editors. Brunner & Suddarth's Textbook of Medical-Surgical Nursing. 15th ed. Wolters Kluwer; 2022:33-55.
4. Centers for Medicare and Medicaid Services. Medicare Program Integrity Manual. Chapter 6 - Medical Review of Inpatient Hospital Claims for Part A Payment Rev. 10365 [Internet] Centers for Medicare and Medicaid Services. 2020 Oct 02 Accessed at: <https://www.cms.gov/medicare/regulations-guidance/manuals/internet-only-manuals-ioms>. [accessed 2025 Sep 09]

Speech-language pathology (SLP) progression

- SLP progression includes **ALL** of the following(1)(2)(3)(4)(5):
 - Ongoing evaluation of treatment plan with measurable short-term and long-term goals
 - Documentation of effectiveness and efficiency toward achieving established goals
 - Patient demonstrates resolving of barriers to transition of care and **1 or more** of the following:
 - Objective measurements document progress toward goals in reasonable and predictable time frame based on treatment plan.[A](6)
 - Minimal progress due to unexpected clinical event, but progress expected to resume toward goals within 3 days as condition improves

References

1. Pryor J, O'Reilly K, Bonser M, Garret G, McKenchnie D. Rehabilitation for the individual and family. In: Chang E, Johnson A, editors. Living With Chronic Illness and Disability. 4th ed. Chatswood NSW 2067: Elsevier; 2022:161-182.
2. The essential ingredients of good therapy: basic skills. In: Roth FP, Worthington CK, editors. Treatment Resource Manual for Speech-Language Pathology. 7th ed. San Diego, CA: Plural Publishing, Inc; 2025:16-38.
3. Shyu SG, Liang HW. The physiatric history and physical examination. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:3-16.
4. Centers for Medicare and Medicaid Services. Medicare Benefit Policy Manual. Chapter 1 - Inpatient Hospital Services Covered Under Part A Rev. 10892 [Internet] Centers for Medicare and Medicaid Services. 2021 Aug Accessed at: <https://www.cms.gov/manuals/>. [accessed 2025 Sep 09]

5. Centers for Medicare and Medicaid Services. "Skilled services requirements." 42 CFR Pt. 409.44 Washington, DC 2023 Oct 02 [accessed 2025 Jul 23]
Accessed at: <https://www.ecfr.gov/>.
6. Carter WE III, Peretiatko S. Quality and outcome measures for medical rehabilitation. In: Cifu DX, Lew HL, editors. Braddom's Rehabilitation Care: A Clinical Handbook. 2nd ed. Elsevier; 2025:55-62.
7. National Outcomes Measurement System (NOMS). [Internet] American Speech-Language-Hearing Association. Accessed at: <https://www.asha.org/noms/>.
Updated 2025 [accessed 2025 Feb 11]

Footnotes

- A. Improvements may be measured by accuracy (percentage or number of correct trials), increased number of repetitions, decreased assistance level (maximum, moderate, or minimal assistance), decreased pain level, decreased level of cues or prompts required (in conjunction with level of assistance), or improvement in standardized functional outcome measures (eg, Continuity Assessment Record and Evaluation (CARE), WeeFIM®, National Outcomes Measurement System (NOMS)). Generally, it is expected that measured improvements should be demonstrable in targeted areas over a 1-week to 2-week period of skilled therapy. The speed of recovery is variable and may be linked to intensity of treatment, patient condition, age, comorbidities, and other factors.(2)(6)(7)

Transition of care planning completed

- Transition of care planning completed; including **ALL** of the following(1)(2):
 - Transition plan communicated to all members of patient's healthcare team
 - Summary of inpatient rehabilitation facility admission, current list of medications, and discharge plan communicated to primary care provider at next level of care
 - Medication reconciliation complete
 - Education on condition management and complications to report provided to patient or caregiver
 - Follow-up appointments scheduled
 - Referrals to continue rehabilitation goals (eg, outpatient or home therapy) arranged, if needed
 - Services (eg, equipment, environment modifications, transportation) required for transition to next level of care arranged, if needed
 - Psychosocial issues addressed, or plan for management at next level of care arranged, if needed

References

1. Transition of care. In: Gillingham DC, editor. Foundations of Case Management. Blue Bayou Press; 2021:137-144.
2. Admitting, transfer, and discharge. In: Perry AG, Potter PA, Ostendorf WR, Laplante N, editors. Clinical Nursing Skills and Techniques. 11th ed. Elsevier; 2025:40-51.

Codes

ICD-10 Diagnosis: G72.0, G72.9, G91.2, G95.9, M48.02, M48.54XD, M48.56XD, M51.16, M54.16, M80.08XD, S12.100D, S22.080D, S22.089D, S22.41XD, S22.42XD, S22.49XD, S32.010D, S32.019D, S32.029D, S32.10XD, S32.501D, S32.502D, S32.509D, S32.511D, S32.512D, S32.519D, S32.591D, S32.592D, S32.599D, S32.810D, S32.82XD, S42.201D, S42.202D, S42.209D, S72.401D, S72.402D, S72.409D, S82.141D, S82.142D, S82.143D, S82.841D, S82.842D, S82.843D, Z47.81 [Hide]

Copyright © 2026 MCG Health, LLC
All Rights Reserved

Last Update: 1/25/2026 12:56:47 AM
Build Number: 30.0.2026012500524.025256